

A11100 988630


NBS
PUBLICATIONS

NAT'L INST OF STANDARDS & TECH R.I.C.



A11100988630

/NBS monograph
QC100 .U556 V25-13;1976 C.1 NBS-PUB-C 19



NBS MONOGRAPH 25 — SECTION 13

U.S. DEPARTMENT OF COMMERCE / National Bureau of Standards

Standard X-ray Diffraction Powder Patterns

QC
100
U556
No.25-13
1976
c.2

NATIONAL BUREAU OF STANDARDS

The National Bureau of Standards¹ was established by an act of Congress March 3, 1901. The Bureau's overall goal is to strengthen and advance the Nation's science and technology and facilitate their effective application for public benefit. To this end, the Bureau conducts research and provides: (1) a basis for the Nation's physical measurement system, (2) scientific and technological services for industry and government, (3) a technical basis for equity in trade, and (4) technical services to promote public safety. The Bureau consists of the Institute for Basic Standards, the Institute for Materials Research, the Institute for Applied Technology, the Institute for Computer Sciences and Technology, and the Office for Information Programs.

THE INSTITUTE FOR BASIC STANDARDS provides the central basis within the United States of a complete and consistent system of physical measurement; coordinates that system with measurement systems of other nations; and furnishes essential services leading to accurate and uniform physical measurements throughout the Nation's scientific community, industry, and commerce. The Institute consists of the Office of Measurement Services, the Office of Radiation Measurement and the following Center and divisions:

Applied Mathematics — Electricity — Mechanics — Heat — Optical Physics — Center for Radiation Research: Nuclear Sciences; Applied Radiation — Laboratory Astrophysics² — Cryogenics² — Electromagnetics² — Time and Frequency².

THE INSTITUTE FOR MATERIALS RESEARCH conducts materials research leading to improved methods of measurement, standards, and data on the properties of well-characterized materials needed by industry, commerce, educational institutions, and Government; provides advisory and research services to other Government agencies; and develops, produces, and distributes standard reference materials. The Institute consists of the Office of Standard Reference Materials, the Office of Air and Water Measurement, and the following divisions:

Analytical Chemistry — Polymers — Metallurgy — Inorganic Materials — Reactor Radiation — Physical Chemistry.

THE INSTITUTE FOR APPLIED TECHNOLOGY provides technical services to promote the use of available technology and to facilitate technological innovation in industry and Government; cooperates with public and private organizations leading to the development of technological standards (including mandatory safety standards), codes and methods of test; and provides technical advice and services to Government agencies upon request. The Institute consists of the following divisions and Centers:

Standards Application and Analysis — Electronic Technology — Center for Consumer Product Technology: Product Systems Analysis; Product Engineering — Center for Building Technology: Structures, Materials, and Life Safety; Building Environment; Technical Evaluation and Application — Center for Fire Research: Fire Science; Fire Safety Engineering.

THE INSTITUTE FOR COMPUTER SCIENCES AND TECHNOLOGY conducts research and provides technical services designed to aid Government agencies in improving cost effectiveness in the conduct of their programs through the selection, acquisition, and effective utilization of automatic data processing equipment; and serves as the principal focus within the executive branch for the development of Federal standards for automatic data processing equipment, techniques, and computer languages. The Institute consists of the following divisions:

Computer Services — Systems and Software — Computer Systems Engineering — Information Technology.

THE OFFICE FOR INFORMATION PROGRAMS promotes optimum dissemination and accessibility of scientific information generated within NBS and other agencies of the Federal Government; promotes the development of the National Standard Reference Data System and a system of information analysis centers dealing with the broader aspects of the National Measurement System; provides appropriate services to ensure that the NBS staff has optimum accessibility to the scientific information of the world. The Office consists of the following organizational units:

Office of Standard Reference Data — Office of Information Activities — Office of Technical Publications — Library — Office of International Relations — Office of International Standards.

¹ Headquarters and Laboratories at Gaithersburg, Maryland, unless otherwise noted; mailing address Washington, D.C. 20234.

² Located at Boulder, Colorado 80302.

U.S. Bureau of Standards
OCT 7 1977
rotacc


Standard X-ray Diffraction Powder Patterns

Section 13—Data for 58 Substances

Monograph no. 25-13

Marlene C. Morris, Howard F. McMurdie, Eloise H. Evans,
Boris Paretzkin, Johan H. de Groot,
Camden R. Hubbard and Simon J. Carmel

Institute for Materials Research
National Bureau of Standards
Washington, D.C. 20234



U.S. DEPARTMENT OF COMMERCE, Elliot L. Richardson, *Secretary*

Dr. Betsy Ancker-Johnson, *Assistant Secretary for Science and Technology*

NATIONAL BUREAU OF STANDARDS, Ernest Ambler, *Acting Director*

Issued June 1976

Library of Congress Catalog Card Number: 53-61386

National Bureau of Standards Monograph 25

Section 13—Data for 58 Substances

Nat. Bur. Stand. (U.S.), Monogr. 25—Sec. 13, 114 pages (June 1976)

CODEN: NBSMA6

U.S. GOVERNMENT PRINTING OFFICE

WASHINGTON: 1976

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402
(Order by SD Catalog No. C13.44:25/Sec. 13). Price \$1.80 (Add 25 percent additional for other than U.S. mailing).

CONTENTS

	Page		Page
Introduction.....	1	Cobalt gallium tantalum, Co_2GaTa ...	76
Experimental patterns:		Cobalt gallium titanium, Co_2GaTi ...	77
Arsenic iodide, AsI_3	7	Cobalt gallium vanadium, Co_2GaV	78
Barium silicate, $\beta\text{-BaSiO}_3$	8	Cobalt germanium manganese,	
Barium silicate, (sanbornite),		Co_2GeMn	79
$\beta\text{-BaSi}_2\text{O}_5$	10	Cobalt germanium titanium, Co_2GeTi .	80
Barium silicate, Ba_2SiO_4	12	Cobalt indium, CoIn_3	81
Barium silicate, $\text{Ba}_2\text{Si}_3\text{O}_8$	13	Cobalt lanthanum, CoLa_3	83
Barium silicate, Ba_3SiO_5	15	Cobalt lutetium, Co_2Lu	86
Barium silicate, $\text{Ba}_3\text{Si}_5\text{O}_{13}$	17	Cobalt neodymium, Co_2Nd	87
Cadmium silicate, Cd_2SiO_4	19	Cobalt nickel tin,	
Cadmium silicate, Cd_3SiO_5	20	$\text{Co}_{.75}\text{Ni}_{.75}\text{Sn}_{.75}$	88
Calcium hydrogen phosphate hydrate,		Cobalt samarium, Co_5Sm	90
$\text{Ca}_8\text{H}_2(\text{PO}_4)_6 \cdot 5\text{H}_2\text{O}$	21	Cobalt tin, Co_3Sn_2	92
Cobalt phosphate, $\text{Co}(\text{PO}_3)_2$	23	Cumulative indices	
Copper imidazole nitrate,		(Circular 539, Volumes 1-10 and	
$\text{Cu}(\text{C}_3\text{H}_4\text{N}_2)_4(\text{NO}_3)_2$	24	Monograph 25, Sections 1-13 inclusive)	
Lead chloride fluoride(matlockite),		1. Inorganic.....	94
PbClF	25	2. Organic.....	107
Magnesium phosphate, $\text{Mg}(\text{PO}_3)_2$	26	3. Mineral.....	108
Magnesium tungsten oxide, MgWO_4	27		
Mercury chloride, HgCl_2	29		
Mercury chloride (calomel), Hg_2Cl_2 .	30		
Nickel acetate hydrate,			
$\text{Ni}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 4\text{H}_2\text{O}$	31		
Potassium lead chloride, KPb_2Cl_5 ...	33		
Potassium platinum chloride,			
K_2PtCl_6	34		
Silicon, Si	35		
Silver carbonate, Ag_2CO_3	36		
Silver sulfate, Ag_2SO_4	37		
Sodium phosphate hydrate,			
$\alpha\text{-Na}_4\text{P}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$	39		
Strontium hydroxide, $\text{Sr}(\text{OH})_2$	41		
Strontium hydroxide hydrate,			
$\text{Sr}(\text{OH})_2 \cdot \text{H}_2\text{O}$	42		
Strontium hydroxide hydrate,			
$\text{Sr}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$	43		
Strontium silicate, Sr_3SiO_5	44		
Tin hydrogen phosphate, SnHPO_4	46		
Zinc borate, $\text{Zn}_4\text{B}_6\text{O}_{13}$	48		
Zinc titanium oxide, ZnTiO_3	49		
Calculated patterns:			
Cerium cobalt, CeCo_2	50		
Cerium cobalt, $\text{Ce}_{24}\text{Co}_{11}$	51		
Cerium gallium, CeGa_2	54		
Cerium magnesium, CeMg_3	56		
Cerium nickel, CeNi_2	58		
Cerium thallium, CeTl	59		
Cerium thallium, CeTl_3	60		
Cerium thallium, Ce_3Tl	61		
Cobalt dysprosium, Co_2Dy	63		
Cobalt erbium, Co_2Er	64		
Cobalt erbium, Co_7Er_2	65		
Cobalt gadolinium, CoGd_3	68		
Cobalt gadolinium, Co_2Gd	71		
Cobalt gadolinium, Co_7Gd_2	72		
Cobalt gallium manganese, Co_2GaMn ..	75		

STANDARD X-RAY DIFFRACTION POWDER PATTERNS

The following copies may be obtained from the National Technical Information Service, 5285 Port Royal Road, Springfield, Virginia, 22161. Where these publications are identified with a number, it must be used in ordering. They are available in hardcopy or microfiche; the price is not fixed and will be furnished on request.

Also, until the present supply is exhausted, the publication numbers marked by an asterisk are for sale from the Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402. Order respectively by SD catalog No. C13.44:25/Sec. 11, Price \$1.55 or SD catalog No. C13.44:25/Sec. 12, Price \$1.50.

NBS Publication	Number	NBS Publication	Number
Circular 539, Volume 1.....	PB 178 902	Monograph 25, Section 1.....	PB 178 429
Volume 2.....	PB 178 903	Section 2.....	PB 178 430
Volume 3.....	PB 178 904	Section 3.....	PB 178 431
Volume 4.....	PB 178 905	Section 4	
Volume 5.....	PB 178 906	Section 5	
Volume 6.....	PB 178 907	Section 6	
Volume 7.....	PB 178 908	Section 7	
Volume 8.....	PB 178 909	Section 8.....	PB 194 872
Volume 9.....	PB 178 910	Section 9.....	COM 72-50002
Volume 10.....	PB 178 911	Section 10.....	COM 72-51079
		*Section 11.....	COM 74-50183
		*Section 12.....	COM 75-50162

STANDARD X-RAY DIFFRACTION POWDER PATTERNS

Section 13. --- Data for 58 Substances

by

Marlene C. Morris, Howard F. McMurdie, Eloise H. Evans,
Boris Paretzkin, and Johan H. de Groot
Joint Committee on Powder Diffraction Standards

and

Camden R. Hubbard and Simon J. Carmel
National Bureau of Standards

Standard x-ray diffraction patterns are presented for 58 substances. Thirty-one of these patterns represent experimental data and 27 are calculated. The experimental x-ray powder diffraction patterns were obtained with an x-ray diffractometer. All d-values were assigned Miller indices determined by comparison with computed interplanar spacings consistent with space group extinctions. The densities and lattice constants were calculated and the refractive indices were measured whenever possible. The calculated x-ray powder diffraction patterns were computed from published crystal structure data. Both peak height and integrated intensities are reported for the calculated patterns.

Key words: Crystal structure; integrated intensities; lattice constants; peak intensities; powder patterns; reference intensities; standard; x-ray diffraction.

INTRODUCTION

The Powder Diffraction File is a continuing compilation of diffraction patterns gathered from many sources. Produced and published by the Joint Committee on Powder Diffraction Standards,¹ the File is used for identification of crystalline materials by matching d-spacings and diffraction intensity measurements. Under the partial sponsorship of the Joint Committee, the program at the National Bureau of Standards contributes new data to this File. Our work also aids in the evaluation and revision of published x-ray data and in the development of diffraction techniques. This report presents information for 58 compounds (31 experimental and 27 calculated patterns), and is the twenty-third of the series of "Standard X-ray Diffraction Powder Patterns."²

¹Joint Committee on Powder Diffraction Standards, 1601 Park Lane, Swarthmore, PA. 19081. This Pennsylvania non-profit corporation functions in cooperation with the American Ceramic Society, the American Crystallographic Association, the American Society for Testing and Materials, The Clay Minerals Society, The Institute of Physics, the Mineralogical Association of Canada, the Mineralogical Society of America, The Mineralogical Society of Great Britain and Ireland, the National Association of Corrosion Engineers, and the Société Française de Minéralogie et de Cristallographie.

²See previous page for other published volumes.

EXPERIMENTAL POWDER PATTERNS

Sample. The samples used to make NBS patterns were obtained from a variety of sources or were prepared in small quantities in our laboratory. Appropriate annealing or recrystallization of the sample improved the quality of most of the patterns. A check of phase purity was provided by indexing the x-ray pattern. Unless otherwise noted, the spectrographic analyses were done at NBS after preparation of the sample was completed; the limit of detection for the alkali elements was 0.05 weight percent.

Optical data, color. A microscopic inspection for phase purity was also made on the non-opaque materials during the refractive index determination. The latter was done by grain-immersion methods in white light, using oils standardized in sodium light, in the refractive index range 1.40 to 2.1 [Hartshorne and Stuart, 1970].

The names of the sample colors were selected from the ISCC-NBS Centroid Color Charts [1965].

Interplanar spacings. For spacing determinations, a shallow holder was packed with a sample mixed with an internal standard (approximately 5 wt. percent tungsten powder). If tungsten lines were found to interfere with sample lines, silver or silicon was used in place of tungsten. If the internal standard correction varied along the length of the pattern, linear interpolations were used. To avoid errors associated with aberrations at the very top of peaks, the readings of 2θ were taken at positions about 20 percent of the way down from the top, and in the center of the peak width. The internal standard correction for each region was then applied to the measured value of 2θ . We have reported all data as $K\alpha_1$ peaks because the internal standard corrections for all regions were established in terms of the $K\alpha_1$ wavelength.

The internal standards used were of high purity (99.99%). The lattice constants used for them at 25 °C are given in the table below; the 2 θ angles were computed using cell dimensions uncorrected for index of refraction.

Calculated 2 θ Angles, CuK α_1 λ = 1.540598Å			
hkl	W a=3.16524Å ±.00004	Ag a=4.08651Å ±.00002	Si a=5.43088Å ±.00004
110	40.262		
111		38.112	28.443
200	58.251	44.295	
211	73.184		
220	86.996	64.437	47.303
310	100.632		
311		77.390	56.123
222	114.923	81.533	
321	131.171		
400	153.535	97.875	69.131
331		110.499	76.377
420		114.914	
422		134.871	88.032
511/333		156.737	94.954
440			106.710
531			114.094
620			127.547
533			136.897
444			158.638

The new internal standard Si powder is available as Standard Reference Material 640 [1974]. The lattice constant for the Si was refined from multiple powder data measurements made with tungsten as an internal standard [Swanson et al., 1966]. Cell parameter data were also collected for a single crystal from the boules ground to prepare the powder. The lattice parameters from the two methods agreed within 3 parts in 10⁵ [Hubbard et al. 1975]. D-spacing results using SRM 640 will be in agreement with patterns recorded in this series of monographs since 1966.

All of our spacing measurements were recorded at 25 ± 1 °C on a diffractometer equipped with a focusing graphite or lithium fluoride crystal monochromator located between the sample and the scintillation counter. Pulse height discrimination was used as well. All measurements were performed using copper radiation: λ (CuK α_1 , peak)=1.540598Å [Deslattes and Henins, 1973].

Structure, lattice constants. The space groups were listed with short Hermann-Mauguin symbols as well as the space group numbers given in the *International Tables for X-ray Crystallography*, Vol. I [1952].


Orthorhombic cell dimensions were arranged according to the Dana convention $b > a > c$ [Palache et al., 1944]. Monoclinic and triclinic lattice constants were transformed if necessary in order to follow the convention of *Crystal Data* [1973]. For primitive cells, the transformed cell axes are an alternate labelling of the reduced cell


axes. For centered monoclinic cells, the transformed cell is the centered cell with the three shortest non-coplanar vectors.

A computer program [Evans et al., 1963] assigned hkl's and refined the lattice constants. Cell refinement was based only upon 2 θ_{obs} values which could be indexed without ambiguity. The program minimized the value $\Sigma(\theta_{obs}-\theta_{calc})^2$. The estimated standard deviations (e.s.d.'s) of the reciprocal cell parameters were determined from the inverse matrix of the normal equations. The program calculated the e.s.d.'s of the direct cell constants by the method of propagation of errors. Since 1973, the e.s.d.'s derived by the computer program have been increased by 50% in order to reflect more truly the uncertainty in the lattice constants. A similar increase should also be applied to all lattice constants in earlier publications of this series. In indexing cubic patterns, multiple hkl's were not utilized in the refinement or reported. Instead, the single appropriate index having the largest h was listed. The number of significant figures reported for d-values varied with the symmetry and crystallinity of each sample.

Densities. These were calculated from the specified lattice constants, the Avogadro number 6.0220943×10^{23} [Deslattes et al., 1974] and atomic weights based on carbon 12 [International Union, 1961].

Intensity measurements. It was found that samples which gave satisfactory intensity patterns usually had an average particle size smaller than 10 μ m, as recommended by Alexander et al. [1948]. In order to avoid the orientation effects which occur when powdered samples are packed or pressed, a sample holder was made that had in its top face a rectangular cavity which extended to one end of the holder. To prepare the sample, a glass slide was clamped over the top face to form a temporary cavity wall (see Figure 1), and the powdered sample was allowed to drift into the end opening while the holder was held in a vertical





position. With the sample holder returned to a horizontal position, the glass slide was carefully removed so that the sample could be exposed to the x-ray beam (as shown in Figure 2). If the sample powder did not flow readily, or was prone to orient excessively, approximately 50 volume percent of finely ground silica-gel was added as a diluent. The intensities of the diffraction lines were measured as peak heights above background and were expressed in percentages of the strongest line. At least three patterns for intensity measurements were prepared for each sample to check reproducibility.

Reference Intensity Ratio, I/I_{corundum} . For reference intensity measurements, $\alpha\text{-Al}_2\text{O}_3$ (corundum) was chosen as an internal standard to be mixed 1:1 by weight with the sample. This mixture of two components was mounted in our regular intensity sample holder (see Figures 1 & 2), and the pattern was taken. The reference intensity was then calculated as the direct ratio of the strongest line of the sample to the strongest line of corundum (hexagonal reflection (113)). In a few instances, the strongest line of one of the components coincided with a line of the other. In that case, the second strongest line was measured, and the value for the strongest line was then calculated.

CALCULATED POWDER PATTERNS

Since some substances of interest are not readily available for experimental work, powder patterns were calculated from published crystal structure data. The FORTRAN program used for the computations was developed by Clark, Smith and Johnson [1973] and modified at NBS.

Lattice parameters. Before the computations of the patterns, any necessary changes were made in the lattice constants in order to make them consistent with the revised value of $\lambda(\text{CuK}\alpha_1) = 1.540598\text{\AA}$ [Deslattes and Henins, 1973]. Both the altered and the original published values are given. Monoclinic and triclinic lattice constants

were transformed if necessary, to follow the convention of *Crystal Data* [1973]. For primitive cells, the transformed cell axes are an alternate labelling of the reduced cell axes. For centered monoclinic cells, the transformed cell is the centered cell with the three shortest non-coplanar vectors.

Scattering factors. Whenever possible, the same scattering factors were used which the author of the reference article specified. Otherwise, the factors were taken directly from the *International Tables for X-ray Crystallography*, Vol. III, [1962]. The factors were corrected for dispersion if the author had done so.

Thermal parameters. The computer program used thermal parameter data of only two forms, the isotropic B's or the anisotropic β_{ij} 's in the following expressions:

$$e^{(-B \sin^2\theta)/\lambda^2}$$

or

$$e^{-(h^2\beta_{11}+k^2\beta_{22}+l^2\beta_{33}+2hk\beta_{12}+2hl\beta_{13}+2kl\beta_{23})}.$$

Other thermal parameters were converted to one of these two forms. The isotropic parameters were used directly, if given by the structure reference. In a few of our patterns, anisotropic parameters were also used directly as given by the structure reference; in other work, instead of using given anisotropic parameters, approximately equivalent isotropic values were substituted as defined by:

$$B = 4 \left[\frac{\beta_{11}\beta_{22}\beta_{33}}{a^2 b^2 c^2} \right]^{1/3}$$

Structural information. The atom positions used in these calculated patterns varied somewhat in the degree of reliability. When the expression "the structure was determined by..." was used, the atomic parameters in the reference cited had been calculated from refinement of single crystal data. When only the space group and structure type were given, the atomic positions had been derived by analogy with similar compounds whose structure was known. In cases where isostructural relationships were used, the atoms were in fixed special positions or the ionic radii were closely related to the corresponding radii of the atoms in the known structure.

Integrated intensities. The theoretical integrated intensity of reflection i on the "absolute/relative" scale is computed from the right hand side of the equation:

$$\frac{I_i^{\text{abs}}}{K} = \frac{M_i L_p_i |F_i T_i|^2}{2\mu V^2}$$

where:

F is the structure factor
T is the thermal correction

$L_p = \frac{1+\cos^2 2\theta}{\sin^2 \theta \cos \theta}$ is the Lorentz-polarization term

M is the multiplicity for the reflection i
 μ is the linear absorption coefficient
V is the volume of the unit cell

When the largest integrated intensity was assigned a relative value of 100 and all other reflections were scaled relative to it, the intensities were placed on the relative intensity scale (I^{rel}). Relative intensities were rounded to the nearest integer value before being listed, and reflections with I^{rel} less than 0.7 were omitted.

Scale factor (integrated intensities). The scale factor, γ , was defined to convert the tabulated I^{rel} to the "absolute/relative" scale [Hubbard, Evans and Smith]. That is:

$$\gamma = \frac{M' L_p' |F' T'|^2}{200 \mu V^2}$$

and

$$\frac{I^{abs}}{K} = \gamma I^{rel}$$

The primes denoted the values for the largest integrated intensity. In earlier Monographs (1969-1975), a different scale factor, k_{NBS} , was reported which is related to γ :

$$\frac{\gamma}{k_{NBS}} = \frac{1}{2 \mu V^2}$$

From γ , the theoretical value of the Reference Intensity Ratio, I/I_c , was calculated:

$$I/I_c = \frac{\mu \gamma \rho_c}{\mu_c \gamma_c \rho}$$

where ρ is the density and the subscript c represents corundum (α - Al_2O_3).

Peak intensities. The purpose of calculating peak intensities was to provide a tabulated pattern similar to what might be obtained from experimental diffractometer measurements. For each predicted reflection, Cauchy profiles centered at both the α_1 and the α_2 peak positions were calculated and summed, forming a simulated

powder pattern. The full width at half-maximum (FWHM) was allowed to vary to represent the changing FWHM as a function of 2θ . [The values of the FWHM vs 2θ are given in the table below]. The resultant simulated powder pattern was then analyzed for peaks. In the regions of the predicted reflections several reflections could have identical or similar 2θ angles and produce only one composite peak in the simulated pattern. The 2θ angle of the composite peak was assigned the hkl of the reflection having the greatest contribution to the peak intensity. If any other peak contributed more than 10% of the intensity toward the composite peak intensity, a plus sign (+) was appended to the hkl . Peaks due solely to α_2 lines were omitted. If an α_1 peak and an α_2 peak overlapped, the α_1 reflection was listed only when it contributed a significant intensity (>10%) at the peak 2θ .

The peak search routine located peaks only at 2θ angles which were a multiple of 0.02° .

2θ CuK α_1	FWHM	2θ CuK α_1	FWHM
0°	0.12°	140	0.230
20	.12	145	.255
40	.12	150	.285
60	.125	155	.315
80	.130	160	.360
100	.135	162.5	.410
120	.155	165	.500
130	.185		

REFERENCES

- Alexander, L., Klug, H. P. and Kummer, E. (1948). J. Appl. Phys., 19, No. 8, 742.
- Clark, C.M., Smith, D.K., and Johnson, G.G., Jr. (1973). A FORTRAN IV Program for Calculating X-ray Powder Diffraction Patterns, Version 5, Dept. of Geosciences, Pennsylvania State Univ., PA 16802.
- Crystal Data (1973). (3rd. Ed. Published jointly by the U. S. Department of Commerce, National Bureau of Standards, Washington, D. C. 20234, and the Joint Committee on Powder Diffraction Standards, Swarthmore, PA 19081).
- Deslattes, R.D. and Henins, A. (1973). Phys. Rev. Lett. 31, 972.
- Deslattes, R. D., Henins, A., Bowman, H. A., Schoonover, R.M., Carroll, C.L., Barnes, I.L., Machlan, L.A., Moore, L. J., and Shields, W.R. (1974). Phys. Rev. Lett. 33, 463.
- Evans, H.T., Jr., Appleman, D. E. and Handwerker, D.S. (1963), Report #PB 216188, U. S. Dept. of Commerce, National Technical Information Center 5285 Port Royal Rd., Springfield, VA 22161, \$3.50.
- Hartshorne, N.H. and Stuart, A. (1970). *Crystals and the Polarizing Microscope* (Edward Arnold and Co., London, 4th Ed.)
- Hubbard, C. R., Evans, E. H., and Smith, D. K., J. Appl. Cryst. (to be published).

Hubbard, C. R., Swanson, H. E., and Mauer, F. A.
(1975). *J. Appl. Cryst.* 8, 45.
International Tables for X-ray Crystallography, I
(1952). (The Kynoch Press, Birmingham, Eng.),
Ibid. III (1962). Pgs. 202, 210, 213, 214.
International Union of Pure and Applied Chemistry
(1961). *Chem. Eng. News* 39 (4), Nov. 20, 42.
ISCC-NBS Centroid Color Charts, SRM 2106 (1965),
obtainable from the Office of Standard Refer-
ence Materials, Room B311, Chemistry Building,
National Bureau of Standards, Washington, D.C.
20234. \$5.00 per set.

Palache, C., Berman, H. and Frondel, C. (1944).
Dana's System of Mineralogy (John Wiley and
Sons, New York, 7th Ed.), I, 6.
Standard Reference Material 640, Silicon Powder,
X-ray Diffraction Standard, obtainable from
the Office of Standard Reference Materials,
Room B311, Chemistry Building, National Bureau
of Standards, Washington, D.C. 20234. \$52.00
per 10 gram unit.
Swanson, H. E., Morris, M. C., and Evans, E. H.
(1966). *Nat'l Bur. Std. U.S. Monograph* 25,
Sec. 4, 3.

Arsenic iodide, AsI₃

Sample

The sample was obtained from the City Chemical Company of New York.

Major impurities

0.001 to 0.01% each Bi, Sb
0.0001 to 0.001% each Fe, Si

Color

Bright orange

Structure

Hexagonal, $R\bar{3}(148)$, $Z = 6$ [Braekken, 1930].

NBS lattice constants of this sample:

$$a = 7.2093(8) \text{ \AA}$$

$$c = 21.449(3)$$

Density

(calculated) 4.702 g/cm³

Reference intensity

$I/I_{\text{corundum}} = 1.3$

Additional patterns

1. PDF card 7-272 [Swanson et al., 1956].
2. Hanawalt et al. [1938].
3. Heyworth [1931].

References

- Braekken, H. (1930). Z. Kristallogr. 74, 67.
Hanawalt, J. D., Rinn, H. W., and Frevel, L. K. (1938). Ind. Eng. Chem. Anal. Ed. 10, 457.
Heyworth, D. (1931). Phys. Rev. 38, 351.
Swanson, H. E., Gilfrich, N. T., and Cook, M. I. (1956). Nat. Bur. Stand. (U.S.) Circ. 539, 6, 17.

$\text{CuK}\alpha_1 \lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1 \text{ }^\circ\text{C}$

Internal standard W, $a = 3.16524 \text{ \AA}$

$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
7.14	4	003	12.39
5.39	9	012	16.43
3.573	50	006	24.90
3.218	100	113	27.70
2.752	2	107	32.51
2.539	20	116	35.32
2.464	2	018	36.43
2.384	1	009	37.71
2.081	25	300	43.46
2.028	3	1•0•10	44.65
1.9882	20	119	45.59
1.8611	2	0•1•11	48.90
1.7982	16	306	50.73
1.7873	8	0•0•12	51.06
1.7680	2	0•2•10	51.66
1.7472	10	223	52.32
1.6097	5	226	57.18
1.6012	6	1•1•12	57.51
1.4378	4	229	64.79
1.4295	3	0•0•15,321	65.21
1.3560	4	3•0•12	69.23
1.3388	6	413	70.25
1.3291	5	1•1•15	70.84
1.2730	3	416	74.47
1.2693	3	2•2•12	74.73
1.1914	3	0•0•18,3•2•10	80.56
1.1825	4	419	81.30
1.0337	1	2•4•10	96.35

Barium silicate, β -BaSiO₃

Sample

The sample was prepared by repeated grindings and heatings at about 1100 °C of a 1:1 molar mixture of Ba(OH)₂ and silica gel.

Color

Colorless

Structure

Orthorhombic, Pmmm(47), Z=4, isostructural with BaGeO₃ and NH₄BeF₃ [Liebau, 1957; Toropov and Grebenshchikov, 1956].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 5.6182(5) \text{ \AA} \\ b &= 12.445(1) \\ c &= 4.5816(5) \end{aligned}$$

Density

(calculated) 4.421 g/cm³

Reference intensity

I/I_{corundum} = 2.6

Polymorphism

Funk [1958] reports a second form below about 990°. Grebenshchikov et al. [1967] confirm this and report that the transformation is irreversible. They also suggest a third form (β').

Additional patterns

1. PDF card 6-247 [Levin and Ugrinic, 1953].
2. PDF card 12-651 [Funk, 1958].
3. PDF card 21-83 [Grebenshchikov et al., 1967].
4. Austin [1947].
5. Toropov and Grebenshchikov [1956].

References

- Austin, A. E. (1947). J. Amer. Ceram. Soc. 30, 218.
- Funk, H. (1958). Z. Anorg. Allg. Chem. 296, 46.
- Grebenshchikov, R.G., Shitova, V.I., and Toropov, N.A. (1967). Inorg. Mater. (USSR) 3, 1410.
- Levin, E. M. and Ugrinic, G. M. (1953). J. Res. Nat. Bur. Stand. 51, 37.
- Liebau, F. (1957). Acta. Crystallogr. 10, 290.
- Toropov, N. A. and Grebenshchikov, R. G. (1956). J. Inorg. Chem. (USSR) 1, [12], 41.

CuK α_1 λ = 1.540598 Å; temp. 25±1 °C			
Internal standard Si, a = 5.43088 Å			
d (Å)	I	hkl	2 θ (°)
6.23	2	020	14.20
5.12	14	110	17.30
4.174	11	120	21.27
3.693	35	021	24.08
3.552	15	101	25.05
3.418	100	111	26.05
3.339	65	130	26.68
3.112	55	040	28.66
2.808	25	200	31.84
2.740	10	210	32.65
2.723	10	140	32.86
2.699	14	131	33.17
2.574	8	041	34.82
2.353	17	211	38.22
2.342	11	141	38.41
2.325	4	230	38.69
2.293	17	002	39.26
2.235	25	221	40.32
2.186	2	051	41.26
2.123	<1	102	42.55
2.085	12	240	43.37
2.075	20	231,060	43.59
2.039	30	151	44.39
2.007	5	032	45.13
1.946	2	160	46.63
1.897	8	241	47.91
1.889	20	132,061	48.12
1.8629	4	250	48.85
1.8459	10	042	49.33
1.7932	6	320	50.88
1.7766	6	070,202	51.39
1.7588	5	212	51.95
1.7541	4	142	52.10
1.7337	7	301	52.76
1.7174	4	311	53.30
1.7067	5	330	53.66
1.6950	15	170	54.06
1.6685	2	260	54.99
1.6328	2	232	56.30
1.6151	1	152	56.97
1.6046	4	340	57.38
1.6002	4	331	57.55
1.5682	4	261	58.84
1.5554	1	080	59.37
1.5427	4	242	59.91
1.5387	4	062	60.08
1.5144	7	341	61.15
1.4989	2	180	61.85
1.4836	4	162	62.56
1.4734	6	081	63.04

Barium silicate, β -BaSiO₃ - continued

d (Å)	I	hkl	2 θ (°)
1.4649	4	113	63.45
1.4458	3	252	64.39
1.4350	3	123,033	64.93
1.4280	6	271	65.29
1.4247	3	181	65.46
1.4126	4	322	66.09
1.3958	4	410	66.99
1.3901	5	360,133	67.30
1.3690	3	332	68.48
1.3629	6	172	68.83
1.3490	1	262	69.64
1.3433	<1	401,203 +	69.98
1.3347	4	411,213	70.50
1.3304	5	430,361	70.76
1.3145	4	342	71.75
1.3126	4	421,223	71.87
1.3046	4	281	72.38
1.2886	4	191	73.42
1.2775	5	431,233	74.17
1.2688	3	153	74.76
1.2548	2	182	75.74
1.2444	2	0•10•0	76.49
1.2409	2	371,290	76.74
1.2304	2	063	77.52
1.2231	2	450	78.07
1.2009	1	0•10•1	79.80

Barium silicate (sanbornite), β -BaSi₂O₅

Sample

The sample was prepared by melting a 1:2 molar mixture of BaCO₃ and silica gel (at about 1430 °C) and annealing for 15 hours at 1325 °C. Because of the presence of a small amount of the high (α) form, the intensities are subject to a slight uncertainty.

Color

Colorless

Structure

Orthorhombic, Pmnb(62), Z=4. The structure has been determined by Douglass [1958].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 7.6922(8) \text{ \AA} \\ b &= 13.525(1) \\ c &= 4.6336(5) \end{aligned}$$

Density

(calculated) 3.769 g/cm³

Polymorphism

There is a high (α) form stable above 1350 °C [Roth and Levin, 1959]. This is given on PDF card 10-45 [Klasens et al., 1957].

Additional patterns

1. PDF card 10-46 [Klasens et al., 1957].
2. PDF card 11-170 (natural mineral) [Douglass, 1958].
3. Levin and Ugrinic [1953].
4. Oehlschlegel [1971].
5. Roth and Levin [1959].

References

- Douglass, R. M. (1958). Amer. Mineral. 43, 517.
 Klasens, H. A., Hockstra, A. H., and Cox, A. P.M. (1957). J. Electrochem. Soc. 104, 93.
 Levin, E. M. and Ugrinic, G. M. (1953). J. Res. Nat. Bur. Stand. 51, 37.
 Oehlschlegel, G. (1971). Glastechn. Ber. 44, 194.
 Roth, R.S. and Levin, E.M. (1959). J. Res. Nat. Bur. Stand. 62, 193.

CuKα ₁ λ = 1.540598 Å; temp. 25±1 °C			
Internal standard Si, a = 5.43088 Å			
d(Å)	I	hkl	2θ(°)
6.77	40	020	13.06
5.08	25	120	17.43
3.973	85	101	22.36
3.844	6	200	23.12
3.808	9	111	23.34
3.424	50	121	26.00
3.382	16	040	26.33
3.343	70	220	26.64
3.234	30	031	27.56
3.097	100	140	28.80
2.981	4	131	29.95
2.892	5	211	30.90
2.732	35	041	32.75
2.712	40	221	33.00
2.575	18	141	34.81
2.539	6	240	35.32
2.399	4	320	37.46
2.337	8	051	38.49
2.317	13	002	38.83
2.285	3	012	39.40
2.244	11	301	40.16
2.236	13	151	40.30
2.227	35	241	40.47
2.192	14	022	41.15
2.164	25	160	41.70
2.130	35	321	42.41
2.108	10	122	42.86
2.059	4	032	43.93
2.043	9	340	44.31
2.027	20	061	44.66
1.996	5	251	45.40
1.9907	8	132	45.53
1.9845	6	202	45.68
1.9233	9	400	47.22
1.9047	11	222	47.71
1.8693	3	341	48.67
1.8547	12	142	49.08
1.8501	10	420	49.21
1.8166	3	232	50.18
1.7832	19	261	50.88
1.7367	3	171	52.66
1.7114	5	242	53.50
1.7049	5	312	53.72
1.6918	16	080	54.17
1.6716	3	440	54.88
1.6663	3	322	55.07
1.6516	5	431,180	55.60
1.6175	3	271	56.88
1.6000	5	252	57.56
1.5881	6	081	58.03

Barium silicate (sanbornite), β -BaSi₂O₅- continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.5812	10	162	58.31
1.5723	8	441	58.67
1.5552	6	181	59.38
1.5327	4	342	60.34
1.5144	2	103	61.15
1.5057	2	023,113	61.54
1.4840	5	451,072	62.54
1.4797	7	402	62.74
1.4676	8	281	63.32
1.4597	7	501	63.70
1.4508	4	511,352	64.14
1.4452	4	422	64.42
1.4352	2	133	64.92
1.4255	3	213	65.42
1.4115	4	380	66.15
1.4004	7	540	66.74
1.3953	8	461	67.02
1.3666	5	362,233	68.62
1.3555	3	442	69.26
1.3506	3	381	69.55
1.3443	4	182	69.92
1.3410	5	053,541	70.12
1.3319	2	1 \cdot 10 \cdot 0	70.67
1.3228	6	303	71.23
1.3196	5	243	71.43
1.2982	5	0 \cdot 10 \cdot 1,452	72.79
1.2801	8	1 \cdot 10 \cdot 1	73.99
1.2758	9	2 \cdot 10 \cdot 0,512	74.28
1.2696	5	480,333	74.71
1.2594	3	620,522	75.42
1.2328	3	532	77.34
1.2321	2	343	77.39
1.2244	3	481	77.97
1.2054	4	382	79.44

Barium silicate, Ba₂SiO₄

Sample

The sample was prepared by heating a 2:1 molar mixture of BaCO₃ and silicic acid at 1000 °C overnight, grinding and reheating at 1400 °C for 2 hours.

Color

Colorless

Structure

Orthorhombic, Pnam(62), Z=4, isostructural with α-K₂SO₄ [O'Daniel and Tscheischwili, 1942].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 7.508(1) \text{ \AA} \\ b &= 10.214(1) \\ c &= 5.8091(8) \end{aligned}$$

Density

(calculated) 5.468 g/cm³

Reference intensity

I/I_{corundum} = 1.8

Additional patterns

1. PDF card 6-366 [Levin and Ugrinic, 1953].
2. Austin [1947].
3. Budnikov and Kulikova [1966].
4. Glushkova and Keler [1957].
5. Grebenshchikov et al. [1956].
6. O'Daniel and Tscheischwili [1942].
7. Shitova and Grebenshchikov [1972].
8. Toropov and Grebenshchikov [1956].

References

- Austin, A.E. (1947). J. Amer. Ceram. Soc. 30, 218.
 Budnikov, P.P. and Kulikova, N. V. (1966). Inorg. Mater. (USSR) 2, 1717.
 Glushkova, V.B. and Keler, E.K. (1957). J. Inorg. Chem. (USSR) 2 [6], 63.
 Grebenshchikov, R.G., Toropov, N. A. and Shitova, V. I. (1965). Inorg. Mater. (USSR) 1, 105.
 Levin, E. M. and Ugrinic, G. (1953). J. Res. Nat. Bur. Stand. 51, 37.
 O'Daniel, H. and Tscheischwili, L. (1942). Z. Kristallogr. 104, 348.
 Shitova, V. I. and Grebenshchikov, R. G. (1972). J. Appl. Chem. USSR 45, 187.
 Toropov, N. A. and Grebenshchikov, R. G. (1956). J. Inorg. Chem. (USSR) 1 [12], 41.

CuKα₁ λ = 1.540598 Å; temp. 25±1 °C

Internal standard Ag, a = 4.08651 Å

d (Å)	I	hkl	2θ (°)
5.11	10	020	17.34
4.22	16	120	21.02
4.20	25	111	21.16
3.524	14	210	25.25
3.415	80	121	26.07

d (Å)	I	hkl	2θ (°)
3.153	25	201	28.28
3.098	20	130	28.79
3.022	70	220	29.53
3.017	100	211	29.59
2.938	95	031	30.40
2.905	70	002	30.75
2.683	13	221	33.37
2.554	9	040	35.11
2.525	20	022	35.53
2.431	40	310	36.95
2.393	20	122	37.55
2.297	5	202	39.18
2.242	20	212	40.19
2.233	19	141	40.35
2.120	20	132	42.62
2.095	30	222	43.14
2.017	12	330	44.91
1.984	6	241	45.69
1.971	16	150	46.00
1.928	4	051	47.11
1.918	2	042	47.37
1.904	17	232	47.72
1.877	3	400	48.47
1.864	25	312	48.82
1.844	4	113	49.38
1.795	3	250	50.84
1.788	6	340	51.04
1.786	5	401	51.11
1.7594	25	411	51.93
1.7203	5	203	53.20
1.7084	35	341	53.60
1.6970	14	213	53.99
1.6832	14	033	54.47
1.6566	6	332	55.42
1.6438	6	430	55.89
1.6309	16	223	56.37
1.5967	5	161	57.69
1.5767	2	402	58.49
1.5507	4	260	59.57
1.5115	6	143	61.28
1.5066	6	422	61.50
1.4976	6	261	61.91
1.4684	4	062	63.28
1.4639	2	441	63.50
1.4524	9	004	64.06
1.4396	8	511	64.70
1.4301	7	432	65.18
1.4151	6	071	65.96
1.3822	2	450	67.74
1.3680	11	361	68.54
1.3479	3	403	69.71
1.3367	5	531	70.38

Barium silicate, Ba₂Si₃O₈

Sample

The sample was prepared by repeated grinding and heating at about 1400 °C of a 2:3 molar mixture of BaCO₃ and silica gel.

Color

Colorless

Structure

Monoclinic, P2₁/a (14), Z = 4 [Kalscher and Liebau, 1965; Oehlschlegel, 1971]. Ba₂Si₃O₈ had earlier been reported with a similar cell with a/2 [Roth and Levin, 1959].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 13.960(3) \text{ \AA} \\ b &= 4.6895(9) \\ c &= 12.486(2) \\ \beta &= 93.54(1)^\circ \end{aligned}$$

Density

(calculated) 3.964 g/cm³

Reference intensity

I/I_{corundum} = 1.8

Polymorphism

Oehlschlegel [1971] reports a reversible transformation of Ba₂Si₃O₈ at 1009 °C.

Additional patterns

1. PDF card 12-694 [Roth and Levin, 1959].
2. Austin [1947].

References

- Austin, A.E. (1947). J. Am. Ceram. Soc. 30, 218.
Kalscher, H. and Liebau, F. (1965). Naturwiss. 52, 512.
Oehlschlegel, G. (1971). Glastechn. Ber. 44, 194.
Roth, R.S. and Levin, E.M. (1959). J. Res. Nat. Bur. Stand. 62, 193.

CuKα₁ λ = 1.540598 Å; temp. 25±1 °C

Internal standard Ag, a = 4.08651 Å

d(Å)	I	hkl	2θ(°)
12.51	10	001	7.06
6.965	20	200	12.70
6.245	5	201	14.17
5.929	16	201	14.93
4.512	2	202	19.66
4.390	2	011	20.21
4.162	2	111,003	21.33
3.890	4	210	22.84
3.746	55	211,012	23.73
3.669	100	203	24.24
3.478	5	400,203	25.59
3.415	5	401	26.07
3.301	70	401,310	26.99
3.250	30	212	27.42
3.121	40	402	28.58
3.114	40	004	28.64
2.864	1	312	31.21
2.792	13	213	32.03
2.782	60	204	32.15
2.756	30	411,403	32.46
2.701	2	411	33.14
2.598	3	412	34.49
2.493	4	005	36.00
2.475	1	214	36.26
2.398	5	404,510	37.48
2.393	5	205,214	37.55
2.376	2	511,413	37.84
2.346	10	020	38.34
2.310	3	601	38.96
2.301	3	205	39.11
2.268	19	121,413	39.71
2.221	13	220,602	40.58
2.201	9	015	40.98
2.180	2	221	41.38
2.133	30	602,215 +	42.34
2.080	8	610,222	43.48
2.074	6	321	43.60
2.042	2	023	44.33
2.032	5	315,414	44.55
1.975	9	223,603	45.90
1.971	11	405	46.02
1.911	6	421	47.53
1.908	4	415	47.62
1.900	2	016	47.83
1.875	4	422,024	48.52
1.834	4	406	49.66
1.820	9	613	50.07
1.807	10	216	50.46
1.793	6	316,224	50.90
1.785	7	712,423	51.12

Barium silicate, $\text{Ba}_2\text{Si}_3\text{O}_8$ - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.780	19	007	51.28
1.776	13	$\bar{6}14, \bar{5}15$	51.42
1.753	6	$\bar{6}05$	52.12
1.751	6	$\bar{2}07$	52.20
1.738	5	423, 406	52.63
1.705	2	$\bar{8}02, \bar{1}25$	53.73
1.649	8	620, 605	55.69
1.645	5	$\bar{6}21$	55.85
1.631	4	$\bar{8}11$	56.36
1.628	6	416	56.46
1.612	3	$\bar{6}22, \bar{5}16$	57.09
1.608	3	811	57.26
1.598	3	217, $\bar{6}06$	57.65
1.578	1	622	58.45
1.556	2	$\bar{6}23, \bar{6}15$	59.34
1.550	4	031, $\bar{8}13$	59.59
1.515	2	032, $\bar{7}21$	61.10
1.508	2	425	61.42
1.500	5	208	61.79
1.476	2	232, $\bar{3}31$	62.93
1.4697	3	910, $\bar{9}11$ +	63.22
1.4636	1	$\bar{2}18, 033$	63.51
1.4448	1	$\bar{4}26$	64.44
1.4208	3	$\bar{4}31$	65.66
1.4180	3	027, $\bar{1}27$	65.81
1.4038	2	127	66.56
1.3914	10	$\bar{6}17$	67.23
1.3485	3	625, $\bar{3}34$	69.67

Barium silicate, Ba₃SiO₅

Sample

The sample was prepared by repeated grindings and heatings at about 1400 °C of a 3:1 molar mixture of BaCO₃ and silica gel.

Color

Colorless

Structure

Tetragonal, I4/mcm (140), Z = 4, isostructural with Cs₃CoCl₅ and other similar compounds [Mansmann, 1965].

NBS lattice constants of this sample:

$$a = 7.3068(2)\text{\AA}$$

$$c = 11.2275(6)$$

Density

(calculated) 5.763 g/cm³

Reference Intensity

$$I/I_{\text{corundum}} = 2.4$$

Polymorphism

Since Glushkova and Keler [1957] and Budnikov and Kulikova [1966] report patterns which differ considerably from the present study, the possibility of polymorphism cannot be ruled out.

Additional patterns

1. PDF card 19-175 [Budnikov and Kulikova, 1966].
2. PDF card 23-1027 [Brisi and Appendino, 1966].
3. Glushkova and Keler [1957].
4. Eysel [1970].

References

- Brisi, C. and Appendino, P. (1966). Ric. Sci. 36, 369.
- Budnikov, P.P. and Kulikova, N.V. (1966). Inorg. Mater. (USSR), 2, 1717.
- Eysel, W. (1970). Neues Jahrb. Mineral. Monatsh. 1970, 534.
- Glushkova, V.B. and Keler, E.K. (1957). J. Inorg. Chem. (USSR) 2 [6], 63.
- Mansmann, M. (1965). Z. Anorg. Allg. Chem. 339, 52.

CuKα ₁ λ = 1.540598 Å; temp. 25±1 °C			
Internal standard W, a = 3.16524 Å			
d(Å)	I	hkl	2θ(°)
5.619	7	002	15.76
5.169	4	110	17.14
3.802	20	112	23.38
3.138	55	211	28.42
3.062	100	202	29.14
2.808	25	004	31.84
2.584	30	220	34.69
2.462	60	213	36.47
2.311	30	310	38.94
1.994	4	321	45.44
1.901	20	224	47.82
1.872	2	006	48.60
1.851	12	215	49.19
1.827	1	400	49.87
1.784	16	314	51.17
1.760	7	116	51.92
1.751	20	411	52.20
1.738	5	402	52.61
1.7224	2	330	53.13
1.6657	14	206	55.09
1.6465	16	332	55.79
1.6338	6	420	56.26
1.6018	16	413	57.49
1.5687	4	422	58.82
1.5311	3	404	60.41
1.5041	1	325	61.61
1.4676	1	334	63.32
1.4492	1	431	64.22
1.4402	4	217	64.67
1.4120	6	424	66.12
1.4034	3	008	66.58
1.3912	11	415	67.24
1.3881	10	512	67.41
1.3614	1	433	68.92
1.3473	1	521	69.74
1.3074	3	406	72.20
1.2917	2	440	73.22
1.2757	2	523	74.29
1.2677	6	336	74.84
1.2584	2	442	75.49
1.2534	5	530	75.84
1.2335	4	228	77.29
1.2307	3	426	77.50
1.2231	3	532	78.07
1.2179	4	600	78.47
1.1994	3	318	79.92
1.1944	3	611	80.32
1.1892	5	417	80.74
1.1737	1	444	82.04
1.1655	3	219	82.74

Barium silicate, Ba_3SiO_5 - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.1615	2	525	83.09
1.1442	8	534	84.63
1.1376	2	516	85.24
1.1350	2	541	85.48
1.1317	6	622	85.79
1.1172	5	604	87.18
1.1129	2	408	87.60
1.0916	1	543	89.77
1.0802	1	437	90.97
1.0732	3	2•0•10	91.74
1.0646	3	428	92.70
1.0591	2	615	93.32
1.0412	2	536	95.43
1.0335	1	710	96.38
1.0201	4	419	98.07
1.0174	3	545	98.42
1.0163	5	712	98.57
0.9997	2	721	100.80
.9972	4	642	101.15
.9830	5	626	103.19
.9742	1	2•1•11	104.51
.9695	3	554,723.	105.22
.9615	2	617	106.48
.9565	1	4•0•10	107.28
.9530	2	644	107.85
.9504	1	448	108.29
.9457	2	732	109.09
.9405	2	3•3•10	109.97
.9347	6	538	111.00
.9298	1	547	111.88
.9252	<1	4•2•10	112.73
.9198	3	608	113.75
.9162	3	725	114.43
.9044	4	716	116.79
.9033	5	741	117.02
.9013	5	802,637	117.45
.8909	2	646	119.67
.8860	3	820	120.78
.8838	4	5•1•10	121.29
.8808	4	743	121.99

Barium silicate, Ba₃Si₅O₁₃

Sample

The sample was prepared by heating a 3:5 molar mixture of BaCO₃ and silica gel at about 1400°C with repeated grindings and reheatings. Because of problems related to orientation, the intensities are subject to some uncertainty.

Color

Colorless

Optical data

Biaxial(+), N_α = 1.612, N_β = 1.616, N_γ = 1.636. 2V is about 35° [Oehlschlegel, 1971].

Structure

Monoclinic, P2₁/c (14), Z = 4 [Roth, 1966; Oehlschlegel, 1971].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 20.208(3) \text{ \AA} \\ b &= 4.7106(5) \\ c &= 13.854(2) \\ \beta &= 98.62(1)^\circ \end{aligned}$$

Density

(calculated) 3.874 g/cm³

Additional patterns

1. PDF card 12-547 [Roth and Levin, 1959].
2. Oehlschlegel [1971].

References

- Oehlschlegel, G. (1971). Glastechn. Ber. 44, 194.
 Roth, R. (1966). Private comm. to Crystal Data (3rd Ed., published jointly by the U.S. Dept. of Commerce, National Bureau of Standards, Washington, D.C. 20234, and the Joint Committee on Powder Diffraction Standards, Swarthmore, Pa., 19081).
 Roth, R. and Levin, E. M. (1959). J. Res. Nat. Bur. Stand. 62, 193.

CuKα₁ λ = 1.540598 Å; temp. 25±1 °C

Internal standard Si, a = 5.43088 Å

d (Å)	I	hkl	2θ (°)
9.96	4	200	8.87
6.80	16	102	13.00
6.202	6	102	14.27
6.091	15	202	14.53
5.181	3	302	17.10

d (Å)	I	hkl	2θ (°)
4.996	2	400	17.74
4.263	5	210	20.82
3.998	3	211, 500	22.22
3.870	11	112	22.96
3.844	55	310	23.12
3.773	100	402	23.56
3.729	10	212	23.84
3.622	2	311	24.56
3.484	3	312	25.55
3.424	7	410, 004	26.00
3.328	20	600	26.77
3.249	85	304, 411 +	27.43
3.199	40	412	27.87
3.100	20	204	28.78
3.045	2	510, 404	29.31
2.909	6	512	30.71
2.875	13	304	31.08
2.855	5	700	31.31
2.788	75	114, 702	32.08
2.769	25	014	32.30
2.718	2	610	32.93
2.698	3	114	33.18
2.644	3	404	33.87
2.607	2	611	34.37
2.589	2	214, 604	34.62
2.558	1	414	35.05
2.497	1	800	35.93
2.469	3	802	36.36
2.443	2	710	36.76
2.429	4	612, 504	36.98
2.356	14	020	38.17
2.283	6	006	39.44
2.273	25	221	39.61
2.269	13	614	39.69
2.241	5	802, 215	40.21
2.230	18	106	40.41
2.224	15	122, 811	40.53
2.211	35	712, 902	40.77
2.206	30	810, 406	40.87
2.175	4	321	41.49
2.158	9	514, 206	41.82
2.124	5	421, 506	42.53
2.068	11	216, 123 +	43.73
2.043	3	316	44.31
2.025	7	902, 812	44.72
2.008	8	910	45.11
2.004	16	904	45.20
1.999	15	422, 10•0•0 +	45.33
1.980	2	423	45.79
1.923	2	620	47.22

Barium silicate, Ba₃Si₅O₁₃ - continued

d(Å)	I	hkl	2θ(°)
1.908	8	$\bar{5}23, \bar{3}24$	47.61
1.888	5	804	48.15
1.8755	4	224	48.50
1.8504	4	$\bar{1}0\cdot0\cdot4$	49.20
1.8445	12	$10\cdot0\cdot2, \bar{9}14$	49.37
1.8410	15	$\bar{1}0\cdot1\cdot2, 10\cdot1\cdot0$	49.47
1.8223	8	$\bar{1}17, 324 +$	50.01
1.8162	35	$11\cdot0\cdot0, 416 +$	50.19
1.7988	8	$\bar{7}22$	50.71
1.7854	11	$025, \bar{7}16$	51.12
1.7635	4	125, 606	51.80
1.7276	6	$\bar{1}08, \bar{9}06$	52.96
1.7040	2	$\bar{8}22, \bar{8}16$	53.75
1.6657	2	$706, 12\cdot0\cdot0 +$	55.09
1.6386	3	$026, 914$	56.08
1.6296	12	$10\cdot1\cdot3, \bar{1}0\cdot0\cdot6$	56.42
1.6253	8	$\bar{9}21, \bar{2}18 +$	56.58
1.6209	8	$\bar{9}16$	56.75
1.6167	8	$\bar{3}18, 815 +$	56.91
1.6118	7	$\bar{9}22, 11\cdot1\cdot4$	57.10
1.5984	3	$\bar{4}18$	57.62
1.5782	2	$\bar{9}23, \bar{5}26$	58.43
1.5547	6	$\bar{2}18$	59.40
1.5509	9	$230, \bar{1}3\cdot0\cdot2$	59.56
1.5360	2	$\bar{6}18, 922$	60.20
1.5284	4	330	60.53
1.5141	2	$\bar{2}27, \bar{1}27 +$	61.16
1.5103	1	$\bar{1}2\cdot1\cdot4, 426$	61.33
1.4948	2	$508, \bar{7}18 +$	62.04
1.4859	4	$\bar{1}33, 12\cdot1\cdot2$	62.45
1.4770	2	$\bar{4}32$	62.87
1.4732	2	$923, 824 +$	63.05
1.4616	3	$\bar{5}31, 530 +$	63.61
1.4526	2	$\bar{1}2\cdot0\cdot6, 10\cdot2\cdot2 +$	64.05
1.4498	2	432	64.19
1.4418	7	$\bar{1}1\cdot2\cdot2, \bar{1}4\cdot0\cdot2$	64.59
1.4386	7	$11\cdot2\cdot0, 10\cdot1\cdot5$	64.75
1.4272	2	$034, 14\cdot0\cdot0$	65.33
1.4155	4	$12\cdot0\cdot4, 916 +$	65.94
1.4117	4	626	66.14
1.4003	3	$\bar{9}18, \bar{7}27$	66.75
1.3936	8	$\bar{1}4\cdot0\cdot4, \bar{1}28$	67.11
1.3852	2	$\bar{2}\cdot0\cdot10, 028$	67.57
1.3782	2	$\bar{4}28, 334$	67.96
1.3662	4	$\bar{1}35, 12\cdot2\cdot2 +$	68.64
1.3583	1	$11\cdot1\cdot5, 527$	69.10

Cadmium silicate, Cd_2SiO_4

Sample

The sample was prepared by heating at 1200 °C, for several hours, a 2:1 molar mixture of CdO and silica gel. The sample was ground and reheated several times at 1000 °C for one hour each time. A small amount of Cd_3SiO_5 was present and this may slightly distort the intensity measurements.

Color

Colorless

Structure

Orthorhombic, $Fddd(70)$, $Z=8$. Isostructural with $\text{Na}_2\text{SO}_4(V)$. The structure was studied by Glasser and Glasser [1964].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 9.805(1)\text{\AA} \\ b &= 11.807(2) \\ c &= 6.013(1) \end{aligned}$$

Density

(calculated) 6.047 g/cm³

Reference intensity

$$I/I_{\text{corundum}} = 2.1$$

Additional pattern

1. PDF card 17-258 [Glasser and Glasser, 1964].

References

Glasser, L.S.D. and Glasser, F.P. (1964). Inorg. Chem. 3, 1228.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. 25±1 °C			
Internal standard W, $a = 3.16524 \text{ \AA}$			
d(Å)	I	hkl	2 θ (°)
4.704	15	111	18.85
3.770	45	220	23.58
2.951	65	040	30.26
2.790	100	311	32.06
2.678	90	022	33.43
2.563	25	202	34.98
2.449	3	400	36.66
2.352	6	222	38.24
2.320	35	331	38.78
2.145	8	151	42.10
1.936	20	113,242	46.90
1.841	20	511	49.48
1.824	35	260,351	49.97
1.758	3	133	51.99
1.691	20	313	54.19
1.685	13	531	54.39
1.647	20	062	55.78
1.603	2	171	57.45
1.575	7	620	58.55
1.567	14	333	58.88
1.504	5	004	61.63
1.476	4	080	62.91
1.454	10	371	63.96
1.436	6	602	64.90
1.396	5	224	66.97
1.393	5	513	67.17
1.3843	6	353	67.62
1.3396	8	044	70.20
1.3199	2	533	71.41
1.2909	5	642	73.27
1.2789	3	282	74.07
1.2571	3	660	75.58
1.2259	1	800	77.86
1.2052	3	553	79.46
1.2003	4	373	79.85
1.1932	3	391	80.42

Cadmium silicate, Cd_3SiO_5

Sample

The sample was made by heating a 3:1 molar mixture of CdO and silica gel at 1100°C for 2 hours. The product was then ground and reheated at 700°C for 20 hours. The sample showed some hydration products after standing in air and also contained a very slight percentage of Cd_2SiO_4 ; therefore, the intensities may be slightly in error.

Color

Greenish yellow.

Structure

Tetragonal, $P4/nmm$ (129), $Z = 2$ [Eysel, 1970]. Eysel (1970) suggested also a possible monoclinic cell. The broadening of some lines in patterns from this sample indicates that it probably is of lower symmetry.

NBS lattice constants of this sample:

$$a = 6.842(2)\text{\AA}$$

$$c = 4.952(2)$$

Density

(calculated) 6.379 g/cm^3

Reference intensity

$$I/I_{\text{corundum}} = 5.4$$

Additional patterns

1. PDF card 17-257 [Dent Glasser and Glasser, 1964].
2. Eysel [1970].

References

- Dent Glasser, L. S. and Glasser, F. P. (1964). *Inorg. Chem.* 3, 1228.
- Eysel, W. (1970). *Neues Jahrb. Mineral. Monatsh.* 1970, 534.

CuK α_1 $\lambda = 1.540598\text{\AA}$; temp. $25 \pm 1^\circ\text{C}$			
Internal standard W, $a = 3.16524\text{\AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
4.96	5	001	17.88
4.85	8	110	18.29
4.015	3	101	22.12
3.462	5	111	25.71
3.420	2	200	26.03
2.814	100	201	31.77
2.604	<1	211	34.41
2.476	11	002	36.25
2.419	20	220	37.14
2.327	4	102	38.66
2.206	3	112	40.88
2.173	5	221	41.53
2.165	5	310	41.68
2.071	2	301	43.68
2.006	3	202	45.16
1.982	3	311	45.70
1.924	1	212	47.20
1.730	20	222	52.87
1.710	10	400	53.55
1.677	1	302	54.68
1.629	1	312	56.44
1.616	1	401	56.94
1.613	2	330	57.04
1.562	1	113	59.08
1.530	1	420	60.44
1.5066	1	322	61.50
1.4861	7	203	62.44
1.4616	14	421	63.61
1.4072	6	402	66.38
1.3782	1	412	67.96

Calcium hydrogen phosphate hydrate, $\text{Ca}_8\text{H}_2(\text{PO}_4)_6 \cdot 5\text{H}_2\text{O}$

Sample

The sample was obtained from B. Dickens at NBS. Brown et al. [1962] prepared the sample. The intensity of the strongest line was very high compared to the other reflections. Therefore, the intensity of the second strongest line ($d=2.833$) was assigned the value of 100 and all other reflections were scaled to it. On that scale the strongest line at $d=18.67$ has $I \approx 300$.

Color

Colorless

Optical data

Biaxial(-), $N_\alpha = 1.576$, $N_\beta = 1.583$, $N_\gamma = 1.585$. $2V$ is $\approx 50^\circ$ [Brown et al, 1962].

Structure

Triclinic, $Z = 2$. The structure was determined by Brown et al. [1962] and refined by Dickens et al. [1973].

NBS lattice constants of this sample:

$a = 9.529(3)\text{\AA}$
 $b = 18.994(4)$
 $c = 6.855(3)$
 $\alpha = 92.33(3)^\circ$
 $\beta = 90.13(3)$
 $\gamma = 79.93(2)$

Density

(calculated) 2.673 g/cm^3 .

Reference intensity

$I/I_{\text{corundum}} = 0.5$. This measurement is based on the line at 2.833\AA (designated as 100).

Additional patterns

1. PDF card 11-184 [Bjerrum, 1958].
2. PDF card 13-391 [Hayek et al, 1960].
3. Lehr et al. [1967].

References

- Bjerrum, N. (1958). Kgl. Dan. Vidensk. Selsk. Mat. Fys. Medd. 31, Nr. 7, 22.
- Brown, W.E., Smith, J.P., Lehr, J.R., and Frazier, A. W. (1962). Nature (London) 196, 1050.
- Dickens, B., Schroeder, L. W., and Brown, W. E. (1973). Am. Crys. Assoc. (Abs.-Winter Meeting) B2, 26.
- Hayek, E., Newesely, H., Hassenteufel, W., and Krismer, B. (1960). Monatsh. Chem. 91, 249.
- Lehr, J. R., Brown, E. H., Frazier, A. W., Smith, J.P., and Thrasher, R.D. (1967). Tenn. Val. Auth. (Chem. Eng. Bull.) No. 6.

$\text{CuK}\alpha_1, \lambda = 1.540598 \text{ \AA}; \text{ temp. } 25 \pm 1^\circ \text{C}$ Internal standard W, $a = 3.16524 \text{ \AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
18.67	300	010	4.73
9.36	45	100,020	9.44
9.05	40	110	9.77
6.10	6	120	14.51
5.52	25	101	16.04
5.417	7	111,021	16.35
5.211	4	111	17.00
5.101	12	111	17.37
4.815	6	130	18.41
4.706	5	031	18.84
4.670	4	040	18.99
4.514	10	031,140 +	19.65
4.492	10	121	19.75
4.294	7	131	20.67
4.111	5	230	21.60
3.919	16	220,140 +	22.67
3.879	12	201,131	22.91
3.862	10	201	23.01
3.786	10	041	23.48
3.745	14	221	23.74
3.660	30	211	24.30
3.492	25	231	25.49
3.441	50	221	25.87
3.424	60	002	26.00
3.378	18	221	26.36
3.311	20	151	26.91
3.278	18	150	27.18
3.209	25	102,250 +	27.78
3.180	25	241,310	28.04
3.132	10	122,300 +	28.48
3.117	7	112,060	28.62
3.055	14	032,240	29.21
3.015	8	330	29.61
2.946	14	122,251	30.31
2.914	12	151	30.66
2.873	30	251	31.10
2.833	100	260	31.55
2.820	95	320,241	31.70
2.779	45	142,331	32.18
2.745	35	132,331	32.59
2.707	25	222,042	33.06
2.671	50	070	33.53
2.637	35	161,350	33.97
2.617	20	330	34.23
2.606	20	222,341	34.38
2.567	16	161,152	34.93
2.544	12	171,251	35.25
2.486	5	251	36.10
2.475	8	052,171	36.27
2.458	5	170	36.52

Calcium hydrogen phosphate hydrate, $\text{Ca}_8\text{H}_2(\text{PO}_4)_6 \cdot 5\text{H}_2\text{O}$ - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
2.365	7	180	38.01
2.335	8	080, 271 +	38.52
2.304	7	252, 302 +	39.06
2.271	5	361, 312	39.66
2.265	6	181, 162	39.77
2.258	7	062, 341	39.89
2.215	16	162, 350	40.71
2.158	5	322, 441	41.83
2.136	7	441	42.27
2.106	9	190	42.90
2.088	7	252	43.29
2.063	6	272, 213 +	43.85
2.036	5	133, 360 +	44.46
2.002	8	262	45.25
1.998	9	431	45.35
1.990	10	191	45.55
1.957	7	422, 190	46.36
1.948	17	342, 381 +	46.58
1.936	18	361	46.88
1.929	11	291, 233	47.06
1.914	11	053	47.46
1.897	10	100, 372 +	47.92
1.891	10	530	48.07
1.848	20	303, 1101	49.27
1.837	20	352, 391 +	49.59
1.832	18	253, 511 +	49.72
1.804	15	391, 2101	50.54
1.745	8	291, 462	52.39
1.743	8	551	52.47
1.725	10	1110	53.05
1.710	25	490	53.55

Cobalt phosphate, $\text{Co}(\text{PO}_3)_2$

Sample

The sample was prepared by heating a 1:2 molar mixture of CoCO_3 and H_3PO_4 to about 640 °C for 15 hours.

Color

Deep purplish red.

Structure

Monoclinic, $I2/a(15)$ or $Ia(9)$, $Z = 8$ [Beucher and Grenier, 1968]. These authors gave the cell in the settings $C2/c(15)$ or $Cc(9)$.

NBS lattice constants of this sample:

$$\begin{aligned} a &= 11.189(3) \text{ \AA} \\ b &= 8.287(2) \\ c &= 9.926(4) \\ \beta &= 112.42(3)^\circ \end{aligned}$$

Density

(calculated) 3.386 g/cm³

Reference intensity

$$I/I_{\text{corundum}} = 1.4$$

Additional pattern

1. PDF card 19-351 (Sarver, 1966).

References

Beucher, M. and Grenier, J.-C. (1968). Mater. Res. Bull. 3, 643.
Sarver, J.F. (1966). Trans. Brit. Ceram. Soc. 65, 191.

$\text{CuK}\alpha_1$ $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1 \text{ }^\circ\text{C}$

Internal standard W, $a = 3.16524 \text{ \AA}$

$d(\text{Å})$	I	hkl	$2\theta(^\circ)$
6.45	6	110	13.71
6.146	40	011	14.40
4.576	20	$\bar{2}11$	19.38
4.251	35	$\bar{1}12$	20.88
3.742	6	$\bar{1}21$	23.76
3.538	20	211	25.15
3.378	30	112,121	26.36
3.232	12	220	27.58
3.184	30	310	28.00
3.001	100	$\bar{2}22$	29.75
2.868	20	013	31.16
2.670	3	$\bar{1}30$	33.54
2.635	4	$\bar{4}11$	34.00
2.586	20	400	34.66
2.466	4	$\bar{2}31$	36.40
2.389	20	222	37.62
2.378	9	$\bar{3}23$	37.80
2.279	6	314	39.51
2.193	7	420,411	41.13
2.177	7	404	41.45
2.156	3	330, $\bar{5}12$	41.86
2.099	20	$\bar{2}33$	43.07
2.071	4	040	43.68
2.016	8	114, $\bar{1}41$	44.93
1.956	6	402	46.38
1.928	4	$\bar{4}24$	47.10
1.895	2	$\bar{5}23$	47.96
1.872	3	$\bar{2}42$	48.61
1.854	3	$\bar{4}33$	49.10
1.824	2	$\bar{4}15$	49.97
1.799	5	$\bar{3}34$	50.71
1.757	8	233	52.00
1.754	6	431, $\bar{1}43$	52.09
1.735	6	$\bar{5}32$	52.70
1.701	7	$\bar{6}22$	53.85
1.6314	10	051, $\bar{5}12$	56.35
1.6167	7	440	56.91
1.6115	7	$\bar{2}35$,314	57.11
1.5376	6	044	60.13
1.5295	6	006	60.48
1.5250	7	$\bar{6}33$	60.68
1.5009	4	444	61.76

Copper imidazole nitrate, $\text{Cu}(\text{C}_3\text{H}_4\text{N}_2)_4(\text{NO}_3)_2$

Sample

The sample was prepared at NBS by C. W. Reimann by evaporating an aqueous solution of $\text{Cu}(\text{NO}_3)_2$ and imidazole ($\text{C}_3\text{H}_4\text{N}_2$) at room temperature. It was difficult to obtain intensities because the sample deteriorated somewhat when exposed to x-rays.

Color

Unground: deep blue

Optical Data

Biaxial (-), $N_\alpha = 1.584$, $N_\beta = 1.610$, $N_\gamma = 1.645$. $2V \approx 40^\circ$. The sample shows pleochroism.

Structure

Orthorhombic, $\text{Pmnb}(62)$, $Z=4$, [Mighell, Santoro, and Reimann, private comm.].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 13.396(3) \text{ \AA} \\ b &= 13.858(3) \\ c &= 9.825(2) \end{aligned}$$

Density

(calculated) 1.675 g/cm^3

Reference intensity

$I/I_{\text{corundum}} = 1.0$

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1^\circ \text{C}$			
Internal standard Ag, $a = 4.08651 \text{ \AA}$			
d(Å)	I	hkl	2 θ (°)
8.01	8	011	11.04
6.87	19	111	12.87
6.697	6	200	13.21
6.159	25	120	14.37
5.142	25	211	17.23
4.911	5	002	18.05
4.629	3	012	19.16
4.375	16	112	20.28
4.182	4	031	21.23
4.066	10	301	21.84
3.962	100	202	22.42
3.906	25	311	22.75
3.839	18	122	23.15
3.807	18	212	23.35
3.756	40	320	23.67

d(Å)	I	hkl	2 θ (°)
3.548	8	231	25.08
3.464	85	040	25.70
3.352	8	140,400	26.57
3.264	9	041,132	27.30
3.179	7	103,141	28.05
3.092	25	411	28.85
3.051	7	331	29.25
3.009	10	232	29.67
2.984	10	322	29.92
2.938	3	241	30.40
2.883	8	421	30.99
2.832	1	042	31.57
2.768	16	142,402	32.32
2.736	6	340	32.70
2.707	8	223	33.06
2.666	5	051	33.59
2.638	14	341	33.96
2.615	25	151,431	34.26
2.608	16	242	34.36
2.541	2	511	35.30
2.498	4	520	35.92
2.477	4	251	36.23
2.456	5	004	36.56
2.379	4	114,043 +	37.79
2.338	4	441	38.47
2.306	13	204	39.03
2.277	20	160,214	39.55
2.243	6	243	40.18
2.228	11	522	40.46
2.183	2	260	41.32
2.161	6	442	41.76
2.119	2	540	42.63
2.087	5	433,451	43.31
2.065	6	162	43.80
2.050	10	360,513	44.14
2.033	5	602	44.53
2.004	2	044	45.20
1.980	4	144,404	45.79
1.958	1	452	46.33
1.941	3	071,443	46.77
1.921	6	171,244	47.29
1.891	8	025,551 +	48.08
1.866	4	461	48.75
1.861	5	711,632	48.89
1.845	2	720,641	49.36
1.828	4	613,344	49.85
1.780	7	371,543	51.28
1.752	3	642,524	52.16
1.749	2	560	52.26
1.733	3	080	52.78
1.727	3	722	52.99
1.710	2	045	53.56

Lead chloride fluoride (matlockite), PbClF

Sample

The sample was prepared by melting a 1:1 molar mixture of PbCl₂ and PbF₂ at about 600 °C.

Color

Yellowish gray

Structure

Tetragonal, P4/nmm (129), Z = 2, isostructural with BaClF and other similar double halides. The structure of PbClF was determined by Bannister [1934].

NBS lattice constants of this sample:

$$a = 4.1104(2) \text{ \AA}$$

$$c = 7.2325(5)$$

Density

(calculated) 7.111 g/cm³

Reference intensity

I/I_{corundum} = 6.2

Additional patterns

1. PDF card 4-460 [Swanson et al., 1953].
2. Nieuwenkamp and Bijvoet [1932].

References

- Bannister, F.A. (1934). Mineral. Mag. 23, 587.
 Nieuwenkamp, W. and Bijvoet, J. M. (1932). Z. Krist. 81, 469.
 Swanson, H. E. and Tatge, E. (1953). Nat. Bur. Stand. (U.S.) Circ. 539, 1, 76.

CuKα ₁ λ = 1.540598 Å; temp. 25±1 °C			
Internal standard W, a = 3.16524 Å			
d(Å)	I	hkl	2θ(°)
7.22	13	001	12.25
3.617	40	002	24.59
3.574	100	101	24.89
2.906	45	110	30.74
2.715	35	102	32.96
2.412	4	003	37.25
2.265	40	112	39.76
2.079	16	103	43.49
2.055	20	200	44.04
1.976	2	201	45.88
1.855	9	113	49.06
1.808	1	004	50.43
1.786	17	202	51.10
1.781	25	211	51.26
1.6552	11	104	55.47
1.6386	9	212	56.08
1.5636	2	203	59.03
1.5350	1	114	60.24
1.4618	7	213	63.60
1.4528	5	220	64.04
1.4466	3	005	64.35
1.4249	1	221	65.45
1.3645	1	105	68.74
1.3478	5	222	69.71
1.3458	4	301	69.83
1.3001	5	310	72.67
1.2952	6	115	72.99
1.2891	8	214	73.39
1.2814	2	302	73.90
1.2232	5	312	78.06
1.1910	1	303	80.60
1.1831	3	205	81.25
1.1567	2	106	83.51
1.1440	2	313	84.65
1.1368	1	215	85.31
1.1259	2	321	86.34
1.0920	2	304	89.72
1.0873	2	322	90.22
1.0331	1	007	96.43
1.0252	2	225	97.42
1.0079	1	216	99.68
0.9878	2	411	102.49
.9736	1	117	104.59
.9669	2	315	105.63
.9645	3	324	106.01

Magnesium phosphate, $\text{Mg}(\text{PO}_3)_2$

Sample

The sample was prepared by heating a 1:2 molar mixture of MgCO_3 and H_3PO_4 to 710 °C. It was then reground and reheated at 710 °C several times.

Color

Colorless

Structure

Monoclinic, $\text{I}2/\text{a}(15)$ or $\text{Ia}(9)$, $Z=8$ [Beucher and Grenier, 1968]. Those authors gave the cell in the settings $\text{C}2/\text{c}(15)$ or $\text{Cc}(9)$.

NBS lattice constants of this sample:

$$\begin{aligned} a &= 11.119(3) \text{ \AA} \\ b &= 8.268(2) \\ c &= 9.920(3) \\ \beta &= 112.44(3)^\circ \end{aligned}$$

Density

(calculated) 2.872 g/cm³

Reference intensity

$$I/I_{\text{corundum}} = 1.4$$

Additional pattern

1. PDF card 11-41 [Sarver and Hummel, 1959].

References

Beucher, M. and Grenier, J.-C. (1968). Mater. Res. Bull. 3, 643.
Sarver, J. F. and Hummel, F. A. (1959). J. Electrochem. Soc. 106, 500.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1 \text{ }^\circ\text{C}$			
Internal standard W, $a = 3.16524 \text{ \AA}$			
$d(\text{Å})$	I	hkl	$2\theta(^\circ)$
6.44	2	110	13.75
6.14	16	011	14.42
5.13	4	200	17.26
4.59	60	002	19.34
4.34	4	202	20.47
4.243	35	$\bar{1}12$	20.92
3.731	1	$\bar{1}21$	23.83
3.519	20	211	25.29
3.371	25	112,121	26.42
3.219	50	220	27.69
3.181	30	$\bar{3}12$	28.03
3.164	30	310	28.18
3.070	2	022	29.06
2.993	100	$\bar{2}22$	29.83
2.865	16	013	31.19
2.728	6	$\bar{4}02$	32.80
2.662	1	130	33.64
2.576	20	$\bar{1}23$	34.80
2.380	16	222	37.77
2.373	12	323	37.89
2.358	6	321	38.14
2.277	6	$\bar{4}22, \bar{3}14$	39.55
2.248	4	231	40.07
2.242	3	123	40.19
2.181	8	420,411	41.37
2.147	2	330	42.05
2.094	17	$\bar{2}33$	43.16
2.068	2	040	43.74
1.946	3	141,402	46.63
1.938	3	$\bar{5}21$	46.85
1.929	4	$\bar{2}15$	47.07
1.885	3	042	48.23
1.865	2	$\bar{2}42$	48.78
1.848	2	$\bar{4}33, 204$	49.27
1.828	2	134	49.83
1.805	4	$\bar{3}41$	50.53
1.796	6	$\bar{3}34$	50.81
1.755	6	125	52.06
1.754	6	233	52.10
1.751	4	143	52.19
1.728	6	$\bar{5}32$	52.96
1.692	7	622	54.18
1.648	4	442,530	55.74
1.627	7	051	56.51
1.623	8	512	56.67
1.6102	10	440	57.16
1.5353	8	044	60.23
1.5184	6	$\bar{6}33$	60.97
1.5148	6	$\bar{6}31$	61.13
1.4974	4	444	61.92

Magnesium tungsten oxide, MgWO_4

Sample

The sample was prepared by treating an aqueous solution of Na_2WO_4 with concentrated MgCl_2 at 80°C . The precipitate was filtered, washed with alcohol and heated at 850°C for 30 minutes.

Color

Colorless

Structure

Monoclinic, $P2/a$ (13), $Z=2$, isostructural with wolframite, $(\text{Fe,Mn})\text{WO}_4$ [Broch, 1929].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 4.9288(6)\text{\AA} \\ b &= 5.6751(8) \\ c &= 4.6879(5) \\ \beta &= 90.70(1)^\circ \end{aligned}$$

Density

(calculated) 6.893 g/cm^3

Reference intensity

$I/I_{\text{corundum}} = 3.0$

Polymorphism

The monoclinic, wolframite type reported here is stable below 1165°C . Chang et al. [1966] reported a high temperature modification stable above 1165°C . Their data are given on PDF card 19-776. Dunning et al., [1947] reported the existence of a cubic modification, formed between 90° and 300°C .

Additional patterns

1. PDF card 7-190 [Swanson et al. 1953].
2. Broch [1929].
3. Dunning and Megaw [1946].
4. Fonda [1944].

References

- Broch, E. (1929). Skrifter Norske Videns. - Akad., Oslo I. Mat. Nat. Klasse 1929, No. 8.
- Chang, L. L. Y., Scroger, M. G., and Phillips, B. (1966). J. Amer. Ceram. Soc. 49, 385.
- Dunning, N. J. and Megaw, H. D., (1946). Trans. Faraday Soc. 42, 705.
- Fonda, G. R. (1944). J. Phys. Chem. 48, 303.
- Swanson, H. E. and Tatge, E. (1953). Nat. Bur. Stand., U.S. Circ. 539, Vol. I, 84.

CuK α_1 $\lambda = 1.540598\text{\AA}$; temp. $25 \pm 1^\circ\text{C}$			
Internal standard W, $a = 3.16524\text{\AA}$			
d(\AA)	I	hkl	$2\theta(^\circ)$
5.67	20	010	15.63
4.68	95	001	18.94
3.719	100	110	23.91
3.610	45	011	24.64
2.929	100	$\bar{1}11$	30.50
2.901	95	111	30.80
2.836	25	020	31.52
2.463	40	200	36.45
2.459	40	120	36.51
2.427	18	021	37.01
2.343	20	002	38.39
2.261	3	210	39.84
2.191	25	$\bar{2}01$	41.16
2.185	14	$\bar{1}21$	41.28
2.172	40	121, 201	41.55
2.044	6	$\bar{2}11$	44.28
2.027	11	211	44.68
1.9919	13	$\bar{1}12$	45.50
1.9751	18	112	45.91
1.8913	8	030	48.07
1.8600	13	220	48.93
1.8068	12	022	50.47
1.7660	3	130	51.72
1.7540	25	031	52.10
1.7346	6	$\bar{2}21$	52.73
1.7243	7	221	53.07
1.7087	15	$\bar{2}02$	53.59
1.7020	17	$\bar{1}22$	53.82
1.6909	20	122	54.20
1.6881	25	202	54.30
1.6552	2	$\bar{1}31$	55.47
1.6508	2	131	55.63
1.6360	2	$\bar{2}12$	56.18
1.6180	2	212	56.86
1.5782	7	310	58.43
1.5626	4	003	59.07
1.5061	3	013	61.52
1.5011	18	$\bar{3}11, 230$	61.75
1.4904	10	311	62.24
1.4720	7	032	63.11
1.4643	6	$\bar{2}22$	63.48
1.4508	6	222	64.14
1.4458	11	$\bar{1}13$	64.39
1.4360	14	113	64.88
1.4327	17	$\bar{2}31$	65.05
1.4264	12	231	65.37
1.4222	15	320	65.59
1.3690	3	023	68.48
1.3641	13	$\bar{3}21, 140$	68.76
1.3565	4	321	69.20

Magnesium tungsten oxide, MgWO_4

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.3271	2	$\bar{2}03$	70.96
1.3224	6	$\bar{1}23$	71.25
1.3160	4	$\bar{3}12$	71.65
1.3121	7	203	71.90
1.3102	8	$\bar{1}41$	72.02
1.3019	4	312	72.55
1.2923	2	$\bar{2}13$	73.17
1.2786	2	$\bar{2}13$	74.09
1.2681	4	$\bar{2}32$	74.81

Mercury chloride, HgCl_2

Sample

The sample was commercially prepared mercuric chloride.

Color

Colorless

Optical data

Biaxial(-), $N_\alpha = 1.725$, $N_\beta = 1.859$, $N_\gamma = 1.965$,
 $2V = 85^\circ$ [Merwin, 1920].

Structure

Orthorhombic, Pmnb (62), $Z = 4$. The structure was determined by Braekken and Scholten [1934].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 5.9756(8) \text{ \AA} \\ b &= 12.768(2) \\ c &= 4.3347(6) \end{aligned}$$

Density

(calculated) 5.453 g/cm^3

Reference intensity

$$I/I_{\text{corundum}} = 3.2$$

Additional pattern

1. PDF card 4-331 [Swanson and Tatge, 1953].

References

Braekken, H. and Scholten, W. (1934). *Z. Krist.* 89, 448.
 Merwin, H.E. (1920). *J. Am. Chem. Soc.* 42, 2432.
 Swanson, H. E. and Tatge, E. (1953). *Nat. Bur. Stand. (U.S.) Circ.* 539, 1, 73.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1^\circ \text{C}$			
Internal standard W, $a = 3.16524 \text{ \AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
4.365	100	120	20.33
4.107	40	011	21.62
3.587	7	021	24.80
3.511	5	101	25.35
3.386	10	111	26.30
3.192	12	040	27.93
3.075	12	121	29.01
3.038	30	031	29.38
2.989	50	200	29.87
2.708	40	131, 220	33.05

$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
2.418	20	211	37.16
2.361	4	141	38.08
2.295	3	221	39.22
2.199	8	051	41.01
2.181	6	240	41.36
2.130	14	231, 060	42.41
2.065	16	151	43.81
2.012	10	112	45.01
2.006	16	160	45.16
1.941	13	122	46.76
1.902	7	320	47.78
1.838	2	132	49.55
1.820	1	161	50.09
1.810	1	301	50.38
1.793	10	042, 311	50.90
1.771	4	251	51.56
1.754	3	202	52.09
1.740	2	321	52.54
1.738	2	212	52.61
1.681	3	071	54.55
1.666	5	331	55.09
1.653	2	052	55.55
1.621	4	232	56.73
1.618	3	171	56.85
1.595	4	080	57.74
1.574	1	341	58.61
1.537	5	242	60.15
1.493	3	400	62.10
1.477	3	351	62.87
1.472	4	162	63.09
1.454	4	420, 360	63.98
1.436	1	013	64.89
1.429	4	322	65.22
1.408	3	280	66.32
1.4043	5	103, 411	66.53
1.3958	2	113, 072	66.99
1.3788	<1	421, 361	67.93
1.3538	1	262	69.36
1.3406	2	431	70.14
1.3151	1	191	71.71
1.2939	1	213	73.07
1.2850	2	082, 371	73.66
1.2745	1	223	74.37
1.2360	1	451	77.10
1.2304	2	153, 402	77.52
1.1805	1	282	81.46
1.1748	1	520	81.94
1.1694	1	303	82.40
1.1648	1	313	82.80
1.1478	2	442, 511	84.31

Mercury chloride (calomel), Hg_2Cl_2

Sample

The sample was obtained from British Drug House, Ltd.

Optical data

Uniaxial(+), $N_o = 2.6559$, $N_e = 1.97325$ [Groth, 1904].

Structure

Tetragonal, $I4/mmm$ (139), $Z = 2$, isostructural with Hg_2Br_2 , Hg_2F_2 , and Hg_2I_2 [Havighurst, 1925 and Mark and Steinback, 1926].

NBS lattice constants of this sample:

$$a = 4.4801(2)\text{\AA}$$

$$c = 10.9060(6)$$

Density

(calculated) 7.162 g/cm^3

Reference intensity

$I/I_{\text{corundum}} = 5.0$

Additional patterns

1. PDF card 4-581 [Swanson and Tatge, 1953].
2. Havighurst [1925].
3. Hylleraas [1926].
4. Ruff et al. [1928].
5. Hanawalt et al. [1938].

References

- Hanawalt, J. D., Rinn, H. W., and Frevel, L. K. (1938). Ind. Eng. Chem. Anal. Ed. 10, 457.
- Havighurst, R.J. (1925). Amer. J. Sci. 10, 15.
- Hylleraas, E. (1926). Z. Phys. 36, 859.
- Groth, H. (1904). Chemische Krystallographie, Vol. 1, 124, Engelmann, Leipzig.
- Mark, H. and Steinbach, J. (1926). Z. Krist. 64, 79.
- Ruff, O., Ebert, F., and Luft, F. (1928). Z. Anorg. Allg. Chem. 170, 49.
- Swanson, H.E. and Tatge, E. (1953). NBS Circular 539, 1, 72.

CuK α_1 $\lambda = 1.540598\text{\AA}$; temp. $25 \pm 1\text{ }^\circ\text{C}$			
Internal standard Ag, $a = 4.08651\text{\AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
4.147	75	101	21.41
3.170	100	110	28.13
2.824	12	103	31.66
2.727	30	004	32.81
2.240	15	200	40.22
2.067	40	114	43.76
1.970	17	211	46.03
1.962	30	105	46.24
1.818	<1	006	50.14
1.756	4	213	52.05
1.732	12	204	52.83
1.5841	6	220	58.19
1.4755	11	215	62.94
1.4164	3	310	65.89
1.3815	1	303	67.78
1.3696	6	224	68.45
1.3633	3	008	68.81
1.2569	4	314	75.59
1.2522	5	118	75.93
1.2343	2	321	77.23
1.2319	2	305	77.41
1.1756	1	323	81.88
1.1697	5	109	82.38
1.1648	3	208	82.80
1.1202	<1	400	86.89
1.0908	<1	0•0•10	89.85
1.0800	2	325	91.00
1.0563	1	330	93.65
1.0410	1	413	95.46
1.0370	3	219, 332	95.95
1.0312	2	1•1•10	96.66
1.0018	1	420	100.52
0.9846	2	334	102.95
.9823	2	318	103.29
.9728	2	415	104.72
.9405	1	424	109.97
.9089	1	0•0•12	115.89
.8930	<1	431	119.22
.8736	1	1•1•12	123.72
.8675	2	329	125.23
.8287	1	435	136.71
.8246	2	1•0•13	138.20

Nickel acetate hydrate, $\text{Ni}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 4\text{H}_2\text{O}$

Sample

The sample was prepared by slow evaporation at room temperature of an aqueous solution of $\text{Ni}(\text{C}_2\text{H}_3\text{O}_2)_2$.

Color

Brilliant bluish green.

Optical data

Biaxial (-). $N_\alpha = 1.441$, $N_\gamma = 1.560$. 2V is very small.

Structure

Monoclinic $P2_1/c(14)$, $Z = 2$, isostructural with $\text{Co}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 4\text{H}_2\text{O}$. The structure was determined by van Niekerk and Schoening [1953] and refined by Downie et al. [1971].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 4.7749(9) \text{ \AA} \\ b &= 11.772(2) \\ c &= 8.435(1) \\ \beta &= 93.86(1)^\circ \end{aligned}$$

Density

(calculated) 1.747 g/cm^3

Reference intensity

$$I/I_{\text{corundum}} = 4.6$$

Additional patterns

1. PDF card 14-721 [Hanawalt et al., 1938].
2. PDF card 24-1360. This is data from card 14-721 indexed by University College, Cardiff, Wales.

References

- Downie, T. C., Harrison, W., Rafer, E. S., and Hepworth, M. A. (1971). Acta Crystallogr. B27, 706.
- Hanawalt, J. D., Rinn, H. W., and Frevel, L. K. (1938). Ind. Eng. Chem. Anal. Ed. 10, 457.
- van Niekerk, J. N. and Schoening, F. R. (1953). Acta Crystallogr. 6, 609.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1^\circ \text{C}$			
Internal standard W, $a = 3.16524 \text{ \AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
6.84	100	011	12.94
5.886	4	020	15.04
4.828	6	021	18.36
4.762	30	100	18.62
4.416	2	110	20.09
4.209	13	002	21.09
4.014	19	$\bar{1}11$	22.13
3.962	5	012	22.42
3.811	7	111	23.32
3.702	<1	120	24.02
3.555	11	031	25.03
3.454	<1	$\bar{1}21$	25.77
3.326	2	121	26.78
3.265	2	$\bar{1}02$	27.29
3.147	20	$\bar{1}12$	28.34
3.053	3	102	29.23
3.029	2	130	29.47
2.956	4	112	30.21
2.946	2	040	30.32
2.890	8	$\bar{1}31$	30.92
2.869	7	032	31.15
2.813	1	131	31.79
2.711	5	122	33.02
2.531	1	023	35.44
2.504	8	140	35.83
2.438	2	$\bar{1}13$	36.83
2.410	3	042,132	37.28
2.383	2	200	37.72
2.378	4	141	37.80
2.304	7	113	39.07
2.294	7	$\bar{1}23$	39.24
2.281	2	033	39.47
2.267	1	051	39.70
2.186	5	$\bar{1}42,123$	41.27
2.136	1	202	42.27
2.111	5	150	42.81
2.104	4	004,221, $\bar{1}33$	42.95
2.063	3	$\bar{1}51$	43.85
2.055	2	052	44.04
2.032	1	151,043	44.56
2.016	3	202,133	44.92
2.007	4	$\bar{2}22$	45.13
1.982	1	024	45.75
1.953	1	231	46.46
1.948	2	$\bar{1}14$	46.59
1.907	3	222	47.64
1.879	2	104	48.41
1.875	2	$\bar{2}32$	48.51
1.872	2	$\bar{1}24$	48.61
1.855	6	$\bar{2}13,114,034$	49.06

Nickel acetate hydrate, $\text{Ni}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 4\text{H}_2\text{O}$ - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.852	5	240	49.16
1.836	1	143	49.61
1.803	2	053	50.57
1.791	3	$\bar{2}23$	50.95
1.789	2	124,241	51.01
1.778	3	062	51.36
1.7645	<1	161, $\bar{1}$ 34	51.77
1.7392	2	213	52.58
1.7282	1	$\bar{2}42$	52.94
1.7108	<1	044, $\bar{1}$ 53	53.52
1.6944	2	$\bar{2}33$,134	54.08
1.6812	<1	$\bar{1}62$	54.54
1.6660	2	015	55.08
1.6629	2	242,153	55.19
1.6272	1	251	56.51
1.6074	2	063, $\bar{1}$ 15	57.27
1.6043	1	233	57.39

Potassium lead chloride, KPb_2Cl_5

Sample

The sample was prepared by melting a 1:2 molar mixture of KCl and PbCl_2 at 480 °C and cooling in air.

Color

Colorless

Structure

Orthorhombic, $Z=4$, isostructural with RbPb_2Cl_5 and other similar compounds [Jansen, 1968].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 8.865(2) \text{ \AA} \\ b &= 12.498(2) \\ c &= 7.934(1) \end{aligned}$$

Density

(calculated) 4.767 g/cm³

Additional pattern

1. PDF card 23-484 [Jansen, 1968].

Reference

Jansen, P. W. J. (1968). Rec. Trav. Chim. Pays-Bas, 87, 1021.

$\text{CuK}\alpha_1$ $\lambda = 1.540598 \text{ \AA}$; temp. 25±1 °C

Internal standard Ag , $a = 4.08651 \text{ \AA}$

d(Å)	I	hkl	2θ(°)
8.83	25	100	10.01
6.69	11	011	13.22
6.25	5	020	14.16
5.90	7	101	15.00
5.34	11	111	16.58
5.10	10	120	17.37
4.292	7	121	20.68
3.968	25	002	22.39
3.693	100	211,031	24.08
3.616	40	102,220	24.60
3.478	8	112	25.59
3.406	9	131	26.14
3.350	2	022	26.59
3.290	3	221	27.08
3.129	5	122	28.50
3.478	8	112	25.59
3.406	9	131	26.14
3.350	2	022	26.59
3.290	3	221	27.08
3.129	5	122	28.50

d(Å)	I	hkl	2θ(°)
3.123	3	040	28.56
2.952	4	202,300	30.25
2.907	4	041	30.73
2.876	11	310,032	31.07
2.836	6	231	31.52
2.764	9	141	32.36
2.733	9	132	32.74
2.703	<2	311	33.11
2.671	50	222,320	33.52
2.587	5	013	34.65
2.553	20	240	35.12
2.534	9	103	35.39
2.483	2	113	36.14
2.454	2	042	36.59
2.371	4	302	37.91
2.350	14	123	38.27
2.328	7	312	38.65
2.303	10	151	39.09
2.232	18	033	40.38
2.216	16	400,322	40.68
2.166	6	133	41.67
2.148	8	242,340	42.04
2.101	19	251	43.02
2.083	7	060	43.41
2.059	5	332	43.94
2.027	3	160	44.66
1.994	3	233	45.44
1.968	7	143	46.08
1.935	9	104,402	46.92
1.912	5	114,412	47.51
1.900	13	431	47.84
1.879	6	323	48.40
1.856	<2	351	49.04
1.848	5	422	49.27
1.845	9	062	49.36
1.817	<2	053	50.18
1.810	5	204	50.38
1.805	3	162	50.51
1.781	4	333	51.24
1.762	<2	441	51.84
1.756	3	510,134	52.04
1.740	3	224	52.56
1.680	2	253	54.58
1.6679	3	521	55.01
1.6467	2	304	55.78
1.6330	<2	314	56.29
1.6232	2	451	56.66
1.6190	2	502	56.82
1.6092	3	163	57.20
1.5730	5	433	58.64
1.5624	2	080,105	59.08
1.5314	3	334	60.40
1.5177	4	460	61.00
1.5155	5	125	61.10

Potassium platinum chloride, K_2PtCl_6

Sample

The sample was prepared by reaction of KCl and H_2PtCl_6 .

Major impurities

0.01 to 0.1% Na and Ba
0.001 to 0.01% Al, Ca, Cr, and Si
0.0001 to 0.001% Ag, Fe, Mg, and Mn

Color

Bright yellow

Optical data

Isotropic, $N = 1.823$

Structure

Cubic, $Fm\bar{3}m$ (225) $Z=4$, isostructural with other similar alkali platinum halides. The structure of K_2PtCl_6 was determined by Ewing and Pauling [1928].

NBS lattice constant of this sample:

$$a = 9.7560(1) \text{ \AA}$$

Density

(calculated) 3.478 g/cm^3

Reference intensity

$I/I_{\text{corundum}} = 5.7$

Additional patterns

1. PDF card 7-199 [Swanson et al., 1955].
2. Hanawalt et al. [1938].

References

- Ewing, F.J. and Pauling, L. (1928). Z. Krist. 68, 223.
Hanawalt, J. D., Rinn, H. W., and Frevel, L. K. (1938). Ind. Eng. Chem. Anal. Ed. 10, 457.
Swanson, H. E., Gilfrich, N.T., and Ugrinic, G.M. (1955). Nat. Bur. Stand. (U.S.) Circ. 539, 5, 49.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1 \text{ }^\circ\text{C}$			
Internal standard Ag, $a = 4.08651 \text{ \AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
5.633	100	111	15.72
4.878	40	200	18.17
3.4491	45	220	25.81
2.9417	45	311	30.36
2.8160	5	222	31.75
2.4391	40	400	36.82
2.2383	13	331	40.26
2.1817	15	420	41.35
1.9915	14	422	45.51
1.8773	14	511	48.45
1.7246	18	440	53.06
1.6492	13	531	55.69
1.6259	6	600	56.56
1.5425	5	620	59.92
1.4878	4	533	62.36
1.4083	4	444	66.32
1.3659	6	711	68.66
1.3531	3	640	69.40
1.3035	4	642	72.45
1.2700	4	731	74.68
1.2194	1	800	78.35
1.1918	1	733	80.53
1.1829	2	820	81.26
1.1496	1	660	84.14
1.1264	2	751	86.29
1.0907	3	840	89.86
1.0708	2	911	92.00
1.0645	2	842	92.71
1.0401	1	664	95.57
1.0226	2	931	97.75
0.9957	2	844	101.36
.9805	2	933	103.56
.9757	1	10•0•0	104.28
.9567	2	10•2•0	107.25
.9431	2	951	109.49
.9098	1	953	115.71
.9059	2	10•4•0	116.50
.8905	1	10•4•2	119.76
.8796	1	11•1•1	122.26
.8624	1	880	126.57
.8525	1	11•3•1	129.28
.8492	1	10•4•4	130.22
.8366	1	10•6•0	134.08
.8275	1	11•3•3	137.14
.8130	1	12•0•0	142.69
.8064	1	11•5•1	146.40
.7912	1	12•2•2	153.60
.7835	<1	11•5•3	158.90

Silicon, Si

Sample

The sample was very pure vacuum floated, zone refined silicon. This sample is NBS Standard Reference Material # 640, Silicon Powder, X-Ray Diffraction Standard.*

Major impurities (after grinding of the sample):

0.001-0.0001% each of Ca, Cu.

Color

Gray

Structure

Cubic, Fd3m (227), Z = 8 [Debye and Scherrer, 1916].

NBS lattice constant of this sample:

$$a = 5.43088(4)\text{\AA}$$

Density

(calculated) 2.329 g/cm³

Reference intensity

$$I/I_{\text{corundum}} = 4.7$$

Polymorphism

Kasper and Richards [1964] reported that a second, dense form with space group Ia3(206) is formed under pressure.

Additional pattern

1. PDF card 5-565 [Swanson and Fuyat, 1953].
The Swanson and Fuyat [1953] reference lists a large number of early powder patterns.

References

- Debye, P. and Scherrer, P. (1916). Phys. Z. 17, 277.
Kasper, J. S. and Richards, S. M. (1964). Acta Crystallogr. 17, 752.
Swanson, H. E. and Fuyat, R. K. (1953). Nat. Bur. Stand. (U.S.) Circ. 539, 2, 6.

* Samples may be obtained from the Office of Standard Reference Materials, Room B311, Chemistry Building, National Bureau of Standards, Washington, D. C. 20234, \$52 per 10 gram unit.

CuK α_1 λ = 1.540598 \AA ; temp. 25 \pm 1 $^{\circ}\text{C}$			
Internal standard W, a = 3.16524 \AA			
d(\AA)	I	hkl	2 θ ($^{\circ}$)
3.13552	100	111	28.443
1.92011	55	220	47.303
1.63747	30	311	56.123
1.35772	6	400	69.131
1.24593	11	331	76.377
1.10857	12	422	88.032
1.04517	6	511	94.954
0.96005	3	440	106.710
.91799	7	531	114.094
.85870	8	620	127.547
.82820	3	533	136.897

Silver carbonate, Ag₂CO₃

Sample

The sample was prepared by precipitation, adding K₂CO₃ to AgNO₃ solution.

Major impurities

0.001 to 0.01%, Al and Si
0.0001 to 0.001%, Ca, Cu, Fe, and Mg

Color

Greenish yellow

Structure

Monoclinic, P2₁(4), Z=2 [Donahue and Helmholz, 1944].

NBS lattice constants of this sample:

a = 4.8510(7) Å
b = 9.544(2)
c = 3.2533(6)
β = 91.96°(2)

Density

(calculated) 6.084 g/cm³

Additional patterns

1. PDF card 12-766 [Swanson et al., 1962].
2. Hanawalt et al [1938].

References

- Donahue, J. and Helmholz, L. (1944). J. Am. Chem. Soc. 66, 295.
Hanawalt, J. D., Rinn, H. W., and Frevel, L. K. (1938). Ind. Eng. Chem. Anal. Ed. 10, 457.
Swanson, H. E., Morris, M.C., Stinchfield, R. P., and Evans, E.H. (1962). Nat. Bur. Stand. (U.S.) Monogr. 25, Sec. 1, 44.

CuKα ₁ λ = 1.540598 Å; temp. 25±1 °C			
Internal standard W, a = 3.16524 Å			
d(Å)	I	hkl	2θ(°)
4.85	15	100	18.29
4.78	35	020	18.56
4.32	30	110	20.52
3.41	2	120	26.14
3.252	3	001	27.40
3.078	8	011	28.99
2.745	60	101	32.60
2.660	100	130,101	33.66
2.561	6	111	35.01
2.423	2	200	37.07
2.385	11	040	37.68
2.381	13	121	37.76
2.351	8	210	38.26
2.322	14	121	38.75
2.275	35	031	39.59
2.161	11	220	41.76
2.041	10	131	44.35
1.976	2	201	45.89
1.935	6	211	46.92
1.929	9	230	47.08
1.912	4	201	47.51
1.875	6	211	48.52
1.801	3	141	50.65
1.777	13	150,141	51.39
1.701	3	240	53.87
1.678	9	231	54.65
1.646	6	051	55.82
1.639	10	231	56.07
1.626	6	002	56.56
1.616	1	300	56.92
1.591	9	060	57.93
1.538	2	022,112	60.11
1.530	3	320	60.44
1.526	3	102	60.64
1.511	2	160	61.29
1.507	3	112	61.47
1.4676	<1	301	63.32
1.4526	<1	122	64.05
1.4500	1	311	64.18
1.4412	2	330	64.62
1.4278	1	301	65.30
1.4115	1	311	66.15
1.3987	5	132	66.83

Silver sulfate, Ag₂SO₄

Sample

The sample was obtained from J. T. Baker Chemical Company.

Major impurities

0.001 to 0.01% each of Al, Fe, Mg, and Si.
0.0001 to 0.001% each of Ca and Pb.

Color

Colorless

Optical data

Biaxial (-), $n_\alpha = 1.756$, $n_\beta = 1.775$, and $n_\gamma = 1.782$.

Structure

Orthorhombic, Fddd(70), Z=8, Na₂SO₄ type structure [Herrmann and Ilge, 1931].

NBS lattice constants of this sample:

$a = 10.2699(5) \text{ \AA}$
 $b = 12.7069(7)$
 $c = 5.8181(3)$

Density

(calculated) 5.455 g/cm³

Reference intensity

$I/I_{\text{corundum}} = 2.2$

Additional patterns

- PDF card 7-203 [Swanson et al., 1957].
- Hanawalt et al. [1938].

References

- Hanawalt, J. D., Rinn, H. W., and Frevel, L. K. (1938). Ind. Eng. Chem., Anal. Ed. 10, 457.
Herrmann, K. and Ilge, W. (1931). Z. Krist. 80, 402.
Swanson, H. E., Gilfrich, N. T., and Cook, M. I. (1957). Nat. Bur. Stand. (U.S.) Circ. 539, 7, 46.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1 \text{ }^\circ\text{C}$

Internal standard W, $a = 3.16524 \text{ \AA}$

d(Å)	I	hkl	2 θ (°)
4.699	10	111	18.87
3.994	25	220	22.24
3.249	3	131	27.43
3.177	70	040	28.06
2.873	100	311	31.10
2.644	90	022	33.87
2.568	1	400	34.91
2.530	17	202	35.45
2.421	30	331	37.10
2.352	3	222	38.24
2.272	8	151	39.64
1.980	11	242	45.78
1.957	8	260	46.35
1.926	30	351	47.15
1.915	12	511	47.44
1.884	5	113	48.26
1.762	3	531	51.85
1.7376	3	133	52.63
1.7123	17	062	53.47
1.6730	12	313	54.83
1.6527	4	620	55.56
1.6243	1	262	56.62
1.5881	3	080	58.03
1.5675	13	333	58.87
1.5462	8	371	59.76
1.5404	6	551	60.01
1.4751	4	602	62.96
1.4542	3	004	63.97
1.4057	6	353	66.46
1.3668	3	224	68.61
1.3598	1	191	69.01
1.3457	2	282	69.84
1.3380	6	642,533	70.30
1.3312	6	660	70.71
1.3225	6	044	71.25
1.2837	1	800	73.75
1.2736	3	391	74.43
1.2359	3	373	77.11
1.2331	3	2·10·0,553	77.32
1.1905	1	840	80.64
1.1678	1	264	82.54
1.1645	3	0·10·2	82.83
1.1549	1	822	83.67
1.1408	1	591	84.94
1.1154	1	911,135	87.36
1.1136	1	573	87.53
1.0975	2	315	89.15
1.0919	3	624	89.73
1.0829	4	393,931	90.69
1.0809	3	682	90.90

Silver sulfate, Ag_2SO_4 - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.0758	3	3•11•1	91.46
1.0727	3	084	91.80
1.0663	2	335	92.51
1.0589	1	0•12•0	93.34
1.0272	1	862	97.17
1.0246	1	951	97.49
1.0203	1	6•10•0	98.05
1.0138	1	10•2•0	98.89
1.0109	1	355	99.28
0.9976	1	593	101.10
.9920	1	5•11•1	101.88
.9820	2	664	103.33
.9586	1	026	106.95
.9531	1	3•11•3,971	107.84
.9419	1	375	109.74
.9407	1	2•10•4,555	109.94
.9279	1	3•13•1,4•12•2	112.23
.9210	1	844	113.51

Sodium phosphate hydrate, α -Na₄P₄O₁₂·4H₂O

Sample

The sample was prepared by H.M. Ondik by hydrolytic cleavage of the α form of P₂O₅ below 15 °C. The material was neutralized by NaOH, then purified by salting out with NaCl, followed by repeated recrystallizations with H₂O and ethanol.

Major impurities

0.001 to 0.01% each of Ba, Ca, Si, and Sr.

Color

Colorless

Optical data

Biaxial(+), $N_{\alpha} = 1.440$, $N_{\beta} = 1.458$, $N_{\gamma} = 1.476$.

Structure

Monoclinic, P2₁/a (14), Z=2. The structure of α -Na₄P₄O₁₂·4H₂O was determined by Ondik et al. [1961].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 9.691(2) \text{ \AA} \\ b &= 12.342(2) \\ c &= 6.187(2) \\ \beta &= 92.58(1)^{\circ} \end{aligned}$$

Density

(calculated) 2.156 g/cm³

Polymorphism

Thilo and Ratz [1949] reported a β , high temperature form of Na₄P₄O₁₂·4H₂O.

Additional patterns

1. PDF card 11-15 [Swanson et al., 1960].
2. Bell et al. [1952].
3. Thilo and Ratz [1949].

References

- Bell, R.N., Audrieth, L.F., and Hill, O.F. (1952). Ind. Eng. Chem. 44, 568.
- Ondik, H. M., MacGillavry, C .H., and Block, S. (1961). Acta Crystallogr. 14, 555.
- Swanson, H. E., Cook, M. I., Evans, E. H., and de Groot, J.H. (1960). Nat. Bur. Stand. (U.S.) Circ. 539, 10, 52.
- Thilo, E. and Ratz, R. (1949). Z. Anorg. Allg. Chem. 260, 255.

CuK α_1 $\lambda = 1.540598$ Å; temp. 25 \pm 1 °C			
Internal standard W, $a = 3.16524$ Å			
d (Å)	I	hkl	2 θ (°)
7.64	75	110	11.58
6.17	90	001,020	14.34
5.21	5	120	17.00
4.844	65	200	18.30
4.719	60	111	18.79
4.510	3	210	19.67
4.369	17	021	20.31
3.933	20	121	22.59
3.894	2	$\bar{2}$ 01	22.82
3.805	95	220	23.36
3.728	25	201	23.85
3.572	8	211	24.91
3.424	10	031	26.00
3.295	100	$\bar{2}$ 21	27.04
3.255	70	131	27.38
3.194	20	221	27.91
3.133	19	230	28.47
3.122	19	310	28.57
3.089	25	002,040	28.88
2.938	9	140	30.40
2.827	70	$\bar{2}$ 31,112	31.62
2.760	19	022,041 +	32.41
2.739	20	311	32.67
2.685	18	$\bar{1}$ 22	33.34
2.669	7	$\bar{1}$ 41	33.55
2.638	14	141	33.95
2.633	15	321	34.02
2.602	7	240, $\bar{2}$ 12	34.44
2.554	14	202	35.11
2.539	35	330	35.32
2.418	18	$\bar{2}$ 41, $\bar{1}$ 32	37.16
2.392	8	150	37.57
2.375	8	410,132	37.85
2.360	5	222	38.10
2.293	5	051	39.26
2.253	40	420	39.98
2.247	25	$\bar{3}$ 12	40.10
2.222	16	151	40.56
2.199	11	250	41.01
2.185	16	411,042	41.29
2.170	5	232	41.59
2.148	5	312	42.02
2.146	3	$\bar{4}$ 21, $\bar{1}$ 42	42.08
2.120	4	341	42.62
2.115	5	142	42.71
2.087	4	430, $\bar{2}$ 51	43.32
2.060	2	003,251	43.92
2.015	8	$\bar{2}$ 42	44.94
2.001	6	$\bar{4}$ 31	45.29
1.996	1	332	45.40

Sodium phosphate hydrate, $\alpha\text{-Na}_4\text{P}_4\text{O}_{12}\cdot 4\text{H}_2\text{O}$ - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.968	8	242	46.08
1.961	10	350	46.25
1.955	8	023, 431	46.40
1.949	8	$\bar{4}02$	46.56
1.928	10	332, 052 +	47.09
1.924	7	$\bar{4}12$	47.19
1.913	6	510	47.49
1.907	10	161	47.65
1.903	14	$\bar{2}13, \bar{1}52$ +	47.75
1.895	9	260	47.97
1.854	18	351	49.11
1.850	18	$\bar{5}11$	49.20
1.842	9	033	49.44
1.837	9	$\bar{4}41, \bar{3}42$	49.58
1.823	10	$\bar{1}33$	50.00
1.786	8	223, 422	51.11
1.776	8	252	51.42
1.7547	6	$\bar{3}13$	52.08
1.7500	5	521	52.23
1.7456	6	$\bar{2}33$	52.37
1.7349	5	360, 170	52.72
1.7123	3	043, 062	53.47
1.6991	6	233, 432 +	53.92
1.6806	12	$\bar{3}61$	54.56
1.6761	16	143, $\bar{3}52$	54.72

Strontium hydroxide, $\text{Sr}(\text{OH})_2$

Sample

The sample was prepared by heating $\text{Sr}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$ for 24 hours at about 200 °C.

Color

Colorless

Structure

Orthorhombic, Pbnm (62), $Z=4$, [Bärnighausen and Weidlein, 1965]. The structure was determined by Grueninger and Bärnighausen [1969].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 6.1201(6) \text{ \AA} \\ b &= 9.892(1) \\ c &= 3.9193(5) \end{aligned}$$

Density

(calculated) 3.405 g/cm³

Reference intensity

$$I/I_{\text{corundum}} = 2.7$$

Additional patterns

1. PDF card 18-1273 [Bärnighausen and Weidlein, 1965].
2. PDF card 19-1276 [Mercer and Miller, 1966]. This pattern is labeled as anhydrous but is for $\text{Sr}(\text{OH})_2 \cdot \text{H}_2\text{O}$.
3. Berggren and Brown [1971].

References

- Bärnighausen, H. and Weidlein, J. (1965). Acta Crystallogr. **19**, 1048.
- Berggren, G. and Brown, A. (1971). Acta Chem. Scand. **25**, 1377.
- Grueninger, H. W. and Bärnighausen, H (1969). Z. Anorg. Allgem. Chem. **368**, 53.
- Mercer, R. A. and Miller, R.P. (1966). J. Inorg. Nucl. Chem. **28**, 61.

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
3.068	25	021,200	29.08
2.922	25	210	30.57
2.903	45	130	30.78
2.745	5	121	32.59
2.602	2	220	34.44
2.473	25	040	36.30
2.343	45	211	38.38
2.293	10	140	39.26
2.244	2	230	40.16
2.168	10	221	41.62
2.091	25	041	43.23
1.998	3	310	45.36
1.979	6	141	45.81
1.959	20	002	46.31
1.947	50	231	46.62
1.886	17	320	48.21
1.834	6	112	49.68
1.822	3	022	50.03
1.810	15	301	50.38
1.780	1	311	51.29
1.746	6	122	52.35
1.735	1	330	52.73
1.727	2	241	52.97
1.6970	17	151	53.99
1.6615	6	250	55.24
1.6486	6	060	55.71
1.6280	11	212	56.48
1.6245	16	132	56.61
1.5735	5	340	58.62
1.5360	7	042	60.20
1.5304	6	400,251	60.44
1.5126	3	410	61.23
1.4898	4	142	62.27
1.4751	3	232,161	62.96
1.4608	5	341	63.65
1.4518	1	260	64.09
1.4109	1	411	66.18
1.3874	1	430	67.45
1.3768	2	170	68.04
1.3696	2	421	68.45
1.3589	8	322	69.06
1.3353	1	351	70.46
1.3080	3	431	72.16
1.3012	4	440	72.60
1.2992	4	171,332	72.73
1.2829	4	270	73.80
1.2672	8	252,113	74.87
1.2618	5	062	75.25

$\text{CuK}\alpha_1 \lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1 \text{ }^{\circ}\text{C}$

Internal standard W, $a = 3.16524 \text{ \AA}$

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
5.19	55	110	17.06
4.94	25	020	17.94
3.846	40	120	23.11
3.300	40	101	27.00
3.130	100	111	28.49

Strontium hydroxide hydrate $\text{Sr}(\text{OH})_2 \cdot \text{H}_2\text{O}$

Sample

The sample was prepared by heating $\text{Sr}(\text{OH})_2$ in a partly closed tube with about 1 ml H_2O at 100°C for 24 hours. The sample contained a small amount of SrCO_3 . Because of this and the tendency to lose H_2O when exposed to air, the intensities may be slightly in error.

Color

Colorless

Structure

Orthorhombic, $\text{Pb}2_1\text{m}$ (26), $Z = 2$. Isostructural with $\text{Eu}(\text{OH})_2 \cdot \text{H}_2\text{O}$ and $\text{Ba}(\text{OH})_2 \cdot \text{H}_2\text{O}$ [Bärnighausen, 1966]. The structure of $\text{Sr}(\text{OH})_2 \cdot \text{H}_2\text{O}$ was determined by Bärnighausen and Weidlein [1967].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 6.201(1) \text{ \AA} \\ b &= 6.716(1) \\ c &= 3.6483(6) \end{aligned}$$

Density

(calculated) 3.053 g/cm^3

Reference intensity

$$I/I_{\text{corundum}} = 0.8$$

Additional patterns

1. PDF card 19-1276 [Mercer and Miller, 1966]. This pattern was labeled as $\text{Sr}(\text{OH})_2$
2. Bärnighausen [1966].
3. Berggren and Brown [1971].
4. Carlson [1954].
5. Lutz [1965].

References

- Bärnighausen, H. (1966). Z. Anorg. Allgem. Chem. 342, 233.
- Bärnighausen, H., and Weidlein, J. (1967). Acta. Crystallogr. 22, 252.
- Berggren, G., and Brown, A. (1971). Acta. Chem. Scand. 25, 1377.
- Carlson, E. T. (1954). J. Res. Nat. Bur. Stand. 53, 371.
- Lutz, H. D. (1965). Z. Naturforsch. 20b, 61.
- Mercer, R. A., and Miller, R. P., (1966). J. Inorg. Nucl. Chem. 28, 61.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1^\circ\text{C}$			
Internal standard Ag, $a = 4.08651 \text{ \AA}$			
$d(\text{ \AA})$	I	hkl	$2\theta(^\circ)$
6.20	65	100	14.28
4.556	80	110	19.47
3.651	40	001	24.36
3.360	55	020	26.51
3.147	45	101	28.34
2.954	15	120	30.23
2.848	60	111	31.39
2.814	100	210	31.77
2.472	50	021	36.31
2.363	10	201	38.05
2.296	70	121	39.21
2.230	85	211	40.42
2.106	25	130	42.91
2.067	10	300	43.77
1.977	18	310	45.87
1.825	35	002,131	49.94
1.816	25	230	50.21
1.798	7	301	50.72
1.760	11	320	51.90
1.750	7	102	52.24
1.738	12	311	52.63
1.6938	11	112	54.10
1.6792	7	040	54.61
1.6248	25	231	56.60
1.6033	9	022	57.43
1.5854	8	321	58.14
1.5515	5	122	59.53
1.5500	7	400	59.60
1.5311	13	212	60.41
1.5186	6	330	60.96
1.4808	9	141	62.69
1.4266	7	401	65.36
1.4073	7	420	66.37
1.4024	8	331	66.62
1.3784	7	132	67.95
1.3678	3	302	68.55
1.3398	7	312	70.19
1.3124	11	150	71.88
1.3029	5	340	72.49
1.2862	8	232	73.58
1.2664	6	322	74.93
1.2396	4	500	76.83
1.2350	9	042,151	77.18

Strontium hydroxide hydrate, $\text{Sr}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$

Sample

The sample was prepared by treating SrCO_3 with HCl followed by NaOH at boiling temperatures. On cooling, the first crystals formed were removed, redissolved and reprecipitated by slow evaporation at room temperature.

Color

Colorless

Optical Data

Uniaxial (-), $N_o = 1.497$, $N_e = 1.475$

Structure

Tetragonal, $P4/ncc$ (130), $Z = 4$. The structure was determined by Smith [1953].

NBS lattice constants of this sample:

$$a = 9.019(2)\text{\AA}$$

$$c = 11.614(2)$$

Density

(calculated) 1.868 g/cm^3

Reference intensity

$I/I_{\text{corundum}} = 1.3$

Additional patterns

1. PDF card 1-1263 [Hanawalt et al, 1938].
2. PDF card 2-1262 [Natta, 1928].
3. Berggren and Brown [1971].

References

- Berggren, G., and Brown, A. (1971). Acta. Chem. Scand. 25, 1377.
- Hanawalt, J.D., Rinn, R.W., and Frevel, L.K. (1938). Ind. Eng. Chem. Anal. Ed. 10, 457.
- Natta, G. (1928). Gazz. Chim. Ital. 58, 870.
- Smith, H. G. (1953). Acta. Crystallogr. 6, 604.

$\text{CuK}\alpha_1 \lambda = 1.540598 \text{\AA}$; temp. $25 \pm 1 \text{ }^\circ\text{C}$

Internal standard W, $a = 3.16524 \text{\AA}$

$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
6.37	50	110	13.89
5.81	45	002	15.25
4.50	45	200	19.70
4.292	100	112	20.68
3.559	80	202	25.00
3.190	19	220	27.95
2.904	14	004	30.76
2.795	45	222, 213	31.99
2.768	60	311	32.32
2.640	40	114	33.93
2.560	55	312	35.02
2.442	40	204	36.78
2.357	6	214	38.15
2.296	25	322, 313	39.21
2.253	20	400	39.99
2.147	30	224	42.06
2.102	18	402, 323	42.99
2.035	65	314	44.48
2.017	55	420	44.91
1.997	15	332	45.36
1.935	6	006	46.92
1.905	20	422, 413	47.70
1.852	12	116	49.15
1.801	11	315	50.63
1.780	7	404	51.28
1.779	15	206	51.33
1.715	9	334	53.37
1.692	15	512	54.15
1.655	8	424, 226	55.46
1.6020	13	316	57.48
1.5106	9	514	61.32
1.4950	8	532	62.03
1.4690	6	406	63.25
1.4518	10	008	64.09
1.4331	8	108	65.03
1.4263	12	620	65.37
1.4157	7	118	65.93
1.3964	10	426	66.96
1.3818	8	208	67.76
1.3648	7	534	68.72

Strontium silicate, Sr_3SiO_5

Sample

The sample was prepared by repeated grindings and heatings of a 3:1 molar mixture of SrCO_3 and silica gel. The temperature was about 1350 °C.

Color

Colorless

Structure

Tetragonal, $P4/ncc$ (130), $Z=4$ [Mansmann, 1965].
The structure of Sr_3SiO_5 was studied by Dent Glasser and Glasser [1965].

NBS lattice constants for this sample:

$$a = 6.9476(3)\text{\AA}$$

$$c = 10.7534(6)$$

Density

(calculated) 4.747 g/cm³

Reference intensity

$I/I_{\text{corundum}} = 3.6$

Additional patterns

1. PDF card 18-1282 [Dear, Bull. Mat. Eng. Exp. Sta., 1957].
2. Eysel [1970].
3. Nurse [1952].

References

- Dent Glasser, L.S. and Glasser, R.P. (1965). Acta Crystallogr. 18, 453.
Eysel, W. (1970). Neues Jahrb. Mineral. Montash. 1970, 534.
Mansmann, M. (1965). Z. Anorg. Allg. Chem. 339, 52.
Nurse, R. W. (1952). J. Appl. Chem. (London) 2, 244.

CuK α_1 $\lambda = 1.540598 \text{\AA}$; temp. 25 \pm 1 °C			
Internal standard W, $a = 3.16524 \text{\AA}$			
d (Å)	I	hkl	2 θ (°)
5.38	4	002	16.46
4.92	1	110	18.02
4.249	1	102	20.89
3.629	10	112	24.51
2.984	30	211	29.92
2.919	100	202	30.60
2.690	30	212, 004	33.29
2.508	2	104	35.78
2.458	30	220	36.53
2.348	55	213	38.30
2.198	25	310	41.04
2.035	1	312, 214	44.49
1.897	4	321	47.92
1.814	20	322, 224	50.27
1.793	2	006	50.89
1.7683	9	215	51.65
1.7550	1	304	52.07
1.7370	1	400	52.65
1.7014	9	314	53.84
1.6840	3	116	54.44
1.6646	17	411	55.13
1.6530	3	402	55.55
1.6376	2	330	56.12
1.5926	16	206	57.85
1.5662	10	332, 324	58.92
1.5535	4	420	59.45
1.5366	1	421, 315	60.17
1.5250	12	413	60.68
1.4926	4	422	62.14
1.4589	4	404	63.74
1.4352	2	325	64.92
1.3982	1	334	66.86
1.3889	1	316	67.37
1.3772	4	217	68.02
1.3446	7	424, 008	69.90
1.3262	7	415	71.02
1.3208	5	512	71.35
1.2961	1	118, 433	72.93
1.2810	1	521	73.93
1.2571	2	406	76.29
1.2338	1	218	77.27
1.2279	1	440, 416	77.71
1.2140	3	523	78.77
1.2090	3	336	79.16
1.2012	1	327	79.77
1.1974	2	442	80.08
1.1913	3	530	80.57
1.1790	4	228	81.59
1.1739	2	426	82.02
1.1674	2	435	82.58

Strontium silicate, Sr_3SiO_5 - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.1632	3	532,524 +	82.94
1.1579	2	600	83.40
1.1468	1	318	84.40
1.1352	6	417	85.46
1.1154	3	219	87.36
1.0891	7	534	90.03
1.0846	4	516	90.51
1.0761	4	622	91.42
1.0632	5	604,408	92.85
1.0504	<1	1•1•10,623 +	94.33
1.0388	1	338	95.72
1.0305	1	437	96.75
1.0272	3	2•0•10	97.16
1.0161	2	428,2•1•10	98.59
1.0133	1	446	98.96
1.0088	2	615	99.56
0.9922	1	536	101.85
.9747	2	419	104.43
.9686	1	545	105.36
.9662	2	438,3•1•10 +	105.74
.9597	<1	641	106.77
.9506	1	721	108.26
.9484	2	642	108.63
.9364	2	626	110.69
.9324	2	2•1•11	111.40
.9305	1	643	111.75

Tin hydrogen phosphate, SnHPO_4

Sample

The sample was made by T. H. Jordan of the American Dental Association Health Foundation. Tin Fluoride, SnF_2 , was treated with H_3PO_4 at a pH of 2, followed by slight heating.

Color

Colorless

Structure

Monoclinic, $P2_1/a(14)$, $Z = 4$. The structure was determined by Berndt and Lamberg [1971].

NBS lattice constants of this sample:

$$\begin{aligned} a &= 5.8307(8) \text{ \AA} \\ b &= 13.617(1) \\ c &= 4.6145(6) \\ \beta &= 98.73(1)^\circ \end{aligned}$$

Density

(calculated) 3.937 g/cm^3

Reference Intensity

$$I/I_{\text{corundum}} = 2.0.$$

Reference

Berndt, A. F. and Lamberg, R. (1971). Acta Crystallogr. B27, 1092.

CuK α_1 $\lambda = 1.540598 \text{ \AA}$; temp. $25 \pm 1^\circ \text{C}$			
Internal standard Ag, $a = 4.08651 \text{ \AA}$			
d (Å)	I	hkl	2 θ (°)
6.80	100	020	13.01
5.311	2	110	16.68
4.562	12	001	19.44
4.401	11	120	20.16
4.327	1	011	20.51
3.792	12	021	23.44
3.731	16	111	23.83
3.566	6	130	24.95
3.404	35	040	26.16
3.367	4	121	26.45
3.241	8	111	27.50
2.997	30	121	29.79
2.946	40	131	30.31
2.882	1	200	31.01
2.820	14	210	31.71
2.729	4	041	32.79
2.689	7	131	33.29
2.654	1	220	33.75
2.623	1	201	34.16
2.574	4	211	34.82
2.559	1	141	35.04
2.448	1	221	36.68
2.384	3	141	37.71
2.338	1	051	38.47
2.279	6	002	39.51
2.269	11	231,060	39.69
2.251	10	211,012	40.02
2.228	3	151	40.45
2.210	4	112	40.79
2.163	<1	022	41.72
2.111	1	160,151	42.80
2.077	1	241	43.53
2.032	10	061	44.55
2.008	6	132	45.11
1.979	9	250	45.82
1.936	3	202,122	46.90
1.918	2	212	47.36
1.894	6	042	47.99
1.889	6	251	48.12
1.878	2	161	48.44
1.863	1	222	48.84
1.848	3	320	49.28
1.843	4	132,170	49.41
1.808	4	321	50.43
1.789	1	071	51.01
1.783	1	260	51.19
1.769	2	330	51.62
1.748	5	052	52.28
1.737	6	171	52.64
1.681	2	171	54.53

Tin hydrogen phosphate, SnHPO_4 - continued

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
1.669	1	202, 311	54.96
1.657	2	212	55.42
1.643	3	$\bar{3}41$	55.93
1.632	3	180, 321	56.32
1.609	3	062	57.22
1.578	2	$\bar{2}52$	58.44
1.5585	1	$\bar{1}81$	59.24
1.5514	1	322	59.54
1.5166	4	181	61.05
1.5081	2	162, 341	61.43
1.4989	1	242	61.85
1.4802	1	072	62.72
1.4730	2	$\bar{2}62$	63.06
1.4634	1	$\bar{1}90$	63.52
1.4486	2	$\bar{1}33$	64.25
1.4432	1	$\bar{3}42$	64.52
1.4384	2	$\bar{4}01, \bar{2}03$	64.76
1.4303	1	351, $\bar{4}11$ +	65.17
1.4231	2	252	65.54
1.4092	2	420, $\bar{1}91$	66.27
1.3943	1	$\bar{1}43$	67.07
1.3878	3	043, 123	67.43
1.3781	1	191	67.97
1.3738	1	430, $\bar{2}72$	68.21
1.3670	1	370	68.59
1.3615	1	0•10•0	68.91

Zinc borate, $\text{Zn}_4\text{B}_6\text{O}_{13}$

Sample

The sample was a phosphor preparation obtained from the Radio Corporation of America [Leverenz, 1944].

Color

Colorless

Structure

Cubic, $I\bar{4}3m(217)$, $Z=2$. The structure was determined by Smith et al. [1961], who found the formula to be $\text{Zn}_4\text{O}(\text{BO}_2)_6$.

NBS lattice constant of this sample:

$$a = 7.4734(2)\text{\AA}$$

Density

(calculated) 4.252 g/cm^3

Additional pattern

1. PDF card 14-2 [Swanson and Tatge, 1953]. The formula at that time was mistakenly given as ZnB_2O_4 .

References

- Leverenz, H.W. (1944). Proc. I.R.E. 32, 256.
 Smith, P., García-Blanco, S., and Rivoir, L. (1961). An. Reál Soc. Españ. Fis. Quim. Madrid, A57, 263.
 Swanson, H. E. and Tatge, E. (1953). Nat. Bur. Stand. (U.S.) Circ. 539, 1, 83.

CuK α_1 $\lambda = 1.540598\text{ \AA}$; temp. $25\pm 1\text{ }^\circ\text{C}$ Internal standard W, $a = 3.16524\text{ \AA}$			
$d(\text{\AA})$	I	hkl	$2\theta(^\circ)$
5.29	6	110	16.74
3.74	4	200	23.77
3.05	100	211	29.28
2.364	25	310	38.03
2.158	2	222	41.82
1.997	20	321	45.37
1.869	14	400	48.68
1.761	40	330	51.88
1.672	2	420	54.86
1.594	4	332	57.79
1.526	25	422	60.63
1.466	6	510	63.39
1.364	8	521	68.76
1.321	4	440	71.34
1.282	4	530	73.86
1.246	2	600	76.37
1.213	2	611	78.84
1.1818	2	620	81.36
1.1532	4	541	83.82
1.1026	2	631	88.64
1.0789	2	444	91.12
1.0569	2	710	93.58
1.0366	2	640	96.00
1.0170	4	721	98.48
0.9992	2	642	100.88
.9813	2	730	103.44
.9491	2	732	108.51
.9199	4	811	113.74
.9063	2	820	116.42
.8933	2	653	119.16
.8808	2	822	121.99
.8688	4	831	124.92
.8574	2	662	127.91
.8463	2	752	131.08
.8253	2	910	137.95
.8154	2	842	141.73
.8059	2	921	145.83
.7967	2	664	150.44
.7878	2	930	155.83

Zinc titanium oxide, ZnTiO₃

Sample

The sample was prepared by heating an equimolar mixture of Zn(NO₃)₂ and TiO₂ (anatase) for about two weeks at 900° with several remixings and regrindings. Because of the lack of thermal stability above 943° [Dulin and Rase, 1960], it was impossible to obtain complete reaction and the sample contained small amounts of rutile (TiO₂) and Zn₂TiO₄; therefore there may be a slight error in some intensities. Intensities calculated from the structure were in good agreement with the experimental values.

Color

Colorless

Structure

Hexagonal, R $\bar{3}$ (148), Z=6. ZnTiO₃ is isostructural with FeTiO₃ (ilmenite) and other similar titanates [Bartram and Slepety's, 1961].

NBS lattice constants of this sample:

$$a = 5.0787(3)\text{\AA}$$

$$c = 13.927(1)$$

Density

(calculated) 5.165 g/cm³

Reference intensity

I/I_{corundum} = 2.5

Additional patterns

1. Bartram and Slepety's [1961].
2. Kubo and Kato [1963].

References

- Bartram, S.F. and Slepety's, R.A. (1961). J. Amer. Ceram. Soc. 44, 493.
Dulin, F.H. and Rase, D.E. (1960). J. Amer. Ceram. Soc. 43, 125.
Kubo, T. and Kato, M. (1963). Kogyo Kagaku Zasshi 66, 404.

CuK α_1 λ = 1.540598 \AA ; temp. 25 \pm 1 °C			
Internal standard Ag, a = 4.08651 \AA			
d (\AA)	I	hkl	2 θ (°)
4.63	1	003	19.14
4.191	3	101	21.18
3.717	20	012	23.92
2.729	100	104	32.79
2.540	75	110	35.31
2.355	1	015	38.18
2.321	1	006	38.76
2.228	20	113	40.45
2.173	3	021	41.52
2.097	1	202	43.10
1.860	35	024	48.94
1.813	1	107	50.29
1.713	35	116	53.43
1.651	1	211	55.64
1.619	11	018	56.83
1.500	25	214	61.80
1.466	25	300	63.41
1.428	1	125	65.31
1.399	1	303	66.84
1.3650	4	208	68.71
1.3276	8	1•0•10	70.93
1.3218	4	119	71.29
1.2760	1	217	74.27
1.2696	6	220	74.71
1.2396	2	306	76.84
1.2166	1	0•1•11	78.57
1.2020	6	128,312	79.71
1.1766	3	0•2•10	81.79
1.1512	7	134	84.00
1.1139	5	226	87.50
1.0862	1	042	90.33
1.0674	5	2•1•10	92.38
1.0558	2	1•1•12	93.71
1.0485	3	404	94.56
1.0069	<1	1•2•11	99.82
0.9990	3	318	100.90
.9816	<1	229	103.39
.9702	4	0•1•14	105.12
.9692	7	324	105.27
.9599	5	410	106.74
.9296	2	048	111.92
.9175	3	1•3•10	114.19
.9064	2	2•0•14	116.39
.8868	5	416	120.59

Cerium cobalt, CeCo₂

Structure

Cubic, Fd3m(227), Z=8, C15 type, isostructural with Cu₂Mg [Fülling et al., 1942].

Lattice constant: [Wernick and Geller, 1960]

$$a = 7.161(5)\text{\AA}$$

Density

(calculated) 9.333 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Ce⁰ and Co⁰ [Thomas and Umeda, 1957], corrected for dispersion [Dauben and Templeton, 1955].

Scale factors (integrated intensities)

$$\gamma = 0.501 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 10.3$$

Additional pattern

1. Fülling, Moeller and Vogel [1942].

References

- Dauben, C. H. and Templeton, D. H. (1955). Acta Crystallogr. 8, 841.
 Fülling, W., Moeller, K., and Vogel, R. (1942). Z. Metallk. 34, 253.
 Thomas, L.H. and Umeda, K. (1957). J. Chem. Phys. 26, 293.
 Wernick, J.H. and Geller, S. (1960). Trans. AIME 218, 866.

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
4.13	12	1	1	1	21.48
2.532	66	2	2	0	35.43
2.159	100	3	1	1	41.80
2.067	15	2	2	2	43.76
1.643	3	3	3	1	55.92
1.462	20	4	2	2	63.60
1.378	19	5	1	1	67.97
1.378	6	3	3	3	67.97
1.266	18	4	4	0	74.96
1.210	2	5	3	1	79.05
1.132	8	6	2	0	85.74
1.092	8	5	3	3	89.72
1.080	4	6	2	2	91.05
.957	10	6	4	2	107.21
.932	12	7	3	1	111.43
.932	6	5	5	3	111.43
.895	4	8	0	0	118.76
.844	3	6	6	0	131.78
.844	6	8	2	2	131.78
.827	14	7	5	1	137.36
.827	2	5	5	5	137.36
.821	3	6	6	2	139.36
.786	3	7	5	3	157.04
.786	1	9	1	1	157.04

Calculated Pattern (Peak heights)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
4.13	16	1	1	1	21.48
2.532	68	2	2	0	35.42
2.159	100	3	1	1	41.80
2.067	15	2	2	2	43.76
1.643	3	3	3	1	55.92
1.462	16	4	2	2	63.60
1.378	20	5	1	1 +	67.96
1.266	13	4	4	0	74.96
1.211	1	5	3	1	79.04
1.132	5	6	2	0	85.74
1.092	5	5	3	3	89.72
1.080	2	6	2	2	91.04
1.003	1	7	1	1 +	100.38
.957	6	6	4	2	107.22
.932	9	7	3	1 +	111.44
.895	2	8	0	0	118.76
.844	4	8	2	2 +	131.78
.827	6	7	5	1 +	137.36
.821	1	6	6	2	139.36
.786	1	7	5	3 +	157.04

Cerium cobalt, $\text{Ce}_{24}\text{Co}_{11}$

Structure

Hexagonal, $P6_3mc$ (186), $Z = 2$. The structure was determined by Larson and Cromer [1962].

Lattice constants:

$$a = 9.588 \text{ \AA}$$

$$c = 21.827$$

(published values: $a = 9.587$, $c = 21.825 \text{ \AA}$ [ibid.]).

Density

(calculated) 7.666 g/cm^3

Thermal parameters

Isotropic [Larson and Cromer, 1962].

Scattering factors

Ce^0 , Co^0 [International Tables, 1962].

Scale factors (integrated intensities)

$$\gamma = 0.170 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 3.63$$

References

International Tables for X-ray Crystallography III (1962), 210, 211.

Larson, A. C. and Cromer, D. T. (1962). Acta Crystallogr. 15, 1224.

Calculated Pattern (Peak heights)

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$			
10.91	2	0 0 2	8.10
8.29	1	1 0 0	10.66
6.60	3	1 0 2	13.40
4.79	1	1 1 0	18.50
3.88	1	2 0 2	22.90
3.60	2	2 0 3 +	24.70
3.33	10	1 0 6	26.74
3.30	28	2 0 4	26.96
3.14	32	2 1 0	28.42
3.11	28	2 1 1	28.72
3.02	28	2 1 2	29.60
3.01	23	2 0 5	29.66
2.919	60	1 0 7	30.60
2.899	12	1 1 6	30.82
2.882	19	2 1 3	31.00
2.768	22	3 0 0	32.32
2.745	8	3 0 1	32.60
2.736	22	2 0 6	32.70
2.720	100	2 1 4 +	32.90
2.587	63	3 0 3 +	34.64
2.548	22	2 1 5	35.20
2.493	56	2 0 7	36.00
2.396	56	2 2 0	37.50
2.377	5	2 1 6	37.82
2.338	4	3 0 5	38.48

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$			
2.280	18	2 0 8	39.50
2.253	1	3 1 2	39.98
2.212	2	2 1 7	40.76
2.203	4	3 0 6	40.94
2.196	3	3 1 3	41.06
2.122	8	3 1 4	42.58
2.111	2	1 0 10	42.80
2.094	2	2 0 9	43.16
2.066	3	4 0 1	43.78
2.059	4	2 1 8	43.94
2.039	1	4 0 2	44.40
1.996	1	4 0 3	45.40
1.946	2	3 1 6	46.64
1.940	3	4 0 4 +	46.78
1.932	2	2 0 10 +	47.00
1.919	3	2 1 9	47.34
1.905	1	3 2 0	47.70
1.876	3	3 2 2 +	48.48
1.843	1	3 2 3	49.42
1.824	2	3 0 9	49.96
1.812	1	4 1 0	50.32
1.800	5	2 2 8 +	50.66
1.798	6	3 2 4	50.72
1.791	6	2 0 11 +	50.96
1.777	2	1 0 12	51.38
1.758	9	4 1 3 +	51.96
1.746	1	3 2 5	52.36
1.728	5	4 0 7	52.94
1.701	1	1 1 12	53.86
1.687	2	3 2 6	54.32
1.677	1	2 1 11	54.68
1.673	1	4 1 5	54.82
1.666	4	2 0 12	55.08
1.661	5	5 0 0	55.26
1.656	5	5 0 1	55.44
1.652	4	4 0 8	55.58
1.646	2	1 0 13	55.82
1.642	4	5 0 2	55.96
1.625	27	3 2 7	56.58
1.621	18	5 0 3 +	56.74
1.613	25	3 0 11	57.06
1.598	17	3 3 0	57.64
1.589	17	5 0 4	58.00
1.577	1	4 0 9	58.48
1.562	3	3 2 8	59.10
1.559	5	0 0 14	59.22
1.555	4	2 0 13	59.38
1.552	5	5 0 5	59.50
1.532	1	1 0 14	60.36
1.520	1	3 0 12	60.90

Cerium cobalt, Ce₂₄Co₁₁ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.511	1	5 0 6	61.32
1.504	1	4 0 10	61.60
1.477	1	5 1 2	62.86
1.463	1	3 3 6	63.54
1.449	1	2 2 12	64.24
1.441	1	4 2 6	64.64
1.438	2	5 1 4	64.76
1.435	3	3 2 10 +	64.92
1.427	1	3 1 12	65.32
1.379	3	3 3 8 +	67.92
1.375	2	3 2 11	68.14
1.373	2	2 0 15	68.24
1.370	1	5 0 9	68.42
1.365	1	4 3 0	68.70
1.359	3	6 0 3 +	69.04
1.345	3	5 1 7	69.86
1.338	5	4 1 11	70.30
1.330	2	5 2 0	70.80
1.324	4	4 3 4	71.14
1.320	3	2 1 15 +	71.38
1.316	2	3 2 12	71.68
1.308	7	5 2 3 +	72.16
1.274	1	4 2 10	74.40
1.272	1	5 2 5	74.54
1.260	2	3 2 13	75.40
1.251	1	4 3 7	76.04
1.249	1	5 2 6	76.18
1.206	2	3 2 14	79.36
1.192	1	5 1 11	80.50
1.164	1	2 0 18	82.86
1.159	3	5 3 4	83.30
1.156	3	3 2 15	83.54
1.153	2	5 1 12	83.80
1.151	1	6 2 0	83.98
1.150	2	6 2 1	84.12
1.137	1	6 2 3	85.28
1.135	3	6 0 11	85.48
1.131	4	2 1 18	85.84
1.128	3	5 3 6	86.16
1.127	4	6 2 4	86.26
1.116	2	3 3 14 +	87.30
1.113	2	6 2 5	87.56
1.109	5	5 3 7 +	88.02
1.105	5	5 2 11	88.44
1.100	3	7 1 0	88.92
1.095	1	5 0 15	89.46
1.088	2	5 3 8	90.16
1.079	1	2 1 19	91.12
1.042	1	5 3 10	95.32

Calculated Pattern (Integrated)			
d(Å)	I	hkl	2θ(°) λ = 1.540598Å
10.91	1	0 0 2	8.09
8.30	1	1 0 0	10.65
6.61	2	1 0 2	13.39
4.79	1	1 1 0	18.49
3.88	1	2 0 2	22.90
3.61	1	2 0 3	24.67
3.60	1	1 1 4	24.70
3.33	9	1 0 6	26.73
3.30	28	2 0 4	26.96
3.14	32	2 1 0	28.42
3.11	28	2 1 1	28.71
3.02	27	2 1 2	29.59
3.01	7	2 0 5	29.67
2.919	62	1 0 7	30.60
2.898	10	1 1 6	30.83
2.882	19	2 1 3	31.01
2.768	23	3 0 0	32.32
2.746	4	3 0 1	32.58
2.736	16	2 0 6	32.70
2.728	12	0 0 8	32.80
2.721	100	2 1 4	32.90
2.592	1	1 0 8	34.58
2.587	70	3 0 3	34.65
2.548	24	2 1 5	35.19
2.493	63	2 0 7	35.99
2.397	65	2 2 0	37.49
2.376	4	2 1 6	37.83
2.338	4	3 0 5	38.48
2.280	21	2 0 8	39.49
2.253	1	3 1 2	39.98
2.212	2	2 1 7	40.76
2.203	5	3 0 6	40.94
2.196	2	3 1 3	41.08
2.122	10	3 1 4	42.58
2.111	2	1 0 10	42.80
2.094	2	2 0 9	43.16
2.067	3	4 0 1	43.77
2.059	4	2 1 8	43.94
2.039	1	4 0 2	44.39
1.996	1	4 0 3	45.40
1.946	2	3 1 6	46.64
1.943	2	3 0 8	46.71
1.940	2	4 0 4	46.78
1.932	2	2 0 10	47.00
1.930	1	1 0 11	47.05
1.919	4	2 1 9	47.33
1.905	2	3 2 0	47.70
1.877	3	3 2 2	48.47
1.875	1	4 0 5	48.52
1.843	1	3 2 3	49.42

Cerium cobalt, Ce₂₄Co₁₁ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.824	3	3 0 9	49.96
1.812	1	4 1 0	50.32
1.806	1	4 1 1	50.50
1.803	1	4 0 6	50.58
1.801	4	2 2 8	50.65
1.799	5	3 2 4	50.72
1.792	3	2 1 10	50.92
1.790	5	2 0 11	50.97
1.777	2	1 0 12	51.38
1.760	3	3 1 8	51.92
1.758	10	4 1 3	51.97
1.746	1	3 2 5	52.36
1.728	7	4 0 7	52.95
1.701	1	1 1 12	53.87
1.688	3	3 2 6	54.32
1.677	1	2 1 11	54.68
1.674	1	4 1 5	54.81
1.666	5	2 0 12	55.08
1.661	4	5 0 0	55.27
1.656	4	5 0 1	55.44
1.652	3	4 0 8	55.58
1.646	2	1 0 13	55.82
1.642	3	5 0 2	55.96
1.626	37	3 2 7	56.57
1.622	2	4 1 6	56.71
1.619	3	5 0 3	56.82
1.613	32	3 0 11	57.06
1.598	22	3 3 0	57.64
1.589	22	5 0 4	58.01
1.577	1	4 0 9	58.48
1.562	2	3 2 8	59.10
1.559	6	0 0 14	59.22
1.557	2	2 0 13	59.32
1.552	5	5 0 5	59.51
1.532	2	1 0 14	60.36
1.520	1	3 0 12	60.90
1.511	2	5 0 6	61.31
1.504	1	4 0 10	61.61
1.478	1	5 1 2	62.84
1.463	1	3 3 6	63.54
1.449	1	2 2 12	64.23
1.441	1	4 2 6	64.63
1.439	2	5 1 4	64.75
1.435	2	3 2 10	64.92
1.433	1	1 0 15	65.02
1.427	1	3 1 12	65.32
1.402	1	4 2 7	66.67
1.379	4	3 3 8	67.92
1.374	1	3 2 11	68.19
1.373	1	2 0 15	68.24

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.370	1	5 0 9	68.41
1.365	1	4 3 0	68.71
1.360	3	6 0 3	69.03
1.358	2	3 0 14	69.09
1.345	4	5 1 7	69.86
1.338	7	4 1 11	70.30
1.330	2	5 2 0	70.81
1.324	5	4 3 4	71.14
1.322	1	5 0 10	71.30
1.320	2	2 1 15	71.39
1.316	2	3 2 12	71.68
1.309	3	5 1 8	72.12
1.308	7	5 2 3	72.16
1.307	4	2 2 14	72.23
1.303	1	4 3 5	72.49
1.296	1	2 0 16	72.93
1.278	1	4 3 6	74.13
1.274	1	4 2 10	74.40
1.272	1	5 2 5	74.55
1.260	3	3 2 13	75.40
1.250	1	4 3 7	76.05
1.249	1	5 2 6	76.17
1.207	3	3 2 14	79.35
1.192	1	5 1 11	80.50
1.164	1	2 0 18	82.87
1.159	4	5 3 4	83.30
1.156	3	3 2 15	83.54
1.153	2	5 1 12	83.82
1.151	1	6 2 0	83.97
1.150	2	6 2 1	84.12
1.137	2	6 2 3	85.26
1.135	4	6 0 11	85.47
1.131	5	2 1 18	85.84
1.128	2	5 3 6	86.16
1.127	7	6 2 4	86.27
1.116	4	3 3 14	87.30
1.115	1	5 1 13	87.39
1.113	1	6 2 5	87.55
1.109	2	3 2 16	87.98
1.109	2	7 0 7	88.02
1.109	6	5 3 7	88.02
1.107	1	2 0 19	88.17
1.105	7	5 2 11	88.43
1.100	6	7 1 0	88.92
1.094	1	5 0 15	89.47
1.088	3	5 3 8	90.16
1.079	2	2 1 19	91.13
1.042	1	5 3 10	95.31
1.042	1	5 1 15	95.40

Cerium gallium, CeGa₂

Structure

Hexagonal, P6/mmm(191), Z=1, isostructural with AlB₂ [Laves, 1943].

Lattice constants: Haszko [1961]

$$a = 4.32\text{\AA}$$

$$c = 4.34$$

Density

(calculated) 6.62 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Ce⁰ and Ga⁰ [Thomas and Umeda, 1957], corrected for dispersion [Dauben and Templeton, 1955].

Scale factors (integrated intensities)

$$\gamma = 0.988 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 12.8$$

References

- Dauben, C. H. and Templeton, D. H. (1955). Acta Crystallogr. 8, 841.
 Haszko, S. E. (1961). Trans. AIME 221, 201.
 Laves, F. (1943). Naturwissenschaften 31, 145.
 Thomas, L.H. and Umeda, K. (1957). J. Chem. Phys. 26, 293.

Calculated Pattern (Peak heights)				
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å
3.74	13	1 0 0	23.76	
2.834	100	1 0 1	31.54	
2.170	14	0 0 2	41.58	
2.160	37	1 1 0	41.78	
1.877	3	1 0 2	48.46	
1.871	2	2 0 0	48.62	
1.718	17	2 0 1	53.28	
1.531	18	1 1 2	60.42	
1.417	1	2 0 2	65.86	
1.414	1	2 1 0	66.02	
1.349	6	1 0 3	69.62	
1.345	12	2 1 1	69.90	
1.247	3	3 0 0	76.30	
1.185	1	2 1 2	81.12	
1.144	3	2 0 3	84.62	
1.085	1	0 0 4	90.46	
1.081	4	3 0 2	90.86	
1.080	3	2 2 0	91.00	
1.011	3	2 1 3	99.24	
1.009	4	3 1 1	99.52	
.970	2	1 1 4	105.22	
.967	3	2 2 2	105.62	
.914	1	4 0 1	114.80	
.846	1	1 0 5	131.30	
.843	3	3 1 3	132.00	
.842	2	3 2 1	132.38	
.819	2	3 0 4	140.46	
.816	3	4 1 0	141.30	
.787	1	2 0 5	156.10	
.785	2	4 0 3	157.46	

Cerium gallium, CeGa₂ - continued

Calculated Pattern (Integrated)					
d (Å)	I	hkl			2θ (°) λ = 1.540598 Å
3.74	11	1	0	0	23.76
2.834	100	1	0	1	31.55
2.170	13	0	0	2	41.58
2.160	38	1	1	0	41.79
1.877	3	1	0	2	48.46
1.871	2	2	0	0	48.63
1.718	20	2	0	1	53.28
1.531	22	1	1	2	60.42
1.417	1	2	0	2	65.87
1.414	1	2	1	0	66.01
1.349	8	1	0	3	69.62
1.344	15	2	1	1	69.91
1.247	5	3	0	0	76.29
1.185	1	2	1	2	81.11
1.144	4	2	0	3	84.62
1.085	1	0	0	4	90.46
1.081	6	3	0	2	90.86
1.080	3	2	2	0	91.00
1.011	6	2	1	3	99.24
1.009	6	3	1	1	99.51
.970	5	1	1	4	105.21
.967	5	2	2	2	105.63
.936	1	3	1	2	110.75
.914	3	4	0	1	114.81
.861	1	2	1	4	126.98
.846	3	1	0	5	131.29
.843	6	3	1	3	132.01
.842	6	3	2	1	132.37
.819	6	3	0	4	140.46
.816	6	4	1	0	141.31
.798	1	3	2	2	149.65
.787	5	2	0	5	156.10
.785	5	4	0	3	157.46

Cerium magnesium, CeMg₃

Structure

Cubic, face centered, Z=4. [Rossi and Iandelli, 1934]. Their atomic positions indicate Fm3m; other references include Fd3m as a possibility.

Lattice constant: [Vogel and Heumann, 1947]

$$a = 7.438 \text{ \AA}$$

Density

(calculated) 3.439 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Ce⁰, Mg⁰ [International Tables, 1962].

Atom positions

Rossi and Iandelli [1934].

Scale factors (integrated intensities)

$$\gamma = 0.716 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 10.8$$

References

International Tables for X-ray Crystallography III (1962), 202, 211.

Rossi, A. and Iandelli, A. (1934). Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Ser. 6, V.19, 415.

Vogel, R. and Heumann, Th. (1947). Z. Metallk. 38, 1 (1947).

Calculated Pattern (Peak heights)				
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å
4.30	75	1 1 1	20.66	
3.72	38	2 0 0	23.92	
2.63	100	2 2 0	34.06	
2.243	33	3 1 1	40.18	
2.148	9	2 2 2	42.04	
1.860	15	4 0 0	48.94	
1.706	11	3 3 1	53.68	
1.663	10	4 2 0	55.18	
1.518	25	4 2 2	60.98	
1.431	7	5 1 1 +	65.12	
1.315	6	4 4 0	71.72	
1.257	6	5 3 1	75.56	
1.240	3	4 4 2 +	76.84	
1.176	7	6 2 0	81.84	
1.134	2	5 3 3	85.54	
1.121	2	6 2 2	86.78	
1.074	2	4 4 4	91.70	
1.041	3	7 1 1 +	95.40	
1.032	1	6 4 0	96.62	
.994	7	6 4 2	101.60	
.968	3	7 3 1 +	105.40	
.930	1	8 0 0	111.90	
.909	1	7 3 3	115.92	
.902	2	8 2 0 +	117.30	
.877	4	8 2 2 +	122.98	
.859	2	7 5 1 +	127.50	
.853	1	6 6 2	129.06	
.832	3	8 4 0	135.72	
.816	3	7 5 3 +	141.30	
.812	2	8 4 2	143.30	
.793	3	6 6 4	152.58	

Cerium magnesium, CeMg₃ - continued

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
4.29	66	1	1	1	20.67
3.72	35	2	0	0	23.91
2.63	100	2	2	0	34.07
2.243	34	3	1	1	40.18
2.147	10	2	2	2	42.05
1.860	16	4	0	0	48.94
1.706	14	3	3	1	53.67
1.663	12	4	2	0	55.18
1.518	31	4	2	2	60.98
1.431	7	5	1	1	65.11
1.431	2	3	3	3	65.11
1.315	9	4	4	0	71.72
1.257	9	5	3	1	75.57
1.240	1	6	0	0	76.83
1.240	4	4	4	2	76.83
1.176	11	6	2	0	81.84
1.134	3	5	3	3	85.55
1.121	3	6	2	2	86.78
1.074	3	4	4	4	91.70
1.042	2	5	5	1	95.39
1.042	2	7	1	1	95.39
1.031	2	6	4	0	96.63
.994	13	6	4	2	101.61
.968	4	7	3	1	105.40
.968	2	5	5	3	105.40
.930	2	8	0	0	111.89
.909	2	7	3	3	115.92
.902	2	8	2	0	117.30
.902	2	6	4	4	117.30
.877	3	6	6	0	122.99
.877	6	8	2	2	122.99
.859	4	7	5	1	127.50
.859	1	5	5	5	127.50
.853	2	6	6	2	129.07
.832	7	8	4	0	135.73
.816	6	7	5	3	141.30
.816	3	9	1	1	141.30
.812	6	8	4	2	143.31
.793	10	6	6	4	152.58

Cerium nickel, CeNi₂

Structure

Cubic, Fd3m(227), Z=8, C15 type, isostructural with Cu₂Mg [Fülling et al., 1942].

Lattice constant: [Wernick and Geller, 1960]

$$a = 7.202(5) \text{ \AA}$$

Density

(calculated) 9.158 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Ce⁰ and Ni⁰ [Thomas and Umeda, 1957], corrected for dispersion [Dauben and Templeton, 1955].

Scale factors (integrated intensities)

$$\gamma = 0.774 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 10.1$$

Additional patterns

1. Fülling, Moeller, and Vogel [1942].
2. Nowotny [1942].

References

- Dauben, C. H. and Templeton, D. H. (1955). Acta Crystallogr. 8, 841.
- Fülling W., Moeller, K., and Vogel, R. (1942). Z. Metallk. 34, 253.
- Nowotny, H. (1942). Z. Metallk. 34 #11, 247.
- Thomas, L.H. and Umeda, K. (1957). J. Chem. Phys. 26, 293.
- Wernick, J.H. and Geller, S. (1960). Trans. AIME 218, 866.

Calculated Pattern (Integrated)					
d (Å)	I	hkl			2θ (°)
					λ = 1.540598Å
4.16	13	1	1	1	21.35
2.546	67	2	2	0	35.22
2.171	100	3	1	1	41.55
2.079	15	2	2	2	43.49
1.652	3	3	3	1	55.58
1.470	21	4	2	2	63.20
1.386	19	5	1	1	67.53
1.386	6	3	3	3	67.53
1.273	18	4	4	0	74.46
1.217	3	5	3	1	78.51
1.139	8	6	2	0	85.13
1.098	8	5	3	3	89.07
1.086	3	6	2	2	90.38
1.008	1	7	1	1	99.60
1.008	1	5	5	1	99.60
.962	11	6	4	2	106.33
.938	11	7	3	1	110.48
.938	6	5	5	3	110.48
.900	3	8	0	0	117.66
.880	1	7	3	3	122.20
.849	3	6	6	0	130.34
.849	6	8	2	2	130.34
.832	13	7	5	1	135.72
.832	2	5	5	5	135.72
.826	2	6	6	2	137.63
.805	1	8	4	0	146.13
.791	3	7	5	3	154.02
.791	2	9	1	1	154.02

Calculated Pattern (Peak heights)					
$d(\text{\AA})$	I	hkl			$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$					
4.16	16	1	1	1	21.36
2.546	72	2	2	0	35.22
2.171	100	3	1	1	41.56
2.079	14	2	2	2	43.50
1.652	3	3	3	1	55.58
1.470	17	4	2	2	63.20
1.386	20	5	1	1 +	67.52
1.273	13	4	4	0	74.46
1.217	2	5	3	1	78.50
1.139	5	6	2	0	85.14
1.098	5	5	3	3	89.08
1.086	2	6	2	2	90.38
1.009	1	7	1	1 +	99.60
.962	6	6	4	2	106.34
.938	8	7	3	1 +	110.48
.900	2	8	0	0	117.66
.849	4	8	2	2 +	130.34
.832	6	7	5	1 +	135.72
.826	1	6	6	2	137.64
.791	1	7	5	3 +	154.02

Cerium thallium, CeTl

Structure

Cubic, $Pm\bar{3}m$ (221), $Z = 1$, CsCl type [Bruzzzone and Ferro Ruggiero, 1962].

Lattice constant: [ibid.]

$$a = 3.893 \text{ \AA}$$

Density

(calculated) 9.695 g/cm^3

Thermal parameters

Overall isotropic $B = 1.0$

Scattering factors

Ce^0 , Tl^0 [International Tables, 1962].

Scale factors (integrated intensities)

$$\gamma = 1.90 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 32.3$$

References

International Tables for X-ray Crystallography III (1962), 211, 212.

Bruzzzone, G. and Ferro Ruggiero, A. (1962). Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. 33, 465.

Calculated Pattern (Integrated)					
$d(\text{\AA})$	I	hkl			$2\theta(^{\circ})$ $\lambda = 1.540598\text{\AA}$
3.89	4	1	0	0	22.82
2.753	100	1	1	0	32.50
2.248	1	1	1	1	40.08
1.947	17	2	0	0	46.62
1.741	2	2	1	0	52.52
1.589	34	2	1	1	57.98
1.376	10	2	2	0	68.06
1.231	13	3	1	0	77.47
1.124	3	2	2	2	86.54
1.040	15	3	2	1	95.52
.973	2	4	0	0	104.65
.918	3	3	3	0	114.17
.918	6	4	1	1	114.17
.871	7	4	2	0	124.48
.830	8	3	3	2	136.28
.795	11	4	2	2	151.56

Calculated Pattern (Peak heights)

$d(\text{\AA})$	I	hkl			$2\theta(^{\circ})$ $\lambda = 1.540598\text{\AA}$
3.89	4	1	0	0	22.82
2.753	100	1	1	0	32.50
2.248	1	1	1	1	40.08
1.947	15	2	0	0	46.62
1.741	1	2	1	0	52.52
1.589	27	2	1	1	57.98
1.376	7	2	2	0	68.06
1.231	8	3	1	0	77.46
1.124	2	2	2	2	86.54
1.040	8	3	2	1	95.52
.973	1	4	0	0	104.64
.918	4	4	1	1 +	114.18
.870	3	4	2	0	124.48
.830	3	3	3	2	136.28
.795	3	4	2	2	151.56

Cerium thallium, CeTl₃

Structure

Cubic, Pm3m (221), Z = 1, AuCu₃ type [Bruzzone and Ferro Ruggiero, 1962].

Lattice constant: [ibid.]

$$a = 4.767 \text{ \AA}$$

Density

(calculated) 11.55 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Ce⁰, Tl⁰ [International Tables, 1962].

Scale factors (integrated intensities)

$$\gamma = 1.66 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 25.4$$

References

International Tables for X-ray Crystallography III (1962), 211, 212.

Bruzzone, G. and Ferro Ruggiero, A. (1962). Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. 33, 465.

Calculated Pattern (Integrated)				
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å
4.77	2	1 0 0	18.60	
3.37	2	1 1 0	26.42	
2.752	100	1 1 1	32.51	
2.384	49	2 0 0	37.71	
2.132	1	2 1 0	42.36	
1.846	1	2 1 1	46.63	
1.685	32	2 2 0	54.39	
1.437	35	3 1 1	64.81	
1.376	10	2 2 2	68.08	
1.192	4	4 0 0	80.54	
1.094	13	3 3 1	89.55	
1.066	12	4 2 0	92.55	
.973	10	4 2 2	104.68	
.917	10	5 1 1	114.21	
.917	3	3 3 3	114.21	
.843	6	4 4 0	132.15	
.806	29	5 3 1	145.87	
.795	17	4 4 2	151.64	
.795	4	6 0 0	151.64	

Calculated Pattern (Peak heights)				
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å
4.77	2	1 0 0	18.60	
3.37	2	1 1 0	26.42	
2.753	100	1 1 1	32.50	
2.383	47	2 0 0	37.72	
2.132	1	2 1 0	42.36	
1.685	26	2 2 0	54.40	
1.437	27	3 1 1	64.82	
1.376	7	2 2 2	68.08	
1.192	3	4 0 0	80.54	
1.094	8	3 3 1	89.56	
1.066	7	4 2 0	92.54	
.973	5	4 2 2	104.68	
.917	6	5 1 1 +	114.20	
.843	2	4 4 0	132.16	
.806	9	5 3 1	145.88	
.795	6	4 4 2 +	151.64	

Cerium thallium, Ce_3Tl

Structure

Cubic, $\text{Pm}\bar{3}\text{m}$ (221), $Z = 1$, AuCu_3 type [Jeitschko et al., 1964].

Lattice constant: [ibid.]

$$a = 5.011 \text{ \AA}$$

Density

(calculated) 8.245 g/cm^3

Thermal parameters

Overall isotropic $B = 1.0$

Scattering factors

Ce^0 , Tl^0 [International Tables, 1962].

Scale factors (integrated intensities)

$$\gamma = 2.09 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 37.2$$

References

International Tables for X-ray Crystallography III (1962), 211, 212.

Jeitschko, W., Nowotny, H., and Benesovsky, F. (1964). Monatsh. Chem. 95, 1040.

Calculated Pattern (Peak heights)				
$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$	$\lambda = 1.540598\text{\AA}$
5.01	34	1 0 0	17.68	
3.54	27	1 1 0	25.12	
2.893	100	1 1 1	30.88	
2.505	46	2 0 0	35.82	
2.241	13	2 1 0	40.20	
2.046	10	2 1 1	44.24	
1.772	27	2 2 0	51.54	
1.670	5	2 2 1 +	54.92	
1.584	3	3 1 0	58.18	
1.511	27	3 1 1	61.30	
1.446	7	2 2 2	64.36	
1.390	2	3 2 0	67.32	
1.339	3	3 2 1	70.22	
1.253	3	4 0 0	75.88	
1.215	2	4 1 0 +	78.66	
1.181	1	4 1 1 +	81.42	
1.150	8	3 3 1	84.14	
1.120	7	4 2 0	86.86	
1.094	1	4 2 1	89.56	
1.023	5	4 2 2	97.72	
.983	1	4 3 1 +	103.22	
.964	6	5 1 1 +	106.02	
.930	1	4 3 2 +	111.76	
.915	1	5 2 1	114.70	
.886	2	4 4 0	120.82	
.872	1	5 2 2 +	124.02	
.859	1	5 3 0 +	127.36	
.847	7	5 3 1	130.86	
.835	4	4 4 2 +	134.54	
.813	1	5 3 2 +	142.74	
.792	4	6 2 0	152.92	
.783	2	6 2 1 +	159.66	

Cerium thallium, Ce₃Tl - continued

Calculated Pattern (Integrated)				
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å
5.01	30	1 0 0	17.69	
3.54	26	1 1 0	25.11	
2.893	100	1 1 1	30.88	
2.505	50	2 0 0	35.81	
2.241	14	2 1 0	40.21	
2.046	11	2 1 1	44.24	
1.772	33	2 2 0	51.54	
1.670	1	3 0 0	54.92	
1.670	5	2 2 1	54.92	
1.585	5	3 1 0	58.17	
1.511	37	3 1 1	61.31	
1.447	10	2 2 2	64.35	
1.390	3	3 2 0	67.32	
1.379	5	3 2 1	70.22	
1.253	4	4 0 0	75.89	
1.215	2	4 1 0	78.66	
1.215	2	3 2 2	78.66	
1.191	1	3 3 0	81.41	
1.191	2	4 1 1	81.41	
1.150	13	3 3 1	84.14	
1.120	12	4 2 0	86.86	
1.093	2	4 2 1	89.57	
1.068	1	3 3 2	92.28	
1.023	9	4 2 2	97.72	
1.002	1	4 3 0	100.46	
.993	2	4 3 1	103.23	
.983	1	5 1 0	103.23	
.964	9	5 1 1	106.02	
.964	3	3 3 3	106.02	
.931	1	5 2 0	111.75	
.931	2	4 3 2	111.75	
.915	2	5 2 1	114.70	
.886	4	4 4 0	120.82	
.872	1	4 4 1	124.03	
.872	1	5 2 2	124.03	
.859	1	5 3 0	127.36	
.859	1	4 3 3	127.36	
.847	18	5 3 1	130.85	
.835	2	6 0 0	134.54	
.835	10	4 4 2	134.54	
.824	1	6 1 0	138.47	
.813	2	5 3 2	142.74	
.813	1	6 1 1	142.74	
.792	15	6 2 0	152.93	
.792	2	5 4 0	152.67	
.783	4	6 2 1	159.67	
.783	2	4 4 3	159.67	

Cobalt dysprosium, Co₂Dy

Structure

Cubic, Fd3m(227), Z=8, C15 type, isostructural with Cu₂Mg [Wernick and Geller, 1960].

Lattice constant: [ibid.]

$$a = 7.187(5)\text{\AA}$$

Density

(calculated) 10.033 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Co⁰ and Dy⁰ [Thomas and Umeda, 1957], corrected for dispersion [Dauben and Templeton, 1955].

Scale factors (integrated intensities)

$$\gamma = 0.510 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 9.35$$

References

Dauben, C. H. and Templeton, D. H. (1955). Acta Crystallogr. 8, 841.

Thomas, L.H. and Umeda, K. (1957). J. Chem. Phys. 26, 293.

Wernick, J.H. and Geller, S. (1960). Trans. AIME 218, 866.

Calculated Pattern (Integrated)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
4.15	13	1 1 1	21.40
2.541	66	2 2 0	35.29
2.167	100	3 1 1	41.65
2.075	16	2 2 2	43.59
1.649	3	3 3 1	55.70
1.467	19	4 2 2	63.35
1.383	19	5 1 1	67.69
1.383	6	3 3 3	67.69
1.270	17	4 4 0	74.64
1.215	2	5 3 1	78.70
1.136	7	6 2 0	85.35
1.096	8	5 3 3	89.31
1.083	4	6 2 2	90.62
.960	9	6 4 2	106.65
.936	11	7 3 1	110.83
.936	5	5 5 3	110.83
.898	3	8 0 0	118.06
.847	2	6 6 0	130.86
.847	5	8 2 2	130.86
.830	12	7 5 1	136.31
.830	2	5 5 5	136.31
.824	3	6 6 2	138.25
.789	2	7 5 3	155.08
.789	1	9 1 1	155.08

Calculated Pattern (Peak heights)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
4.15	16	1 1 1	21.40
2.541	70	2 2 0	35.30
2.167	100	3 1 1	41.64
2.075	15	2 2 2	43.58
1.649	2	3 3 1	55.70
1.467	16	4 2 2	63.34
1.383	19	5 1 1 +	67.68
1.271	13	4 4 0	74.64
1.215	1	5 3 1	78.70
1.136	5	6 2 0	85.36
1.096	5	5 3 3	89.30
1.084	2	6 2 2	90.62
.960	5	6 4 2	106.66
.936	8	7 3 1 +	110.82
.898	2	8 0 0	118.06
.847	3	8 2 2 +	130.86
.830	5	7 5 1 +	136.32
.824	1	6 6 2	138.26

Cobalt erbium, Co₂Er

Structure

Cubic, Fd3m(227), Z=8, C15 type, isostructural with Cu₂Mg [Wernick and Geller, 1960].

Lattice constant: [ibid.]

$$a = 7.144(5)\text{\AA}$$

Density

(calculated) 10.388 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Co⁰ and Er⁰ [Thomas and Umeda, 1957], corrected for dispersion [Dauben and Templeton, 1955].

Scale factors (integrated intensities)

$$\gamma = 0.762 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 9.76$$

References

- Dauben, C. H. and Templeton, D. H. (1955). Acta Crystallogr. 8, 841.
 Thomas, L.H. and Umeda, K. (1957). J. Chem. Phys. 26, 293.
 Wernick, J.H. and Geller, S. (1960). Trans. AIME 218, 866.

Calculated Pattern (Integrated)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
4.12	15	1 1 1	21.53
2.526	68	2 2 0	35.51
2.154	100	3 1 1	41.91
2.062	15	2 2 2	43.87
1.639	3	3 3 1	56.07
1.458	20	4 2 2	63.77
1.375	19	5 1 1	68.15
1.375	6	3 3 3	68.15
1.263	17	4 4 0	75.17
1.208	2	5 3 1	79.27
1.130	7	6 2 0	85.99
1.089	8	5 3 3	89.99
1.077	4	6 2 2	91.32
.955	9	6 4 2	107.59
.930	11	7 3 1	111.83
.930	5	5 5 3	111.83
.893	4	8 0 0	119.22
.842	2	6 6 0	132.39
.842	5	8 2 2	132.39
.825	13	7 5 1	138.07
.825	2	5 5 5	138.07
.819	3	6 6 2	140.10
.784	2	7 5 3	158.43
.784	1	9 1 1	158.43

Calculated Pattern (Peak heights)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
4.12	19	1 1 1	21.54
2.525	74	2 2 0	35.52
2.154	100	3 1 1	41.90
2.063	15	2 2 2	43.86
1.639	3	3 3 1	56.06
1.458	16	4 2 2	63.78
1.375	19	5 1 1 +	68.14
1.263	13	4 4 0	75.18
1.207	1	5 3 1	79.28
1.129	5	6 2 0	86.00
1.089	5	5 3 3	90.00
1.077	2	6 2 2	91.32
.955	5	6 4 2	107.58
.930	8	7 3 1 +	111.84
.893	2	8 0 0	119.22
.842	3	8 2 2 +	132.38
.825	6	7 5 1 +	138.06
.819	1	6 6 2	140.10

Cobalt erbium, Co₇Er₂

Structure

Hexagonal, $R\bar{3}m(166)$, $Z = 6$. The structure was determined by Ostertag [1967].

Lattice constants: [ibid.]

$$a = 4.973\text{\AA}$$

$$c = 36.11$$

Density

(measured) 9.620 g/cm³ [ibid.]
(calculated) 9.624 g/cm³

Thermal parameters

Isotropic [Ostertag, op. cit.]

Scattering factors

Co⁰, Er⁰ [International Tables, 1962].

Scale factors (integrated intensities)

$\gamma = 0.467 \times 10^{-3}$
 I/I_c (calculated) 6.70

References

International Tables for X-ray Crystallography III (1962), 204, 212.
Ostertag, W. (1967). J. Less-Common Metals, 13, 385.

Calculated Pattern (Peak heights)

d (Å)	I	hkl	2θ (°)
$\lambda = 1.540598\text{\AA}$			
12.04	5	0 0 3	7.34
6.02	1	0 0 6	14.72
4.28	12	1 0 1	20.76
4.19	2	0 1 2	21.20
4.01	5	0 0 9	22.14
3.89	4	1 0 4	22.86
3.116	2	0 1 8	28.64
3.009	5	0 0 12	29.68
2.767	49	1 0 10	32.34
2.611	39	0 1 11	34.32
2.487	58	1 1 0	36.10
2.407	2	0 0 15	37.32
2.334	2	1 0 13	38.54
2.213	9	0 1 14	40.74
2.150	40	0 2 1	42.00
2.114	100	1 1 9	42.76
2.095	10	0 2 4	43.14
2.064	4	2 0 5	43.84
2.006	14	0 0 18	45.16
1.999	10	1 0 16	45.26

d (Å)	I	hkl	2θ (°)
$\lambda = 1.540598\text{\AA}$			
1.987	8	0 2 7	45.62
1.944	12	2 0 8	46.70
1.917	8	1 1 12	47.40
1.905	7	0 1 17	47.70
1.801	3	2 0 11	50.66
1.730	3	1 1 15	52.90
1.720	4	0 0 21	53.22
1.665	6	0 1 20	55.12
1.653	1	2 0 14	55.54
1.626	2	2 1 1	56.56
1.561	2	1 1 18	59.12
1.558	2	0 2 16	59.28
1.534	3	1 0 22	60.30
1.512	1	2 0 17	61.24
1.505	1	0 0 24	61.58
1.484	13	2 1 10	62.54
1.458	11	1 2 11	63.76
1.436	9	3 0 0	64.90
1.425	6	0 2 19	65.44
1.414	12	1 1 21	66.00
1.384	8	2 0 20	67.66
1.377	4	1 2 14	68.06
1.369	2	1 0 25	68.46
1.352	20	3 0 9+	69.48
1.296	2	3 0 12+	72.96
1.292	4	1 2 17	73.20
1.287	3	1 1 24	73.50
1.269	1	2 0 23	74.78
1.243	19	2 2 0	76.58
1.233	1	3 0 15+	77.32
1.209	4	1 2 20	79.16
1.204	2	0 0 30	79.58
1.196	1	0 1 29	80.18
1.194	1	1 3 1	80.38
1.188	1	2 2 9	80.88
1.178	3	1 1 27	81.68
1.167	2	2 0 26+	82.60
1.156	2	2 1 22	83.60
1.149	1	2 2 12	84.20
1.134	5	1 3 10	85.58
1.124	3	1 0 31	86.48
1.122	5	3 1 11	86.68
1.106	2	0 2 28	88.26
1.102	6	0 3 21+	88.70
1.083	6	1 1 30+	90.62
1.080	4	2 1 25	90.92
1.076	4	4 0 1	91.42
1.057	8	2 2 18	93.60
1.047	1	0 4 8	94.70
1.041	2	3 1 17	95.44

Cobalt erbium, Co₇Er₂ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.039	2	0 3 24+	95.74
1.025	3	0 2 31	97.50
1.007	4	2 2 21	99.74
1.003	2	0 0 36	100.34
.9995	1	2 0 32	100.82
.9962	3	3 1 20	101.30
.9890	1	1 2 29	102.32
.9786	1	3 0 27+	103.84
.9658	1	1 3 22	105.80
.9584	1	2 2 24	106.98
.9530	5	3 2 10+	107.86
.9473	3	2 1 31	108.82
.9461	4	2 3 11	109.02
.9398	6	4 1 0	110.10
.9368	3	4 0 19	110.52
.9304	3	2 0 35+	111.78
.9247	3	0 4 20	112.82
.9224	6	3 0 30+	113.26
.9205	3	1 3 25	113.66
.9150	14	1 4 9+	114.66
.9106	1	2 2 27	115.54
.8971	2	1 4 12+	118.34
.8959	1	2 3 17	118.60
.8895	1	2 1 34+	120.00
.8835	1	1 0 40	121.34
.8755	1	1 4 15+	123.26
.8677	1	1 1 39	125.20
.8667	3	2 3 20	125.42
.8648	5	2 2 30	125.94
.8629	3	0 1 41	126.48
.8509	1	0 4 26+	129.70
.8465	1	3 2 22	131.00
.8378	2	0 5 10	133.68
.8370	1	2 1 37	133.92
.8339	3	1 3 31	134.94
.8331	2	5 0 11	135.20
.8288	3	3 3 0	136.68
.8265	2	4 0 28	137.44
.8247	8	4 1 21+	138.16
.8203	1	3 1 32	139.80
.8170	1	5 0 14	141.08
.8155	2	3 2 25+	141.70
.8137	6	2 4 1	142.42
.8126	3	1 1 42	142.90
.8117	10	3 3 9	143.24
.8106	2	2 4 4	143.70
.8088	1	4 2 5	144.52
.8062	1	0 1 44	145.68
.8040	1	2 4 7	146.72
.8010	2	4 2 8	148.18

Calculated Pattern (Integrated)				
$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$	$\lambda = 1.540598\text{\AA}$
12.04	4	0 0 3	7.34	
6.02	1	0 0 6	14.71	
4.28	10	1 0 1	20.75	
4.19	1	0 1 2	21.19	
4.01	4	0 0 9	22.14	
3.89	3	1 0 4	22.86	
3.116	2	0 1 8	28.63	
3.009	5	0 0 12	29.66	
2.767	46	1 0 10	32.33	
2.611	37	0 1 11	34.32	
2.487	55	1 1 0	36.00	
2.407	2	0 0 15	37.32	
2.334	1	1 0 13	38.54	
2.213	9	0 1 14	40.74	
2.150	38	0 2 1	42.00	
2.138	9	2 0 2	42.23	
2.114	100	1 1 9	42.75	
2.095	6	0 2 4	43.15	
2.064	4	2 0 5	43.84	
2.006	14	0 0 18	45.16	
1.999	1	1 0 16	45.33	
1.987	7	0 2 7	45.61	
1.944	12	2 0 8	46.70	
1.917	7	1 1 12	47.39	
1.905	7	0 1 17	47.70	
1.801	3	2 0 11	50.66	
1.730	3	1 1 15	52.89	
1.720	4	0 0 21	53.23	
1.665	7	0 1 20	55.11	
1.653	1	2 0 14	55.55	
1.626	2	2 1 1	56.55	
1.561	2	1 1 18	59.12	
1.558	1	0 2 16	59.26	
1.534	3	1 0 22	60.30	
1.512	1	2 0 17	61.25	
1.505	1	0 0 24	61.59	
1.484	14	2 1 10	62.54	
1.458	12	1 2 11	63.77	
1.436	10	3 0 0	64.90	
1.425	6	0 2 19	65.45	
1.414	13	1 1 21	66.00	
1.384	9	2 0 20	67.66	
1.377	4	1 2 14	68.05	
1.369	2	1 0 25	68.46	
1.352	11	3 0 9	69.40	
1.352	11	0 3 9	69.40	
1.296	1	3 0 12	72.96	
1.296	1	0 3 12	72.96	
1.292	4	1 2 17	73.20	
1.287	3	1 1 24	73.51	

Cobalt erbium, Co₇Er₂ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.269	1	2 0 23	74.77
1.243	21	2 2 0	76.57
1.209	5	1 2 20	79.16
1.204	1	0 0 30	79.58
1.196	1	0 1 29	80.18
1.194	1	1 3 1	80.37
1.188	1	2 2 9	80.88
1.178	3	1 1 27	81.69
1.167	2	2 0 26	82.60
1.156	2	2 1 22	83.59
1.149	2	2 2 12	84.19
1.134	6	1 3 10	85.57
1.124	2	1 0 31	86.48
1.122	5	3 1 11	86.67
1.106	2	0 2 28	88.25
1.105	1	2 2 15	88.43
1.102	3	3 0 21	88.69
1.102	3	0 3 21	88.69
1.084	2	3 1 14	90.58
1.083	5	1 1 30	90.63
1.080	2	2 1 25	90.96
1.076	4	4 0 1	91.41
1.075	1	0 4 2	91.57
1.057	9	2 2 18	93.59
1.054	1	4 0 7	93.92
1.047	1	0 4 8	94.70
1.041	2	3 1 17	95.44
1.039	1	3 0 24	95.74
1.039	1	0 3 24	95.74
1.025	4	0 2 31	97.50
1.007	5	2 2 21	99.74
1.003	2	0 0 36	100.34
.9995	1	2 0 32	100.83
.9962	3	3 1 20	101.29
.9890	1	1 2 29	102.31
.9786	1	0 3 27	103.85
.9786	1	3 0 27	103.85
.9658	2	1 3 22	105.80
.9584	1	2 2 24	106.98
.9530	5	3 2 10	107.86
.9525	2	0 2 34	107.94
.9473	4	2 1 31	108.81
.9461	4	2 3 11	109.01
.9398	7	4 1 0	110.10
.9368	2	4 0 19	110.62
.9304	3	2 0 35	111.77
.9302	1	1 1 36	111.80
.9274	1	1 2 32	112.32
.9247	4	0 4 20	112.81
.9227	2	2 3 14	113.20

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
.9224	2	3 0 30	113.26
.9224	2	0 3 30	113.26
.9205	2	1 3 25	113.61
.9150	10	4 1 9	114.67
.9150	10	1 4 9	114.67
.9106	1	2 2 27	115.55
.8971	1	4 1 12	118.34
.8971	1	1 4 12	118.34
.8959	2	2 3 17	118.60
.8895	1	2 1 34	120.00
.8835	2	1 0 40	121.34
.8677	1	1 1 39	125.19
.8667	4	2 3 20	125.43
.8648	6	2 2 30	125.94
.8629	2	0 1 41	126.43
.8620	1	3 1 29	126.67
.8509	1	0 4 26	129.71
.8465	2	3 2 22	131.01
.8378	3	0 5 10	133.67
.8370	1	2 1 37	133.93
.8339	5	1 3 31	134.94
.8331	3	5 0 11	135.21
.8325	1	0 2 40	135.41
.8288	5	3 3 0	136.68
.8265	2	4 0 28	137.50
.8247	8	4 1 21	138.16
.8247	8	1 4 21	138.16
.8222	1	0 3 36	139.05
.8222	1	3 0 36	139.05
.8203	1	3 1 32	139.79
.8170	1	5 0 14	141.07
.8155	2	3 2 25	141.67
.8152	2	2 0 41	141.80
.8137	11	2 4 1	142.41
.8131	2	4 2 2	142.67
.8126	2	1 1 42	142.88
.8117	15	3 3 9	143.25
.8106	2	2 4 4	143.71
.8088	1	4 2 5	144.51
.8062	2	0 1 44	145.69
.8040	3	2 4 7	146.73
.8010	4	4 2 8	148.18
.7991	2	3 3 12	149.15
.7982	2	5 0 17	149.60
.7971	3	4 1 24	150.21
.7971	3	1 4 24	150.21
.7937	2	1 3 34	152.11
.7907	8	4 0 31	153.94
.7900	3	4 2 11	154.37
.7895	9	2 1 40	154.69

Cobalt gadolinium, CoGd₃

Structure

Orthorhombic, Pnma(62), Z=4, isostructural with CFe₃, type DO₁₁. The structure was determined by Strydom and Alberts [1970].

Lattice constants: [ibid.]

a = 7.05Å
b = 9.54
c = 6.32

Density

(calculated) 8.29 g/cm³

Thermal parameters

Isotropic [Strydom and Alberts, op. cit.].

Scattering factors

Co⁰, Gd⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$\gamma = 0.168 \times 10^{-3}$
I/I_c (calculated) 4.41

References

Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Strydom, O.A.W. and Alberts, L. (1970). J. Less-Common Metals, 22, 511.

Calculated Pattern (Peak heights)					
d(Å)	I	h	k	l	2θ(°)
λ = 1.540598Å					
4.22	1	1	1	1	21.04
3.525	1	2	0	0	25.26
3.350	18	1	2	1	26.60
3.307	13	2	1	0	26.96
3.160	9	0	0	2	28.22
3.079	15	2	0	1	28.98
2.930	35	2	1	1	30.50
2.884	38	1	0	2	31.00
2.841	100	0	3	1+	31.52
2.760	51	1	1	2	32.42
2.635	40	1	3	1+	34.00
2.587	44	2	2	1	34.66
2.468	14	1	2	2	36.38
2.385	6	0	4	0	37.70
2.361	15	2	3	0	38.08
2.284	8	2	1	2	39.42
2.203	15	3	0	1	40.94
2.146	8	3	1	1	42.08
2.136	6	1	3	2	42.16
2.127	2	1	4	1	42.44

d(Å)	I	h	k	l	2θ(°)
λ = 1.540598Å					
2.110	3	2	2	2	42.82
1.975	7	2	4	0+	45.92
1.891	2	2	3	2	48.08
1.885	3	2	4	1	48.22
1.859	16	1	2	3	48.96
1.827	3	0	5	1	49.88
1.811	3	3	3	1+	50.40
1.777	1	2	1	3	51.40
1.768	1	1	5	1	51.66
1.754	3	3	2	2	52.12
1.704	14	1	3	3	53.74
1.698	16	4	0	1	53.96
1.678	5	2	5	0	54.66
1.671	4	4	1	1	54.80
1.653	1	4	2	0	55.54
1.622	24	3	3	2+	56.72
1.599	2	4	2	1	57.60
1.591	15	1	5	2+	57.92
1.580	3	0	0	4	58.36
1.569	10	3	0	3	58.82
1.542	17	4	3	0+	59.96
1.490	2	3	2	3	62.26
1.482	3	2	5	2	62.64
1.479	2	3	4	2	62.80
1.467	2	1	2	4	63.36
1.465	2	4	2	2	63.44
1.442	2	3	5	1+	64.58
1.426	1	2	1	4	65.42
1.420	1	0	6	2	65.60
1.413	1	2	6	1+	66.08
1.392	3	1	6	2	67.18
1.387	6	1	5	3+	67.50
1.380	3	2	2	4	67.84
1.362	1	5	1	1	68.88
1.341	1	3	5	2	70.10
1.332	1	0	7	1	70.64
1.322	1	5	2	1	71.26
1.317	1	0	4	4+	71.56
1.301	1	4	2	3	72.64
1.295	1	4	5	0+	73.02
1.289	2	3	6	1	73.38
1.268	1	4	5	1	74.80
1.264	1	3	2	4	75.04
1.243	4	5	2	2+	76.58
1.234	1	2	4	4+	77.28
1.232	1	1	7	2	77.38
1.212	4	3	3	4	78.92
1.199	1	1	5	4	79.94
1.198	1	4	5	2	80.02
1.194	1	5	3	2+	80.38

Cobalt gadolinium, CoGd₃ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.190	1	2 0 5	80.64
1.181	1	2 1 5	81.46
1.175	3	0 3 5+	81.96
1.172	3	5 0 3	82.20
1.166	1	6 1 0	82.68
1.163	2	5 1 3	82.96
1.161	4	4 6 1	83.18
1.159	2	1 3 5	83.34
1.156	2	1 8 1	83.58
1.154	1	2 2 5	83.70
1.150	1	2 5 4	84.08
1.141	1	6 2 0+	84.94
1.138	1	5 2 3	85.20
1.133	1	5 4 2	85.66
1.130	2	2 8 0	85.98
1.123	1	6 2 1	86.64
1.121	1	0 6 4	86.84
1.117	3	3 6 3+	87.24
1.112	3	2 8 1	87.68
1.107	2	1 6 4	88.20
1.103	2	4 3 4+	88.54
1.102	2	1 8 2+	88.70
1.086	2	6 3 1	90.38
1.084	1	3 2 5	90.56
1.081	1	3 5 4	90.92
1.078	1	4 7 0	91.20
1.067	1	5 5 2	92.40
1.053	1	0 0 6	94.00
1.045	1	0 9 1+	94.92
1.042	1	1 5 5	95.30
1.027	3	1 8 3+	97.20
1.010	1	2 5 5	99.46
1.001	1	6 5 0	100.70
.9985	2	5 5 3+	100.96
.9882	1	6 5 1	102.44
.9874	1	2 2 6	102.52
.9596	2	7 0 2	106.78
.9540	1	0 10 0	107.70
.9493	1	3 8 3+	108.48
.9433	1	1 8 4+	109.50
.9240	1	3 9 2	112.94
.9213	1	5 5 4	113.46
.9189	1	2 8 4	113.92
.8921	1	2 5 6	119.42
.8906	1	4 7 4+	119.88
.8781	1	0 6 6	122.64
.8749	2	5 8 2+	123.38
.8737	2	7 3 3	123.70
.8697	1	4 3 6	124.68
.8628	1	4 6 5	126.46
.8586	1	8 2 1+	127.60

Calculated Pattern (Integrated)					
d(A)	I	hkl			2θ(°) λ = 1.540598Å
4.22	1	1	1	1	21.03
3.525	1	2	0	0	25.24
3.350	23	1	2	1	26.50
3.307	16	2	1	0	26.94
3.160	12	0	0	2	28.22
3.079	20	2	0	1	28.98
2.930	46	2	1	1	30.49
2.884	48	1	0	2	30.99
2.841	100	0	3	1	31.47
2.835	57	2	2	0	31.53
2.760	69	1	1	2	32.41
2.635	42	1	3	1	34.00
2.634	14	0	2	2	34.00
2.587	61	2	2	1	34.65
2.468	19	1	2	2	36.38
2.385	9	0	4	0	37.69
2.361	21	2	3	0	38.08
2.353	4	2	0	2	38.22
2.284	11	2	1	2	39.41
2.212	2	2	3	1	40.76
2.203	20	3	0	1	40.94
2.146	11	3	1	1	42.07
2.136	3	1	3	2	42.27
2.127	2	1	4	1	42.46
2.110	4	2	2	2	42.82
2.057	1	0	1	3	43.98
1.975	6	2	4	0	45.90
1.975	5	1	1	3	45.92
1.891	2	2	3	2	48.06
1.885	3	2	4	1	48.23
1.859	24	1	2	3	48.96
1.827	4	0	5	1	49.80
1.811	3	3	3	1	50.35
1.808	2	2	0	3	50.42
1.777	1	2	1	3	51.39
1.768	1	1	5	1	51.65
1.754	4	3	2	2	52.11
1.704	20	1	3	3	53.75
1.698	19	4	0	1	53.97
1.691	1	2	2	3	54.20
1.678	8	2	5	0	54.65
1.671	3	4	1	1	54.88
1.653	1	4	2	0	55.54
1.622	24	3	3	2	56.71
1.622	13	2	5	1	56.72
1.618	2	3	4	1	56.85
1.599	1	4	2	1	57.58
1.591	15	1	5	2	57.91
1.590	10	0	6	0	57.95
1.580	3	0	0	4	58.36

Cobalt gadolinium, CoGd₃ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.572	1	2 3 3	58.68
1.569	14	3 0 3	58.82
1.548	1	3 1 3	59.69
1.542	8	1 0 4	59.95
1.542	14	4 3 0	59.96
1.541	5	1 4 3	59.99
1.490	3	3 2 3	62.25
1.482	5	2 5 2	62.63
1.479	1	3 4 2	62.77
1.467	3	1 2 4	63.35
1.465	1	4 2 2	63.45
1.442	2	3 5 1	64.57
1.441	1	2 4 3	64.63
1.426	2	2 1 4	65.41
1.420	1	0 6 2	65.69
1.413	1	2 6 1	66.09
1.392	4	1 6 2	67.18
1.387	2	1 3 4	67.46
1.387	6	1 5 3	67.50
1.385	4	4 3 2	67.56
1.383	1	4 4 1	67.69
1.380	3	2 2 4	67.85
1.362	1	5 1 1	68.88
1.341	2	3 5 2	70.11
1.332	2	0 7 1	70.65
1.322	2	5 2 1	71.26
1.309	1	1 7 1	72.09
1.301	1	4 2 3	72.64
1.295	1	4 5 0	73.02
1.289	3	3 6 1	73.38
1.268	2	4 5 1	74.79
1.264	1	3 2 4	75.07
1.244	1	1 0 5	76.51
1.243	5	5 2 2	76.58
1.234	1	2 4 4	77.26
1.232	1	1 7 2	77.39
1.212	7	3 3 4	78.91
1.212	1	3 5 3	78.95
1.199	1	1 5 4	79.93
1.198	1	4 5 2	80.03
1.194	1	5 3 2	80.39
1.190	1	2 0 5	80.69
1.181	2	2 1 5	81.45
1.175	1	6 0 0	81.93
1.175	4	0 3 5	81.96
1.172	1	5 0 3	82.20
1.166	1	6 1 0	82.68
1.163	3	5 1 3	82.95
1.161	5	4 6 1	83.17
1.159	1	1 3 5	83.34

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.156	1	1 8 1	83.58
1.154	1	2 2 5	83.71
1.150	1	2 5 4	84.08
1.141	1	6 2 0	84.93
1.138	1	5 2 3	85.21
1.133	1	5 4 2	85.66
1.130	2	2 8 0	85.99
1.123	1	6 2 1	86.64
1.121	1	0 6 4	86.83
1.117	4	3 6 3	87.23
1.116	1	5 5 1	87.28
1.116	1	0 8 2	87.33
1.112	4	2 8 1	87.69
1.107	2	1 6 4	88.20
1.105	1	3 7 2	88.43
1.103	2	4 3 4	88.55
1.102	2	1 8 2	88.70
1.101	1	6 0 2	88.76
1.086	2	6 3 1	90.38
1.084	1	3 2 5	90.56
1.081	2	3 5 4	90.93
1.078	1	4 7 0	91.20
1.067	1	5 5 2	92.39
1.053	1	0 0 6	93.99
1.045	1	0 9 1	94.93
1.042	1	1 5 5	95.31
1.027	1	5 2 4	97.15
1.027	2	4 0 5	97.17
1.027	3	1 8 3	97.23
1.010	2	2 5 5	99.45
1.001	1	6 5 0	100.69
.9985	3	5 5 3	100.97
.9882	1	6 5 1	102.43
.9874	1	2 2 6	102.55
.9620	1	2 3 6	106.41
.9596	2	7 0 2	106.79
.9540	1	0 10 0	107.69
.9493	1	3 8 3	108.47
.9492	1	7 3 1	108.48
.9433	1	1 8 4	109.50
.9268	1	0 7 5	112.44
.9240	1	3 9 2	112.95
.9213	1	5 5 4	113.47
.9189	1	2 8 4	113.92
.9084	1	4 9 0	115.99
.9025	1	5 3 5	117.20
.8921	2	2 5 6	119.41
.8802	1	1 2 7	122.13
.8781	1	0 6 6	122.62
.8749	3	5 8 2	123.39

Cobalt gadolinium, Co₂Gd

Structure

Cubic, Fd3m(227), Z=8, C15 type, isostructural with Cu₂Mg [Wernick and Geller, 1960].

Lattice constant: [ibid.]

$$a = 7.255(5)\text{\AA}$$

Density

(calculated) 9.571 g/cm³

Thermal parameters

Overall isotropic B = 1.0

Scattering factors

Co⁰ and Gd⁰ [Thomas and Umeda, 1957], corrected for dispersion [Dauben and Templeton, 1955].

Scale factors (integrated intensities)

$$\gamma = 0.372 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 8.83$$

References

Dauben, C. H. and Templeton, D. H. (1955). Acta Crystallogr. 8, 841.

Thomas, L.H. and Umeda, K. (1957). J. Chem. Phys. 26, 293.

Wernick, J.H. and Geller, S. (1960). Trans. AIME 218, 866.

Calculated Pattern (Integrated)

d(Å)	I	hkl	2θ(°)
λ = 1.540598Å			
4.19	10	1 1 1	21.19
2.565	63	2 2 0	34.95
2.187	100	3 1 1	41.24
2.094	17	2 2 2	43.16
1.664	2	3 3 1	55.14
1.481	18	4 2 2	62.68
1.396	19	5 1 1	66.97
1.396	6	3 3 3	66.97
1.283	18	4 4 0	73.83
1.226	1	5 3 1	77.83
1.147	7	6 2 0	84.37
1.106	8	5 3 3	88.25
1.094	4	6 2 2	89.54
.969	9	6 4 2	105.22
.945	10	7 3 1	109.28
.945	5	5 5 3	109.28
.907	3	8 0 0	116.29
.855	2	6 6 0	128.56
.855	4	8 2 2	128.56
.838	11	7 5 1	133.71
.838	2	5 5 5	133.71
.832	3	6 6 2	135.52
.796	1	7 5 3	150.61

Calculated Pattern (Peak heights)

d(Å)	I	hkl	2θ(°)
λ = 1.540598Å			
4.19	12	1 1 1	21.20
2.564	64	2 2 0	34.96
2.187	100	3 1 1	41.24
2.094	17	2 2 2	43.16
1.664	2	3 3 1	55.14
1.481	15	4 2 2	62.68
1.396	19	5 1 1 +	66.96
1.283	13	4 4 0	73.82
1.226	1	5 3 1	77.82
1.147	4	6 2 0	84.36
1.106	5	5 3 3	88.26
1.094	3	6 2 2	89.54
.970	5	6 4 2	105.22
.945	8	7 3 1 +	109.28
.907	2	8 0 0	116.30
.855	3	8 2 2 +	128.56
.838	5	7 5 1 +	133.70
.832	1	6 6 2	135.52

Cobalt gadolinium, Co₇Gd₂

Structure

Hexagonal, $R\bar{3}m(166)$, $Z = 6$. The structure was determined by Bertaut et al. [1965].

Lattice constants: [Ostertag, 1967]

$$a = 5.023\text{\AA}$$

$$c = 36.29$$

Density

(calculated) 9.135 g/cm³

Thermal parameters

Isotropic [Bertaut et al., op. cit.]

Scattering factors

Co⁰, Gd⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.302 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 6.84$$

References

Bertaut, E. F., Lemaire, F. G. R., and Schweizer, J. (1965). C. R. Acad. Sci. 260, 3595.
Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Ostertag, W. (1967). J. Less-Common Metals, 13, 385.

Calculated Pattern (Peak heights)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
12.10	7	0 0 3	7.30
4.319	7	1 0 1	20.56
4.032	6	0 0 9	22.04
3.922	3	1 0 4	22.66
3.140	4	0 1 8	28.42
3.024	1	0 0 12	29.52
2.787	43	1 0 10	32.10
2.629	25	0 1 11	34.08
2.512	50	1 1 0	35.72
2.459	1	1 1 3	36.52
2.419	1	0 0 15	37.14
2.349	4	1 0 13	38.28
2.227	10	0 1 14	40.42
2.171	35	0 2 1	41.56
2.132	100	1 1 9	42.36
2.115	11	0 2 4	42.70
2.083	4	2 0 5	43.40
2.016	16	0 0 12	44.92
2.006	8	0 2 7	45.16
1.961	9	2 0 8	46.26

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.932	2	1 1 12	47.00
1.916	5	0 1 17	47.40
1.816	1	2 0 11	50.20
1.742	3	1 1 15	52.42
1.728	2	0 0 21	52.94
1.675	4	0 1 20	54.78
1.666	1	2 0 14	55.06
1.642	1	2 1 1	55.94
1.572	4	1 1 18	58.68
1.546	1	1 2 8	59.78
1.524	1	2 0 17	60.74
1.512	1	0 0 24	61.26
1.498	10	2 1 10	61.90
1.483	2	0 1 23	62.58
1.472	7	1 2 11	63.14
1.450	7	3 0 0	64.18
1.435	7	0 2 19	64.92
1.424	5	1 1 21	65.52
1.417	2	2 1 13	65.86
1.393	7	2 0 20	67.12
1.388	5	1 2 14	67.32
1.377	1	1 0 25	68.04
1.364	18	0 3 9+	68.74
1.303	2	1 2 17	72.50
1.295	4	1 1 24	72.98
1.277	2	2 0 23	74.12
1.256	15	2 2 0	75.62
1.244	1	3 0 15+	76.54
1.218	3	1 2 20	78.44
1.210	1	0 0 30	79.10
1.203	1	0 1 29	79.66
1.199	1	2 2 9	79.94
1.185	2	1 1 27	81.02
1.177	1	3 0 18+	81.74
1.175	2	2 0 26	81.96
1.145	4	1 3 10	84.52
1.138	1	1 2 23	85.16
1.133	3	3 1 11	85.66
1.113	1	0 2 28+	87.56
1.111	3	0 3 21+	87.82
1.094	2	3 1 14	89.54
1.090	3	1 1 30	89.96
1.087	4	4 0 1+	90.24
1.066	8	2 2 12	92.56
1.058	1	0 4 8	93.50
1.050	1	3 1 17	94.34
1.047	2	0 3 24+	94.72
1.037	1	1 0 34	96.00
1.031	1	0 2 31	96.70
1.016	2	2 2 21	98.62

Cobalt gadolinium, Co₇Gd₂ - continued

d(Å)	I	hkl	2θ(°)
λ = 1.540598Å			
1.007	2	1 1 33+	99.74
1.005	2	3 1 20	100.12
.9959	1	1 2 29	101.36
.9857	1	0 3 27+	102.78
.9660	1	2 2 24	105.76
.9622	3	3 2 10	106.36
.9582	2	0 2 34+	107.00
.9552	2	2 3 11	107.48
.9536	1	2 1 31	107.76
.9493	4	4 1 0	108.48
.9451	2	4 0 19	109.19
.9359	1	2 0 35+	110.78
.9328	2	0 4 20	111.34
.9313	2	2 3 14	111.60
.9289	3	3 0 30+	112.04
.9278	1	1 3 25	112.22
.9240	11	1 4 9+	112.96
.8953	2	2 1 34+	118.72
.8881	1	1 0 40	120.30
.8762	1	3 0 33+	123.08
.8744	2	2 3 20	123.52
.8726	2	1 1 39	124.00
.8712	2	2 2 30	124.30
.8686	2	3 1 29	124.96
.8588	1	1 4 18+	127.52
.8579	1	0 4 26	127.78
.8460	1	0 5 10	131.14
.8434	1	2 3 23	131.92
.8412	1	5 0 11	132.60
.8402	1	1 3 31	132.94
.8372	2	3 3 0+	133.88
.8320	2	4 1 21+	135.60
.8285	1	1 0 43	136.80
.8273	1	2 2 33	137.20
.8219	3	2 4 1	139.18
.8107	6	3 3 0	140.00
.8103	1	0 1 44	143.84
.8089	1	4 2 8	144.46
.8040	3	4 1 24+	146.72
.7994	1	1 3 34	148.98
.7977	1	4 2 11	150.00
.7968	1	4 0 31	150.38
.7943	1	2 1 40	151.74
.7861	4	2 2 36	156.98
.7831	1	0 3 39+	159.22

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°)
					λ = 1.540598Å
12.10	5	0	0	3	7.30
4.319	6	1	0	1	20.55
4.032	5	0	0	9	22.03
3.922	3	1	0	4	22.65
3.140	4	0	1	8	28.40
3.024	1	0	0	12	29.51
2.787	40	1	0	10	32.09
2.629	24	0	1	11	34.08
2.512	48	1	1	0	35.72
2.459	1	1	1	3	36.51
2.419	1	0	0	15	37.13
2.349	4	1	0	13	38.28
2.227	10	0	1	14	40.48
2.171	33	0	2	1	41.56
2.160	7	2	0	2	41.79
2.132	100	1	1	9	42.36
2.115	5	0	2	4	42.72
2.083	3	2	0	5	43.40
2.016	16	0	0	18	44.92
2.006	5	0	2	7	45.17
1.961	9	2	0	8	46.25
1.932	2	1	1	12	46.99
1.916	5	0	1	17	47.40
1.816	1	2	0	11	50.20
1.742	3	1	1	15	52.47
1.728	2	0	0	21	52.94
1.675	5	0	1	20	54.77
1.666	1	2	0	14	55.07
1.642	1	2	1	1	55.94
1.572	4	1	1	18	58.67
1.570	1	0	2	16	58.77
1.546	1	1	2	8	59.78
1.524	1	2	0	17	60.74
1.512	1	0	0	24	61.25
1.498	11	2	1	10	61.91
1.483	1	0	1	23	62.57
1.472	7	1	2	11	63.13
1.450	8	3	0	0	64.18
1.435	7	0	2	19	64.92
1.424	6	1	1	21	65.51
1.417	1	2	1	13	65.88
1.393	7	2	0	20	67.13
1.398	4	1	2	14	67.39
1.377	1	1	0	25	68.03
1.364	10	3	0	9	68.74
1.364	10	0	3	9	68.74
1.303	2	1	2	17	72.51
1.295	4	1	1	24	72.97
1.277	2	2	0	23	74.19
1.256	17	2	2	0	75.67

Cobalt gadolinium, Co₇Gd₂ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.218	3	1 2 20	78.43
1.210	1	0 0 30	79.11
1.203	1	0 1 29	79.66
1.199	1	2 2 9	79.95
1.185	3	1 1 27	81.09
1.175	2	2 0 26	81.95
1.145	4	1 3 10	84.57
1.138	1	1 2 23	85.16
1.133	3	3 1 11	85.66
1.113	1	0 2 28	87.55
1.111	1	0 3 21	87.81
1.111	1	3 0 21	87.81
1.094	2	3 1 14	89.54
1.090	4	1 1 30	89.95
1.088	1	2 1 25	90.13
1.087	3	4 0 1	90.25
1.066	9	2 2 18	92.55
1.058	1	0 4 8	93.50
1.050	1	3 1 17	94.34
1.047	1	3 0 24	94.79
1.047	1	0 3 24	94.79
1.037	1	1 0 34	95.99
1.031	1	0 2 31	96.71
1.016	2	2 2 21	98.62
1.008	1	0 0 36	99.66
1.007	2	1 1 33	99.76
1.005	2	3 1 20	100.12
.9958	2	1 2 29	101.35
.9857	1	3 0 27	102.79
.9857	1	0 3 27	102.79
.9660	2	2 2 24	105.76
.9622	3	3 2 10	106.36
.9584	1	3 1 23	106.98
.9582	2	0 2 34	107.01
.9552	2	2 3 11	107.49
.9536	1	2 1 31	107.76
.9493	5	4 1 0	108.48
.9451	2	4 0 19	109.19
.9359	1	2 0 35	110.78
.9328	2	0 4 20	111.34
.9313	1	2 3 14	111.60
.9289	1	3 0 30	112.05
.9289	1	0 3 30	112.05
.9278	1	1 3 25	112.24
.9240	7	4 1 9	112.95
.9240	7	1 4 9	112.95
.9041	1	2 3 17	116.87
.8954	1	0 4 23	118.69
.8953	2	2 1 34	118.73
.8881	1	1 0 40	120.30

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
.8762	1	0 3 33	123.07
.8762	1	3 0 33	123.07
.8744	2	2 3 20	123.50
.8726	1	1 1 39	123.97
.8712	3	2 2 30	124.30
.8686	2	3 1 29	124.97
.8588	1	4 1 18	127.51
.8588	1	1 4 18	127.51
.8579	1	0 4 26	127.78
.8460	2	0 5 10	131.14
.8434	1	2 3 23	131.93
.8423	1	2 1 37	132.27
.8412	1	5 0 11	132.60
.8402	1	1 3 31	132.94
.8372	3	3 3 0	133.89
.8331	1	4 0 28	135.23
.8320	2	4 1 21	135.59
.8320	2	1 4 21	135.59
.8285	2	1 0 43	136.79
.8273	2	2 2 33	137.21
.8248	1	5 0 14	138.11
.8224	1	3 2 25	139.00
.8219	6	2 4 1	139.19
.8212	1	4 2 2	139.43
.8197	9	3 3 9	140.02
.8187	1	2 4 4	140.39
.8119	1	2 4 7	143.14
.8103	2	0 1 44	143.83
.8089	1	4 2 8	144.45
.8040	3	1 4 24	146.72
.8040	3	4 1 24	146.72
.7994	3	1 3 34	148.98
.7977	1	4 2 11	149.88
.7968	2	4 0 31	150.39
.7943	4	2 1 49	151.73
.7911	1	3 3 15	153.64
.7868	1	0 2 43	156.49
.7861	12	2 2 36	156.98
.7845	2	5 0 20	158.17
.7836	1	4 2 14	158.84
.7831	2	0 3 39	159.23
.7831	2	3 0 39	159.23

Cobalt gallium manganese, Co₂GaMn

Structure

Cubic, Fm3m(225), Z=4, Heusler alloy, type L2₁, from powder data (x-ray and neutron) [Webster, 1971].

Lattice constant: [ibid.]

$$a = 5.770\text{\AA}$$

Density

(calculated) 8.383 g/cm³

Thermal parameters

Isotropic: overall B = 1.0

Scattering factors

Co⁰, Ga⁰, Mn⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.851 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 12.4$$

References

- Cromer, D. T. and Webster, J. B. (1968). Acta Crystallogr. A24, 321.
Webster, P.J. (1971). J. Phys. Chem. Solids, 32, 1221.

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
3.331	1	1	1	1	26.74
2.040	100	2	2	0	44.37
1.4425	13	4	0	0	64.55
1.1778	23	4	2	2	81.69
1.0200	7	4	4	0	98.08
.9123	11	6	2	0	115.20
.8328	4	4	4	4	135.31

Calculated Pattern (Peak heights)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
3.331	1	1	1	1	26.74
2.040	100	2	2	0	44.38
1.4425	12	4	0	0	64.56
1.1778	20	4	2	2	81.70
1.0200	6	4	4	0	98.08
.9123	8	6	2	0	115.20
.8328	2	4	4	4	135.32

Cobalt gallium tantalum, Co₂GaTa

Structure

Cubic, Fm3m(225), Z=4, Heusler alloy, type L2₁,
from powder data [Markiv et al., 1965].

Lattice constant: [ibid.]

$$a = 5.923\text{\AA}$$

Density

(calculated) 11.780 g/cm³

Thermal parameters

Isotropic: overall B = 1.0

Scattering factors

Co⁰, Ga⁰, Ta⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 1.56 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 18.7$$

References

- Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Markiv, V. Ya., Voroshilov, Yu.V., Kripyakevich, P.I., and Cherkashin, E. E. (1965). Sov. Phys. Crystallogr. 9, 619.

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
3.420	21	1	1	1	26.04
2.961	16	2	0	0	30.15
2.094	100	2	2	0	43.17
1.786	10	3	1	1	51.10
1.710	4	2	2	2	53.55
1.481	14	4	0	0	62.69
1.359	4	3	3	1	69.07
1.324	5	4	2	0	71.13
1.209	26	4	2	2	79.16
1.140	2	5	1	1	85.03
1.140	1	3	3	3	85.03
1.047	8	4	4	0	94.73
1.001	3	5	3	1	100.60
.9872	2	4	4	2	102.58
.9365	13	6	2	0	110.68
.9032	1	5	3	3	117.04
.8929	2	6	2	2	119.24
.8549	4	4	4	4	128.58
.8294	2	5	5	1	136.48
.8294	2	7	1	1	136.48
.8214	3	6	4	0	139.38
.7915	43	6	4	2	153.42

Calculated Pattern (Peak heights)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
3.420	24	1	1	1	26.04
2.961	18	2	0	0	30.16
2.094	100	2	2	0	43.16
1.786	10	3	1	1	51.10
1.710	4	2	2	2	53.56
1.481	13	4	0	0	62.70
1.359	3	3	3	1	69.06
1.324	5	4	2	0	71.12
1.209	22	4	2	2	79.16
1.140	2	5	1	1+	85.02
1.047	6	4	4	0	94.74
1.001	3	5	3	1	100.60
.9872	2	4	4	2+	102.58
.9365	10	6	2	0	110.68
.9032	1	5	3	3	117.04
.8929	2	6	2	2	119.24
.8549	3	4	4	4	128.58
.8294	2	7	1	1+	136.48
.8214	1	6	4	0	139.38
.7915	16	6	4	2	153.42

Cobalt gallium titanium, Co_2GaTi

Structure

Cubic, $\text{Fm}\bar{3}\text{m}$, $Z=4$, Heusler alloy, type L2_1 , from powder data (x-ray and neutron) [Webster and Ziebeck, 1973].

Lattice constant: [ibid.]

$$a = 5.848\text{\AA}$$

Density

(calculated) 7.818 g/cm^3

Thermal parameters

Isotropic: overall $B = 1.0$

Scattering factors

Co^0 , Ga^0 , Ti^0 [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.885 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 11.9$$

References

- Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Webster, P. J. and Ziebeck, K. R. A. (1973). J. Phys. Chem. Solids, 34, 1647.

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°)
λ = 1.540598Å					
3.376	3	1	1	1	26.3 ^a
2.068	100	2	2	0	43.75
1.7632	2	3	1	1	51.8 ^a
1.4620	13	4	0	0	63.5 ^a
1.1937	23	4	2	2	80.3 ^a
1.0338	7	4	4	0	96.34
.9246	11	6	2	0	112.83
.8441	4	4	4	4	131.73

Calculated Pattern (Peak heights)					
$d(\text{\AA})$	I	hkl			$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$					
3.376	3	1	1	1	26.3 ^a
2.068	100	2	2	0	43.76
1.7632	1	3	1	1	51.8 ^a
1.4620	12	4	0	0	63.60
1.1937	20	4	2	2	80.3 ^a
1.0338	6	4	4	0	96.34
.9246	8	6	2	0	112.84
.8441	2	4	4	4	131.7 ^a

Cobalt gallium vanadium, Co₂GaV

Structure

Cubic, Fm3m(225), Z=4, Heusler alloy, type L2₁, from powder data (x-ray and neutron) [Ziebeck and Webster, 1974].

Lattice constant: [ibid.]

$$a = 5.786\text{\AA}$$

Density

(measured) 8.15 g/cm³ [ibid.]

(calculated) 8.177 g/cm³

Thermal parameters

Isotropic: overall B = 1.0

Scattering factors

Co⁰, Ga⁰, V⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.893 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 12.0$$

References

Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.

Ziebeck, K. R. A. and Webster, P. J. (1974). J. Phys. Chem. Solids, 35, 1.

Calculated Pattern (Integrated)					
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å	
3.341	2	1 1 1	26.66		
2.046	100	2 2 0	44.24		
1.7445	1	3 1 1	52.41		
1.4465	13	4 0 0	64.35		
1.1811	23	4 2 2	81.42		
1.0228	7	4 4 0	97.72		
.9148	11	6 2 0	114.70		
.8351	4	4 4 4	134.55		

Calculated Pattern (Peak heights)					
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å	
3.341	3	1 1 1	26.68		
2.046	100	2 2 0	44.24		
1.7445	1	3 1 1	52.40		
1.4465	12	4 0 0	64.36		
1.1811	20	4 2 2	81.42		
1.0228	6	4 4 0	97.72		
.9148	8	6 2 0	114.70		
.8351	2	4 4 4	134.54		

Cobalt germanium manganese, Co_2GeMn

Structure

Cubic, $\text{Fm}\bar{3}\text{m}(225)$, $Z=4$, Heusler alloy, type L2_1 , from powder data (x-ray and neutron) [Webster, 1971].

Lattice constant: [ibid.]

$$a = 5.743\text{\AA}$$

Density

(calculated) 8.594 g/cm^3

Thermal parameters

Isotropic: overall $B = 1.0$

Scattering factors

Co^0 , Ge^0 , Mn^0 [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.851 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 12.4$$

References

- Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Webster, P.J. (1971). J. Phys. Chem. Solids, 32, 1221.

Calculated Pattern (Integrated)

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$			
3.316	2	1 1 1	26.87
2.0305	100	2 2 0	44.59
1.7316	1	3 1 1	52.83
1.4357	13	4 0 0	64.89
1.1723	23	4 2 2	82.16
1.0152	7	4 4 0	98.71
.9080	12	6 2 0	116.05
.8289	4	4 4 4	136.64

Calculated Pattern (Peak heights)

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$			
3.316	2	1 1 1	26.88
2.0305	100	2 2 0	44.60
1.7316	1	3 1 1	52.82
1.4357	12	4 0 0	64.90
1.1723	20	4 2 2	82.16
1.0152	6	4 4 0	98.70
.9080	9	6 2 0	116.06
.8289	2	4 4 4	136.64

Cobalt germanium titanium, Co_2GeTi

Structure

Cubic, $\text{Fm}\bar{3}\text{m}(225)$, $Z=4$, Heusler alloy, type L2_1 , from powder data (Gladyshevskii et al., 1963).

Lattice constant: [ibid.]

$$a = 5.823\text{\AA}$$

Density

(calculated) 8.018 g/cm^3

Thermal parameters

Isotropic: overall $B = 1.0$

Scattering factors

Co^0 , Ge^0 , Ti^0 [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.884 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 12.0$$

References

Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. **A24**, 321.

Gladyshevskii, E. I., Markiv, V. Ya., Kuz'ma, Yu. B. and Cherkashin, E. E. (1963). Titan Ego Splavy, No. 10, 71.

Calculated Pattern (Integrated)					
$d(\text{\AA})$	I	hkl			$2\theta(^{\circ})$ $\lambda = 1.540598\text{\AA}$
3.362	3	1	1	1	26.49
2.059	100	2	2	0	43.94
1.756	2	3	1	1	52.05
1.456	13	4	0	0	63.90
1.189	23	4	2	2	80.79
1.029	7	4	4	0	96.89
.921	11	6	2	0	113.58
.840	4	4	4	4	132.84

Calculated Pattern (Peak heights)					
$d(\text{\AA})$	I	hkl			$2\theta(^{\circ})$ $\lambda = 1.540598\text{\AA}$
3.362	4	1	1	1	26.50
2.059	100	2	2	0	43.94
1.756	2	3	1	1	52.04
1.456	12	4	0	0	63.90
1.189	20	4	2	2	80.80
1.029	6	4	4	0	96.90
.921	9	6	2	0	113.58
.840	2	4	4	4	132.84

Cobalt indium, CoIn₃

Structure

Tetragonal, P4/mbm(127), Z=2. The structure was determined by Stadelmaier et al. [1973].

Lattice constants: [ibid.]

$$a = 6.830 \text{ \AA}$$

$$c = 3.547$$

Density

(measured) 8.09 g/cm³ [ibid.]

(calculated) 8.097 g/cm³

Thermal parameters

Isotropic [Stadelmaier et al., op. cit.].

Scattering factors

Co⁰, In⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.451 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 7.04$$

References

- Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. **A24**, 321.
 Stadelmaier, H. H., Schöbel, J. D., Jones, R. A., and Shumaker, C. A. (1973). Acta Crystallogr. **B29**, 2926.

Calculated Pattern (Peak heights)

d(Å)	I	hkl	2θ(°)
λ = 1.540598 Å			
4.830	2	1 1 0	18.36
3.547	2	0 0 1	25.10
3.054	39	2 1 0	29.22
2.859	53	1 1 1	31.26
2.460	38	2 0 1	36.50
2.415	13	2 2 0	37.20
2.315	100	2 1 1	38.88
2.160	59	3 1 0	41.80
1.996	5	2 2 1	45.40
1.773	15	0 0 2	51.48
1.707	2	4 0 0	53.64
1.671	3	3 2 1	54.90
1.657	3	4 1 0	55.42
1.610	3	3 3 0	57.18
1.539	12	4 0 1	60.10
1.534	12	2 1 2	60.28
1.527	9	4 2 0	60.58
1.501	9	4 1 1	61.76
1.466	13	3 3 1	63.40
1.429	3	2 2 2	65.22

d(Å)	I	hkl	2θ(°)
λ = 1.540598 Å			
1.403	1	4 2 1	66.62
1.371	19	3 1 2	68.38
1.339	1	5 1 0	70.22
1.268	2	5 2 0	74.80
1.253	3	5 1 1	75.86
1.230	1	4 0 2	77.54
1.211	2	4 1 2	79.04
1.207	4	4 4 0	79.28
1.194	11	5 2 1	80.34
1.192	7	3 3 2	80.56
1.171	1	5 3 0	82.24
1.157	4	4 2 2	83.46
1.148	2	1 1 3	84.24
1.138	4	6 0 0	85.18
1.117	2	2 0 3	87.18
1.112	1	5 3 1	87.66
1.103	5	2 1 3	88.64
1.070	1	6 1 1	92.04
1.069	1	5 1 2	92.22
1.067	1	5 4 0	92.48
1.033	3	6 2 1	96.42
1.032	2	5 2 2	96.60
1.021	3	5 4 1	97.90
.9981	3	4 4 2	101.04
.9774	1	5 3 2	104.02
.9720	2	4 0 3	104.84
.9623	2	4 1 3	106.34
.9580	5	6 0 2	107.04
.9529	2	3 3 3	107.86
.9320	3	7 1 1+	111.48
.9151	3	6 4 1	114.66
.9141	1	5 4 2	114.84
.9070	1	7 2 1	116.28
.8968	4	7 3 0	118.40
.8867	2	0 0 4+	120.64
.8648	4	5 2 3	125.92
.8516	1	2 1 4	129.52
.8491	1	6 5 1	130.24
.8472	1	8 1 0	130.84
.8324	1	2 2 4+	135.48
.8240	3	8 1 1	138.40
.8203	4	3 1 4	139.78
.8049	2	6 6 0	146.28
.8003	7	7 3 2	148.52
.7974	2	6 2 3	150.04
.7920	2	5 4 3	153.12

Cobalt indium, CoIn_3 - continued

Calculated Pattern (Integrated)				
$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$ $\lambda = 1.540598\text{\AA}$	
4.830	1	1 1 0	18.36	
3.547	2	0 0 1	25.09	
3.054	37	2 1 0	29.21	
2.859	52	1 1 1	31.26	
2.460	38	2 0 1	36.49	
2.415	13	2 2 0	37.20	
2.315	100	2 1 1	38.88	
2.160	60	3 1 0	41.70	
1.996	5	2 2 1	45.40	
1.773	16	0 0 2	51.49	
1.707	2	4 0 0	53.63	
1.671	3	3 2 1	54.90	
1.657	3	4 1 0	55.42	
1.610	4	3 3 0	57.17	
1.539	12	4 0 1	60.09	
1.534	7	2 1 2	60.30	
1.527	8	4 2 0	60.59	
1.501	10	4 1 1	61.76	
1.466	14	3 3 1	63.40	
1.429	4	2 2 2	65.22	
1.403	2	4 2 1	66.62	
1.371	22	3 1 2	68.39	
1.339	1	5 1 0	70.21	
1.268	2	5 2 0	74.80	
1.253	3	5 1 1	75.86	
1.230	1	4 0 2	77.55	
1.211	2	4 1 2	79.03	
1.207	3	4 4 0	79.28	
1.194	12	5 2 1	80.33	
1.192	2	3 3 2	80.51	
1.171	2	5 3 0	82.24	
1.157	5	4 2 2	83.46	
1.148	2	1 1 3	84.25	
1.138	5	6 0 0	85.17	
1.117	2	2 0 3	87.17	
1.112	1	5 3 1	87.67	
1.103	6	2 1 3	88.63	
1.070	2	6 1 1	92.04	
1.069	1	5 1 2	92.22	
1.067	1	5 4 0	92.47	
1.033	3	6 2 1	96.42	
1.032	2	5 2 2	96.61	
1.021	4	5 4 1	97.89	
.9981	4	4 4 2	101.03	
.9774	2	5 3 2	104.02	
.9720	2	4 0 3	104.83	
.9623	2	4 1 3	106.34	
.9580	6	6 0 2	107.04	
.9529	3	3 3 3	107.87	
.9320	3	7 1 1	111.40	

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$ $\lambda = 1.540598\text{\AA}$
.9151	4	6 4 1	114.66
.9141	1	5 4 2	114.85
.9070	1	7 2 1	116.27
.8968	6	7 3 0	118.39
.8867	2	0 0 4	120.61
.8864	1	5 1 3	120.69
.8648	7	5 2 3	125.92
.8516	1	2 1 4	129.52
.8491	2	6 5 1	130.25
.8472	1	8 1 0	130.81
.8324	1	2 2 4	135.46
.8240	6	8 1 1	138.41
.8240	1	7 4 1	138.41
.8203	8	3 1 4	139.78
.8142	1	6 1 3	142.20
.8066	2	8 2 1	145.51
.8049	3	6 6 0	146.27
.8003	17	7 3 2	148.52
.7974	4	6 2 3	150.05
.7920	5	5 4 3	153.12
.7870	1	4 0 4	156.39

Cobalt lanthanum, CoLa_3

Structure

Orthorhombic, $\text{Pnma}(62)$, $Z=4$, isostructural with CFe_3 , type DO_{11} . The structure was determined by Cromer and Larson [1961].

Lattice constants: [ibid.]

$$a = 7.279\text{\AA}$$

$$b = 10.089$$

$$c = 6.578$$

(published value $b = 10.088$)

Density

(measured) 6.48 g/cm^3 [ibid.]

(calculated) 6.539 g/cm^3

Thermal parameters

Isotropic (Cromer and Larson, op. cit.).

Scattering factors

Co^0 , La^0 [Forsyth and Wells, 1959].

Scale factors (integrated intensities)

$$\gamma = 0.200 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 4.15$$

References

- Cromer, D. T. and Larson, A. C. (1961). Acta Crystallogr. 14, 1226.
Forsyth, J.B. and Wells, M. (1959). Acta Crystallogr. 12, 412.

Calculated Pattern (Peak heights)

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$			
5.51	1	0 1 1	16.08
4.88	2	1 0 1	18.18
4.39	1	1 1 1	20.20
3.508	15	1 2 1	25.38
3.424	14	2 1 0	26.02
3.289	10	0 0 2	27.10
3.185	14	2 0 1	28.00
3.037	31	2 1 1	29.40
2.994	100	0 3 1+	29.82
2.952	33	2 2 0	30.26
2.873	51	1 1 2	31.10
2.769	32	1 3 1	32.30
2.693	45	2 2 1	33.24
2.577	12	1 2 2	34.80
2.522	5	0 4 0	35.56
2.470	16	2 3 0	36.34
2.440	2	2 0 2	36.80
2.372	8	2 1 2	37.90
2.312	1	2 3 1	38.92
2.276	13	3 0 1	39.56

$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$
$\lambda = 1.540598\text{\AA}$			
2.241	3	1 4 1+	40.26
2.221	6	3 1 1	40.60
2.197	3	2 2 2	41.06
2.073	4	2 4 0	43.62
2.055	3	1 1 3	44.02
1.977	5	2 4 1+	45.88
1.953	1	3 0 2	46.48
1.938	15	1 2 3	46.84
1.885	3	3 3 1	48.24
1.878	3	2 0 3	48.38
1.821	2	3 2 2+	50.06
1.781	14	1 3 3	51.26
1.765	6	2 5 0	51.76
1.760	5	2 2 3	51.90
1.754	11	4 0 1	52.10
1.728	2	4 1 1	52.94
1.712	1	4 2 0	53.50
1.704	7	2 5 1	53.74
1.689	15	3 3 2	54.28
1.681	11	0 6 0	54.52
1.674	12	1 5 2	54.80
1.645	2	0 0 4	55.86
1.640	2	2 3 3	56.02
1.627	9	3 0 3	56.52
1.614	4	1 4 3	57.02
1.604	8	1 0 4	57.40
1.600	12	4 3 0	57.54
1.584	1	1 1 4	58.18
1.555	4	2 5 2	59.38
1.548	2	3 2 3	59.66
1.544	2	3 4 2	59.84
1.529	1	1 2 4	60.52
1.518	1	4 2 2	60.96
1.510	1	3 5 1	61.34
1.506	1	2 4 3	61.50
1.4972	1	0 6 2	61.92
1.4869	1	2 6 1	62.40
1.4824	2	2 1 4	62.60
1.4665	3	1 6 2	63.38
1.4548	3	1 5 3	63.94
1.4478	1	1 3 4	64.28
1.4391	3	4 3 2+	64.72
1.4366	3	2 2 4	64.86
1.4075	1	5 1 1+	66.36
1.4032	2	3 5 2	66.58
1.3681	1	5 2 1	68.54
1.3525	3	3 6 1	69.44
1.3493	2	4 2 3+	69.62
1.3237	1	4 5 1	71.18
1.3143	1	3 2 4+	71.78

Cobalt lanthanum, CoLa₃ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.2989	1	1 7 2	72.74
1.2872	4	5 2 2	73.52
1.2618	5	3 3 4	75.24
1.2557	1	1 5 4	75.68
1.2528	1	2 6 3	75.90
1.2378	1	5 3 2+	76.98
1.2280	1	2 1 5	77.70
1.2252	4	0 3 5	77.92
1.2177	1	3 7 1	78.48
1.2138	4	4 6 1+	78.80
1.2082	1	1 3 5	79.22
1.2042	3	5 1 3+	79.54
1.2016	3	2 2 5	79.76
1.1916	1	2 8 0	80.54
1.1848	1	6 1 1+	81.10
1.1795	2	6 2 0+	81.56
1.1773	2	5 4 2+	81.74
1.1725	3	2 8 1	82.14
1.1692	4	3 6 3	82.42
1.1624	2	1 8 2+	83.04
1.1607	3	1 6 4+	83.16
1.1470	1	4 3 4	84.38
1.1412	1	6 3 0+	84.92
1.1382	1	6 0 2	85.18
1.1285	1	3 5 4	86.10
1.1273	1	3 2 5	86.20
1.1244	2	6 3 1	86.48
1.1051	1	0 9 1	88.38
1.0963	1	0 0 6	89.28
1.0896	1	1 5 5+	89.96
1.0811	2	1 8 3	90.88
1.0785	1	6 4 1+	91.16
1.0662	1	4 0 5+	92.54
1.0548	1	2 5 5	93.82
1.0441	1	2 1 6	95.08
1.0395	3	5 5 3+	95.64

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
5.51	1	0	1	1	16.07
5.04	1	0	2	0	17.57
4.88	2	1	0	1	18.16
4.39	1	1	1	1	20.20
3.508	20	1	2	1	25.37
3.424	18	2	1	0	26.01
3.289	13	0	0	2	27.09
3.185	19	2	0	1	28.00
3.037	41	2	1	1	29.39
2.997	43	1	0	2	29.78
2.994	100	0	3	1	29.81
2.952	43	2	2	0	30.26
2.873	72	1	1	2	31.10
2.769	44	1	3	1	32.30
2.755	10	0	2	2	32.47
2.693	66	2	2	1	33.24
2.577	17	1	2	2	34.79
2.522	7	0	4	0	35.56
2.470	23	2	3	0	36.34
2.440	2	2	0	2	36.80
2.372	12	2	1	2	37.90
2.312	1	2	3	1	38.92
2.276	19	3	0	1	39.56
2.241	2	1	4	1	40.21
2.238	2	1	3	2	40.27
2.221	9	3	1	1	40.59
2.197	4	2	2	2	41.06
2.073	6	2	4	0	43.63
2.055	4	1	1	3	44.02
1.977	6	2	4	1	45.86
1.975	2	2	3	2	45.91
1.953	1	3	0	2	46.47
1.938	23	1	2	3	46.83
1.929	3	0	5	1	47.07
1.885	4	3	3	1	48.24
1.878	2	2	0	3	48.43
1.821	2	3	2	2	50.05
1.820	1	4	0	0	50.09
1.781	21	1	3	3	51.26
1.765	8	2	5	0	51.76
1.760	1	2	2	3	51.91
1.754	16	4	0	1	52.11
1.728	4	4	1	1	52.95
1.712	1	4	2	0	53.49
1.704	11	2	5	1	53.74
1.690	1	3	4	1	54.24
1.689	22	3	3	2	54.28
1.681	12	0	6	0	54.53
1.674	17	1	5	2	54.80
1.645	3	0	0	4	55.36

Cobalt lanthanum, CoLa₃ - continued

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.640	1	2 3 3	56.04
1.627	14	3 0 3	56.52
1.614	6	1 4 3	57.03
1.606	2	3 1 3	57.32
1.604	9	1 0 4	57.40
1.600	14	4 3 0	57.54
1.584	1	1 1 4	58.19
1.555	6	2 5 2	59.39
1.548	2	3 2 3	59.67
1.544	1	3 4 2	59.86
1.529	2	1 2 4	60.52
1.518	2	4 2 2	60.97
1.510	2	3 5 1	61.35
1.506	1	2 4 3	61.51
1.4972	1	0 6 2	61.93
1.4869	2	2 6 1	62.40
1.4824	2	2 1 4	62.62
1.4665	5	1 6 2	63.37
1.4548	5	1 5 3	63.94
1.4478	2	1 3 4	64.29
1.4391	4	4 3 2	64.72
1.4366	3	2 2 4	64.85
1.4079	1	0 7 1	66.34
1.4075	1	5 1 1	66.36
1.4032	2	3 5 2	66.59
1.3681	1	5 2 1	68.53
1.3525	4	3 6 1	69.43
1.3493	1	4 2 3	69.62
1.3237	2	4 5 1	71.17
1.3143	1	3 2 4	71.76
1.2989	2	1 7 2	72.75
1.2884	1	2 4 4	73.44
1.2872	7	5 2 2	73.52
1.2618	8	3 3 4	75.25
1.2557	1	1 5 4	75.68
1.2528	1	2 6 3	75.89
1.2378	1	5 3 2	76.97
1.2372	1	2 0 5	77.01
1.2280	2	2 1 5	77.70
1.2252	5	0 3 5	77.91
1.2210	1	1 8 1	78.23
1.2177	1	3 7 1	78.48
1.2138	6	4 6 1	78.78
1.2132	1	6 0 0	78.83
1.2128	1	5 0 3	78.86
1.2082	1	1 3 5	79.22
1.2045	1	6 1 0	79.51
1.2042	4	5 1 3	79.54
1.2031	1	2 5 4	79.62
1.2016	1	2 2 5	79.74

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
1.1916	2	2 8 0	80.55
1.1848	1	6 1 1	81.11
1.1795	1	6 2 0	81.54
1.1775	1	0 8 2	81.71
1.1773	2	5 4 2	81.73
1.1757	1	0 6 4	81.87
1.1725	5	2 8 1	82.14
1.1692	6	3 6 3	82.42
1.1624	2	1 8 2	83.01
1.1620	1	5 5 1	83.04
1.1607	3	1 6 4	83.16
1.1470	2	4 3 4	84.38
1.1412	1	6 3 0	84.91
1.1382	1	6 0 2	85.18
1.1285	1	3 5 4	86.09
1.1273	1	3 2 5	86.21
1.1244	3	6 3 1	86.48
1.1051	2	0 9 1	88.38
1.0963	1	0 0 6	89.27
1.0925	1	1 9 1	89.67
1.0900	1	5 0 4	89.93
1.0896	1	1 5 5	89.97
1.0811	3	1 8 3	90.88
1.0662	2	4 0 5	92.52
1.0655	1	5 2 4	92.60
1.0548	2	2 5 5	93.82
1.0441	1	2 1 6	95.08
1.0397	1	6 5 0	95.61
1.0395	4	5 5 3	95.64

Cobalt lutetium, Co₂Lu

Structure

Cubic, Fd3m(227), Z=8, isostructural with Cu₂Mg, type C15, from powder data [Lemaire, 1971].

Lattice constant: [ibid.]

$$a = 7.102\text{\AA}$$

Density

(calculated) 10.860 g/cm³

Thermal parameters

Isotropic: overall B = 1.0

Scattering factors

Co⁰, Lu⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 1.08 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 14.5$$

References

Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.

Lemaire, F. G. R. (1971). Solid State Commun. 9, 341.

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
4.100	20	1	1	1	21.66
2.511	73	2	2	0	35.73
2.141	100	3	1	1	42.17
2.050	12	2	2	2	44.14
1.775	1	4	0	0	51.42
1.629	5	3	3	1	56.43
1.450	23	4	2	2	64.10
1.367	20	5	1	1	68.61
1.367	7	3	3	3	68.61
1.255	18	4	4	0	75.60
1.200	4	5	3	1	79.83
1.123	9	6	2	0	86.62
1.083	8	5	3	3	90.67
1.071	3	6	2	2	92.02
.9945	1	5	5	1	101.53
.9945	1	7	1	1	101.53
.9490	12	6	4	2	108.52
.9246	6	5	5	3	112.84
.9246	12	7	3	1	112.84
.8877	4	8	0	0	120.38
.8676	1	7	3	3	125.20
.8370	7	8	2	2	133.95
.8370	3	6	6	0	133.95
.8201	15	7	5	1	139.87
.8201	2	5	5	5	139.87
.8147	2	6	6	2	142.01
.7940	2	8	4	0	151.92

Calculated Pattern (Peak heights)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
4.100	24	1	1	1	21.66
2.511	77	2	2	0	35.74
2.141	100	3	1	1	42.18
2.050	13	2	2	2	44.14
1.775	1	4	0	0	51.42
1.629	5	3	3	1	56.44
1.450	21	4	2	2	64.20
1.367	23	5	1	1+	68.60
1.255	16	4	4	0	75.70
1.200	3	5	3	1	79.84
1.123	8	6	2	0	86.62
1.083	7	5	3	3	90.68
1.071	2	6	2	2	92.02
.9945	2	7	1	1+	101.54
.9490	10	6	4	2	108.52
.9246	14	7	3	1+	112.84
.8877	3	8	0	0	120.38
.8676	1	7	3	3	125.20
.8370	6	8	2	2+	133.94
.8201	8	7	5	1+	139.88
.8147	1	6	6	2	142.02
.7940	1	8	4	0	151.92

Cobalt neodymium, Co₂Nd

Structure

Cubic, Fd3m (227), Z = 8, isostructural with Cu₂Mg, type C15, from powder data [Harris et al., 1965].

Lattice constant: [ibid.]

$$a = 7.2986 \text{ \AA}$$

(published value 7.2834 kX)

Density

(calculated) 8.955 g/cm³

Thermal parameters

Isotropic: overall B = 1.0

Scattering factors

Co⁰, Nd⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.571 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 12.2$$

References

- Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Harris, I. R., Mansey, R. C., and Raynor, G. V. (1965). J. Less-Common Metals, 9, 270.

Calculated Pattern (Integrated)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
4.214	11	1 1 1	21.07
2.5804	63	2 2 0	34.74
2.2006	100	3 1 1	40.98
2.1069	17	2 2 2	42.89
1.6744	2	3 3 1	54.78
1.4898	19	4 2 2	62.27
1.4046	19	5 1 1	66.52
1.4046	6	3 3 3	66.52
1.2902	18	4 4 0	73.31
1.2337	2	5 3 1	77.27
1.1540	7	6 2 0	83.75
1.1130	8	5 3 3	87.59
1.1003	4	6 2 2	88.87
.9753	10	6 4 2	104.33
.9502	11	7 3 1	108.32
.9502	5	5 5 3	108.32
.9123	3	8 0 0	115.20
.8601	2	6 6 0	127.16
.8601	5	8 2 2	127.16
.8428	12	7 5 1	132.13
.8428	2	5 5 5	132.13
.8372	3	6 6 2	133.88
.8011	2	7 5 3	148.11
.8011	1	9 1 1	148.11
.7780	13	6 6 4	163.83

Calculated Pattern (Peak heights)

d(Å)	I	hkl	2θ(°) λ = 1.540598Å
4.214	12	1 1 1	21.08
2.5804	65	2 2 0	34.74
2.2006	100	3 1 1	40.98
2.1069	16	2 2 2	42.90
1.6744	2	3 3 1	54.78
1.4898	17	4 2 2	62.26
1.4046	23	5 1 1+	66.52
1.2902	16	4 4 0	73.32
1.2337	2	5 3 1	77.28
1.1540	6	6 2 0	83.74
1.1130	6	5 3 3	87.58
1.1003	3	6 2 2	88.86
1.0220	1	7 1 1+	97.82
.9753	7	6 4 2	104.34
.9502	12	7 3 1+	108.32
.9123	2	8 0 0	115.20
.8601	4	8 2 2+	127.16
.8428	7	7 5 1+	132.14
.8372	1	6 6 2	133.88
.8011	1	7 5 3+	148.10
.7780	3	6 6 4	163.82

Cobalt nickel tin, $\text{Co}_{.75}\text{Ni}_{.75}\text{Sn}_{.75}$

Structure

Hexagonal, $P6_3/mmc$ (194), $Z=2$, from powder data.
The atoms were assigned these positions: 1.5 Co plus 0.5 Ni in 2a; 1.5 Sn plus 0.5 Ni in 2c; and only 0.5 Ni in 2d. [Castelliz, 1953].

Lattice constants: [ibid.]

$$a = 4.095 \text{ \AA}$$

$$c = 5.209$$

Density

(calculated) 7.781 g/cm^3

Thermal parameters

Isotropic: overall $B = 1.0$

Scattering factors

Co^0 , Ni^0 , Sn^0 [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.453 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 6.08$$

References

Castelliz, L. (1953). *Monatsh. Chem.* 84, 49.
Cromer, D. T. and Mann, J. B. (1968). *Acta Crystallogr.* A24, 321.

Calculated Pattern (Peak heights)				
$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$	$\lambda = 1.540598 \text{ \AA}$
2.931	100	1 0 1	30.48	
2.604	7	0 0 2	34.42	
2.099	87	1 0 2	43.06	
2.048	88	1 1 0	44.20	
1.679	14	2 0 1	54.64	
1.610	7	1 1 2	57.18	
1.559	11	1 0 3	59.20	
1.466	20	2 0 2	63.40	
1.302	5	0 0 4	72.54	
1.298	11	2 1 1	72.80	
1.241	4	2 0 3	76.76	
1.192	16	2 1 2	80.52	
1.182	9	3 0 0	81.32	
1.099	14	1 1 4	89.02	
1.076	2	3 0 2	91.39	
1.061	5	2 1 3	93.10	
1.024	5	2 2 0	97.60	
1.000	2	1 0 5	100.82	
.967	3	3 1 1	105.68	
.953	1	2 2 2	107.90	
.920	7	3 1 2	113.68	
.898	2	2 0 5	118.08	
.875	7	3 0 4	123.30	
.874	2	4 0 1	123.60	
.856	3	3 1 3	128.34	
.843	3	1 0 6	131.98	
.839	3	4 0 2	133.22	
.823	3	2 1 5	138.94	
.805	7	2 2 4	146.32	
.804	3	3 2 1	146.78	
.799	1	1 1 6	149.04	
.790	1	4 0 3	154.60	

Cobalt nickel tin, Co_{.75}Ni_{.75}Sn_{.75} - continued

Calculated Pattern (Integrated)				
d(Å)	I	hkl	2θ(°)	λ = 1.540598Å
2.931	100	1 0 1	30.47	
2.604	7	0 0 2	34.41	
2.099	93	1 0 2	43.06	
2.048	95	1 1 0	44.20	
1.679	17	2 0 1	54.63	
1.610	9	1 1 2	57.18	
1.559	13	1 0 3	59.20	
1.466	24	2 0 2	63.41	
1.302	6	0 0 4	72.53	
1.298	12	2 1 1	72.80	
1.241	5	2 0 3	76.77	
1.192	21	2 1 2	80.53	
1.182	11	3 0 0	81.33	
1.099	17	1 1 4	89.02	
1.076	2	3 0 2	91.38	
1.061	6	2 1 3	93.10	
1.024	7	2 2 0	97.60	
1.000	3	1 0 5	100.82	
.967	5	3 1 1	105.60	
.953	2	2 2 2	107.80	
.920	10	3 1 2	113.68	
.898	2	2 0 5	118.09	
.875	12	3 0 4	123.30	
.874	2	4 0 1	123.61	
.856	5	3 1 3	128.34	
.843	6	1 0 6	131.98	
.839	6	4 0 2	133.21	
.823	6	2 1 5	138.93	
.805	17	2 2 4	146.31	
.804	7	3 2 1	146.78	
.799	2	1 1 6	149.05	
.790	4	4 0 3	154.60	

Cobalt samarium, Co₅Sm

Structure

Hexagonal, P6/mmm(191), Z=1, isostructural with CaCu₅, type D2_d, from powder data [Khan and Feldmann, 1973].

Lattice constants: [ibid.]

$$a = 4.997\text{\AA}$$

$$c = 3.978$$

Density

(measured) 8.58 g/cm³ [ibid.]
(calculated) 8.590 g/cm³

Thermal parameters

Isotropic: overall B = 2.0

Scattering factors

Co⁰, Sm⁰ [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.320 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 6.73$$

References

Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Khan, Y. and Feldmann, D. (1973). J. Less-Common Metals, 31, 111.

Calculated Pattern (Peak heights)				
d (Å)	I	hkl	2θ (°)	λ = 1.540598 Å
4.328	2	1 0 0	20.52	
3.978	8	0 0 1	22.34	
2.929	60	1 0 1	30.50	
2.498	36	1 1 0	35.92	
2.164	36	2 0 0	41.72	
2.116	100	1 1 1	42.70	
1.989	24	0 0 2	45.58	
1.901	5	2 0 1	47.82	
1.556	11	1 1 2	59.34	
1.513	10	2 1 1	61.22	
1.464	14	2 0 2	63.48	
1.443	4	3 0 0	64.56	
1.356	14	3 0 1	69.22	
1.268	2	1 0 3	74.84	
1.249	9	2 2 0	76.14	
1.171	7	1 1 3	82.24	
1.168	6	3 0 2	82.52	
1.149	3	3 1 1	84.20	
1.082	2	4 0 0	90.80	
1.058	8	2 2 2	93.46	
1.030	2	2 1 3	96.80	
.994	1	0 0 4	101.54	
.976	3	3 0 3	104.20	
.963	1	3 2 1	106.20	
.950	2	4 0 2	108.30	
.944	1	4 1 0	109.32	
.924	1	1 1 4	112.96	
.919	5	4 1 1	113.94	
.904	2	2 0 4	116.96	
.890	1	3 1 3	119.92	
.853	2	4 1 2	129.10	
.819	1	3 0 4	140.38	
.818	1	4 2 0	140.74	
.815	2	3 3 1	141.80	
.795	1	3 2 3	151.52	

Cobalt samarium, Co₅Sm - continued

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
4.328	2	1	0	0	20.51
3.978	7	0	0	1	22.33
2.929	56	1	0	1	30.50
2.498	35	1	1	0	35.91
2.164	36	2	0	0	41.71
2.116	100	1	1	1	42.70
1.989	25	0	0	2	45.57
1.901	5	2	0	1	47.81
1.556	12	1	1	2	59.34
1.513	11	2	1	1	61.22
1.464	15	2	0	2	63.48
1.443	4	3	0	0	64.55
1.356	16	3	0	1	69.23
1.268	3	1	0	3	74.83
1.249	10	2	2	0	76.14
1.192	1	2	2	1	80.53
1.171	8	1	1	3	82.24
1.168	3	3	0	2	82.55
1.149	3	3	1	1	84.10
1.082	2	4	0	0	90.80
1.058	10	2	2	2	93.46
1.030	2	2	1	3	96.81
.994	1	0	0	4	101.53
.976	4	3	0	3	104.20
.963	2	3	2	1	106.20
.950	3	4	0	2	108.29
.944	2	4	1	0	109.31
.924	2	1	1	4	112.95
.919	7	4	1	1	113.94
.904	2	2	0	4	116.96
.890	2	3	1	3	119.92
.853	3	4	1	2	129.10
.846	1	5	0	1	131.24
.833	1	3	3	0	135.31
.819	2	3	0	4	140.37
.818	3	4	2	0	140.74
.815	4	3	3	1	141.81
.795	2	3	2	3	151.51
.782	2	1	0	5	159.75

Cobalt tin, Co_3Sn_2

Structure

Hexagonal, $P6_3/mmc$ (194), $Z = 1$, isostructural with Ni_3Sn_2 , type $B8_2$, from powder data [Rajeswari and Manohar, 1970].

Lattice constants: [ibid.]

$$\begin{aligned} a &= 4.109 \text{ \AA} \\ c &= 5.180 \end{aligned}$$

Density

(calculated) 9.080 g/cm^3

Thermal parameters

Isotropic: overall $B = 1.0$

Polymorphism

This phase was annealed above 550°C , and quenched to room temperature. A low-temperature, more ordered modification also exists [Rajeswari and Manohar, op. cit.].

Scattering factors

Co^0 , Sn^0 [Cromer and Mann, 1968].

Scale factors (integrated intensities)

$$\gamma = 0.383 \times 10^{-3}$$

$$I/I_c \text{ (calculated)} = 6.62$$

References

- Cromer, D. T. and Mann, J. B. (1968). Acta Crystallogr. A24, 321.
Rajeswari, H. and Manohar, H. (1970). Indian J. Pure Appl. Phys. 8, 363.

Calculated Pattern (Peak heights)				
$d(\text{\AA})$	I	hkl	$2\theta(^{\circ})$	$\lambda = 1.540598 \text{ \AA}$
3.558	2	1 0 0	25.02	
2.933	82	1 0 1	30.46	
2.590	12	0 0 2	34.60	
2.094	88	1 0 2	43.16	
2.055	100	1 1 0	44.04	
1.683	12	2 0 1	54.48	
1.610	14	1 1 2	57.19	
1.553	9	1 0 3	59.46	
1.467	21	2 0 2	63.38	
1.302	9	2 1 1	72.56	
1.295	6	0 0 4	73.00	
1.239	4	2 0 3	76.89	
1.194	18	2 1 2	80.38	
1.186	11	3 0 0	81.00	
1.096	15	1 1 4	89.36	
1.078	3	3 0 2	91.16	
1.061	4	2 1 3	93.10	
1.027	6	2 2 0	97.16	
.995	2	1 0 5	101.50	
.970	3	3 1 1	105.22	
.955	2	2 2 2	107.54	
.922	7	3 1 2	113.28	
.895	1	2 0 5	118.72	
.877	2	4 0 1	122.94	
.875	9	3 0 4	123.44	
.857	3	3 1 3	128.06	
.841	3	4 0 2	132.56	
.839	4	1 0 6	133.28	
.821	2	2 1 5	139.62	
.806	3	3 2 1	145.58	
.805	8	2 2 4	146.34	
.796	2	1 1 6	150.84	
.791	1	4 0 3	153.84	

Cobalt tin, Co₃Sn₂ - continued

Calculated Pattern (Integrated)					
d(Å)	I	hkl			2θ(°) λ = 1.540598Å
3.558	2	1	0	0	25.00
2.933	76	1	0	1	30.45
2.590	12	0	0	2	34.60
2.094	89	1	0	2	43.17
2.055	100	1	1	0	44.04
1.683	13	2	0	1	54.49
1.610	15	1	1	2	57.18
1.553	10	1	0	3	59.45
1.467	24	2	0	2	63.37
1.302	10	2	1	1	72.56
1.295	6	0	0	4	73.00
1.239	4	2	0	3	76.87
1.194	20	2	1	2	80.38
1.186	12	3	0	0	80.99
1.096	18	1	1	4	89.36
1.078	3	3	0	2	91.17
1.061	5	2	1	3	93.10
1.027	7	2	2	0	97.16
.995	2	1	0	5	101.50
.970	4	3	1	1	105.22
.955	3	2	2	2	107.55
.922	10	3	1	2	113.28
.895	2	2	0	5	118.72
.877	2	4	0	1	122.94
.875	13	3	0	4	123.44
.857	4	3	1	3	128.05
.841	5	4	0	2	132.56
.839	6	1	0	6	133.31
.821	5	2	1	5	139.61
.806	6	3	2	1	145.57
.805	18	2	2	4	146.33
.796	4	1	1	6	150.85
.791	4	4	0	3	153.83

CUMULATIVE INORGANIC INDEX

	Vol. or Sec.	Page		Vol. or Sec.	Page
Aluminum, Al	1	11	Ammonium copper bromide hydrate, (NH ₄) ₂ CuBr ₄ ·2H ₂ O	10m	6
Aluminum antimony, AlSb	4	72	Ammonium copper chloride, NH ₄ CuCl ₃	7m	7
Aluminum bismuth oxide, Al ₄ Bi ₂ O ₉ ..	11m	5	Ammonium copper chloride hydrate, (NH ₄) ₂ CuCl ₄ ·2H ₂ O	12m	6
Aluminum chloride, AlCl ₃	9m	61	Ammonium copper fluoride, NH ₄ CuF ₃ ..	11m	8
Aluminum chloride hydrate (chloraluminite), AlCl ₃ ·6H ₂ O	7	3	Ammonium gallium sulfate hydrate, NH ₄ Ga(SO ₄) ₂ ·12H ₂ O	6	9
Aluminum fluorine hydroxide silicate, topaz, Al ₂ (F,OH) ₂ SiO ₄	1m	4	Ammonium germanium fluoride, (NH ₄) ₂ GeF ₆	6	8
Aluminum nitride, AlN	12m	5	Ammonium hydrogen carbonate (teschemacherite), (NH ₄)HCO ₃	9	5
Aluminum nitrate hydrate, Al(NO ₃) ₃ ·9H ₂ O	11m	6	Ammonium hydrogen phosphate, NH ₄ H ₂ PO ₄	4	64
Aluminum oxide (corundum), α-Al ₂ O ₃ ..	9	3	Ammonium iodate, NH ₄ IO ₃	10m	7
Aluminum oxide hydrate (boehmite), α-Al ₂ O ₃ ·H ₂ O	3	38	Ammonium iodide, NH ₄ I	4	56
Aluminum oxide hydrate, diaspore, β-Al ₂ O ₃ ·H ₂ O	3	41	Ammonium iridium chloride, (NH ₄) ₂ IrCl ₆	8	6
Aluminum phosphate, Al(PO ₃) ₃	2m	3	Ammonium iron fluoride, (NH ₄) ₃ FeF ₆ ..	9m	9
Aluminum phosphate (berlinite), AlPO ₄ (trigonal)	10	3	Ammonium iron sulfate, NH ₄ Fe(SO ₄) ₂ ..	10m	8
Aluminum phosphate, AlPO ₄ (orthorhombic)	10	4	Ammonium iron sulfate hydrate, NH ₄ Fe(SO ₄) ₂ ·12H ₂ O	6	10
Aluminum silicate (mullite), Al ₆ Si ₂ O ₁₃	3m	3	Ammonium lead chloride, (NH ₄) ₂ PbCl ₆ ..	11m	10
Aluminum tungsten oxide, Al ₂ (WO ₄) ₃ ..	11m	7	Ammonium magnesium aluminum fluoride, NH ₄ MgAlF ₆	10m	9
Ammonium aluminum fluoride, (NH ₄) ₃ AlF ₆	9m	5	Ammonium magnesium chromium oxide hydrate, (NH ₄) ₂ Mg(CrO ₄) ₂ ·6H ₂ O	8m	10
Ammonium aluminum selenate hydrate, NH ₄ Al(SeO ₄) ₂ ·12H ₂ O	9m	6	Ammonium manganese chloride hydrate, (NH ₄) ₂ MnCl ₄ ·2H ₂ O	11m	11
Ammonium aluminum sulfate, NH ₄ Al(SO ₄) ₂	10m	5	Ammonium manganese(II) fluoride, NH ₄ MnF ₃	5m	8
Ammonium aluminum sulfate hydrate (tschermigite), NH ₄ Al(SO ₄) ₂ ·12H ₂ O ..	6	3	Ammonium manganese sulfate, (NH ₄) ₂ Mn ₂ (SO ₄) ₃	7m	8
Ammonium azide, NH ₄ N ₃	9	4	Ammonium manganese sulfate hydrate, (NH ₄) ₂ Mn(SO ₄) ₂ ·6H ₂ O	8m	12
Ammonium beryllium fluoride, (NH ₄) ₂ BeF ₄	3m	5	Ammonium mercury chloride, NH ₄ HgCl ₃ ..	8m	14
Ammonium boron fluoride, NH ₄ BF ₄ ...	3m	6	Ammonium molybdenum oxide phosphate hydrate, (NH ₄) ₃ (MoO ₃) ₁₂ PO ₄ ·4H ₂ O ...	8	10
Ammonium bromide, NH ₄ Br	2	49	Ammonium nickel(II) chloride, NH ₄ NiCl ₃	6m	6
Ammonium cadmium chloride, NH ₄ CdCl ₃ ..	5m	6	Ammonium nickel chromium oxide hydrate, (NH ₄) ₂ Ni(CrO ₄) ₂ ·6H ₂ O	8m	16
Ammonium cadmium sulfate, (NH ₄) ₂ Cd ₂ (SO ₄) ₃	7m	5	Ammonium nitrate (nitrammite), NH ₄ NO ₃	7	4
Ammonium cadmium sulfate hydrate, (NH ₄) ₂ Cd(SO ₄) ₂ ·6H ₂ O	8m	5	Ammonium osmium bromide, (NH ₄) ₂ OsBr ₆ ..	3	71
Ammonium calcium sulfate, (NH ₄) ₂ Ca ₂ (SO ₄) ₃	8m	7	Ammonium osmium chloride, (NH ₄) ₂ OsCl ₆	1m	6
Ammonium chlorate, NH ₄ ClO ₄ (orthorhombic)	7	6	Ammonium palladium chloride, (NH ₄) ₂ PdCl ₄	6	6
Ammonium chloride (sal-ammoniac), NH ₄ Cl	1	59	Ammonium palladium chloride, (NH ₄) ₂ PdCl ₆	8	7
Ammonium chromium sulfate hydrate, NH ₄ Cr(SO ₄) ₂ ·12H ₂ O	6	7	Ammonium platinum bromide, (NH ₄) ₂ PtBr ₆	9	6
Ammonium cobalt (II) chloride, NH ₄ CoCl ₃	6m	5	Ammonium platinum chloride, (NH ₄) ₂ PtCl ₆	5	3
Ammonium cobalt fluoride, NH ₄ CoF ₃ ..	8m	9	Ammonium rhenium oxide, NH ₄ ReO ₄	9	7
			Ammonium selenium bromide, (NH ₄) ₂ SeBr ₆	8	4
			Ammonium silicon fluoride (cryptohalite), (NH ₄) ₂ SiF ₆	5	5
			Ammonium sulfate (mascagnite), (NH ₄) ₂ SO ₄	9	8
			Ammonium tellurium bromide, (NH ₄) ₂ TeBr ₆	8	5

Further work on this program is in progress, and it is anticipated that additional sections will be issued. Therefore, the accumulative index here is not necessarily the concluding index for the project.

m - Monograph 25.

A mineral name in () indicates a synthetic sample.

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Ammonium tellurium chloride, (NH ₄) ₂ TeCl ₆	8	8	Barium lead nitrate, Ba ₃₃ Pb ₆₇ (NO ₃) ₂	12m	40
Ammonium tin chloride, (NH ₄) ₂ SnCl ₆	5	4	Barium lead nitrate, Ba ₆₇ Pb ₃₃ (NO ₃) ₂	12m	40
Ammonium vanadium oxide, NH ₄ VO ₃ ...	8	9	Barium molybdenum oxide, BaMoO ₄	7	7
Ammonium zinc fluoride, NH ₄ ZnF ₃ ...	8m	18	Barium molybdenum oxide, Ba ₂ MoO ₅ ...	12m	10
Ammonium zirconium fluoride, (NH ₄) ₃ ZrF ₇	6	14	Barium nitrate (nitrobarite), Ba(NO ₃) ₂	11m	14
Antimony, Sb	3	14	Barium oxide, BaO	9m	63
Antimony(III) fluoride, SbF ₃	2m	4	Barium oxide, BaO ₂	6	18
Antimony(III) iodide, SbI ₃	6	16	Barium phosphate, Ba ₃ (PO ₄) ₂	12m	12
Antimony(III) oxide (senarmontite), Sb ₂ O ₃ (cubic)	3	31	Barium selenide, BaSe	5m	61
Antimony(III) oxide, valentinite, Sb ₂ O ₃ (orthorhombic)	10	6	Barium silicate, β-BaSiO ₃	13m	8
Antimony(IV) oxide (cervantite), Sb ₂ O ₄	10	8	Barium silicate (sanbornite), β-BaSi ₂ O ₅	13m	10
Antimony(V) oxide, Sb ₂ O ₅	10	10	Barium silicate, Ba ₂ SiO ₄	13m	12
Antimony selenide, Sb ₂ Se ₃	3m	7	Barium silicate, Ba ₂ Si ₃ O ₈	13m	13
Antimony(III) sulfide (stibnite), Sb ₂ S ₃	5	6	Barium silicate, Ba ₃ SiO ₅	13m	15
Antimony telluride, Sb ₂ Te ₃	3m	8	Barium silicate, Ba ₃ Si ₅ O ₁₃	13m	17
Arsenic, As	3	6	Barium silicon fluoride, BaSiF ₆	4m	7
Arsenic(III) iodide, AsI ₃	13m	7	Barium strontium nitrate, Ba ₂₅ Sr ₇₅ (NO ₃) ₂	12m	42
Arsenic oxide (arsenolite), As ₂ O ₃ (cubic)	1	51	Barium strontium nitrate, Ba ₅₀ Sr ₅₀ (NO ₃) ₂	12m	42
Arsenic oxide, claudetite, As ₂ O ₃ (monoclinic)	3m	9	Barium strontium nitrate, Ba ₇₅ Sr ₂₅ (NO ₃) ₂	12m	42
Barium, Ba	4	7	Barium sulfate (baryte), BaSO ₄	10m	12
Barium aluminum oxide, BaAl ₂ O ₄	5m	11	Barium sulfide, BaS	7	8
Barium aluminum oxide, Ba ₃ Al ₂ O ₆ ...	12m	7	Barium tin oxide, BaSnO ₃	3m	11
Barium arsenate, Ba ₃ (AsO ₄) ₂	2m	6	Barium titanium oxide, BaTiO ₃	3	45
Barium borate, BaB ₄ O ₇	4m	6	Barium titanium silicate (fresnoite), Ba ₂ TiSi ₂ O ₈	9m	14
Barium borate, high form, BaB ₂ O ₄ ..	4m	4	Barium tungsten oxide, BaWO ₄	7	9
Barium borate, BaB ₈ O ₁₃	7m	10	Barium tungsten oxide, Ba ₂ WO ₅	12m	14
Barium bromate hydrate, Ba(BrO ₃) ₂ ·H ₂ O	8m	19	Barium zirconium oxide, BaZrO ₃	5	8
Barium bromide, BaBr ₂	10m	63	Beryllium, alpha, Be	9m	64
Barium bromide fluoride, BaBrF	10m	10	Beryllium aluminum oxide (chrysoberyl), BeAl ₂ O ₄	9	10
Barium bromide hydrate, BaBr ₂ ·H ₂ O	3m	10	Beryllium aluminum silicate, beryl, Be ₃ Al ₂ (SiO ₃) ₆	9	13
Barium calcium nitrate, Ba ₂₅ Ca ₇₅ (NO ₃) ₂	12m	38	Beryllium calcium oxide, Be ₁₇ Ca ₁₂ O ₂₉	7m	89
Barium calcium nitrate, Ba ₅₀ Ca ₅₀ (NO ₃) ₂	12m	38	Beryllium chromium oxide, BeCr ₂ O ₄ ..	10	12
Barium calcium nitrate, Ba ₇₅ Ca ₂₅ (NO ₃) ₂	12m	38	Beryllium cobalt, BeCo	5m	62
Barium calcium tungsten oxide, Ba ₂ CaWO ₆	9m	10	Beryllium germanium oxide, Be ₂ GeO ₄	10	13
Barium carbonate (witherite), BaCO ₃ (orthorhombic)	2	54	Beryllium lanthanum oxide, Be ₂ La ₂ O ₅	9m	65
Barium carbonate, BaCO ₃ (cubic) at 1075 °C	10	11	Beryllium niobium, Be ₂ Nb	7m	92
Barium chlorate hydrate, Ba(ClO ₄) ₂ ·3H ₂ O	2m	7	Beryllium oxide (bromellite), BeO ..	1	36
Barium chlorate hydrate, Ba(ClO ₃) ₂ ·H ₂ O	8m	21	Beryllium palladium, BePd	5m	62
Barium chloride, BaCl ₂ , (cubic) ...	9m	13	Beryllium silicate, phenacite, BeSi ₂ O ₄	8	11
Barium chloride, BaCl ₂ , (orthorhombic)	9m	11	Bismuth, Bi	3	20
Barium chloride fluoride, BaClF ...	10m	11	Bismuth fluoride, BiF ₃	1m	7
Barium chloride hydrate, BaCl ₂ ·2H ₂ O	12m	9	Bismuth(III) iodide, BiI ₃	6	20
Barium fluoride, BaF ₂	1	70	Bismuth oxide (bismite), α-Bi ₂ O ₃ ...	3m	16
Barium hydroxide phosphate, Ba ₅ (OH)(PO ₄) ₃	11m	12	Bismuth oxide bromide, BiOBr	8	14
Barium iodide, BaI ₂	10m	66	Bismuth oxide chloride (bismoclite), BiOCl	4	54
Barium lead chloride, BaPbCl ₄	11m	13	Bismuth oxide iodide, BiOI	9	16
			Bismuth phosphate, BiPO ₄ (monoclinic)	3m	11
			Bismuth phosphate, BiPO ₄ (trigonal)	3m	13
			Bismuth sulfide (bismuthinite), Bi ₂ S ₃	5m	13
			Bismuth telluride, BiTe	4m	50
			Bismuth telluride (tellurobis- muthite), Bi ₂ Te ₃	3m	16

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Bismuth vanadium oxide, low form, BiVO ₄ (tetragonal)	3m	14	Calcium fluoride phosphate (fluorapatite), Ca ₅ F(PO ₄) ₃	3m	22
Bismuth vanadium oxide, high form, BiVO ₄ (monoclinic)	3m	14	Calcium gallium germanium oxide, Ca ₃ Ga ₂ (GeO ₄) ₃	10	18
Boron oxide, B ₂ O ₃ , phase 1	10m	70	Calcium hydrogen phosphate hydrate, Ca ₈ H ₂ (PO ₄) ₆ ·5H ₂ O	13m	21
Cadmium, Cd	3	10	Calcium hydroxide (portlandite), Ca(OH) ₂	1	58
Cadmium ammine chloride, Cd(NH ₃) ₂ Cl ₂	10m	14	Calcium iron germanium oxide, Ca ₃ Fe ₂ (GeO ₄) ₃	10	19
Cadmium bromide, CdBr ₂	9	17	Calcium iron silicate (andradite), Ca ₃ Fe ₂ Si ₃ O ₁₂	9	22
Cadmium bromide chloride, CdBrCl ...	11m	15	Calcium iron silicate hydroxide, jul- goldite, Ca ₂ Fe ₃ Si ₃ O ₁₀ (OH,O) ₂ (OH) ₂	10m	72
Cadmium carbonate (otavite), CdCO ₃ ..	7	11	Calcium lead nitrate, Ca ₃₃ Pb ₆₇ (NO ₃) ₂	12m	44
Cadmium chlorate hydrate, Cd(ClO ₄) ₂ ·6H ₂ O	3m	19	Calcium lead nitrate, Ca ₆₇ Pb ₃₃ (NO ₃) ₂	12m	44
Cadmium chloride, CdCl ₂	9	18	Calcium magnesium silicate (diopside), CaMg(SiO ₃) ₂	5m	17
Cadmium chromium oxide, CdCr ₂ O ₄	5m	16	Calcium molybdenum oxide (powellite), CaMoO ₄	6	22
Cadmium cyanide, Cd(CN) ₂	2m	8	Calcium nitrate, Ca(NO ₃) ₂	7	14
Cadmium fluoride, CdF ₂	10m	15	Calcium oxide, CaO	1	43
Cadmium iron oxide, CdFe ₂ O ₄	9m	16	Calcium oxide phosphate, Ca ₄ O(PO ₄) ₂	12m	17
Cadmium manganese oxide, CdMn ₂ O ₄ ...	10m	16	Calcium phosphate, β-Ca ₂ P ₂ O ₇	7m	95
Cadmium molybdenum oxide, CdMoO ₄ ...	6	21	Calcium platinum oxide, Ca ₄ PtO ₆	10m	18
Cadmium nitrate hydrate, Cd(NO ₃) ₂ ·4H ₂ O	7m	93	Calcium selenide, CaSe	5m	64
Cadmium oxide, CdO	2	27	Calcium strontium nitrate, Ca ₃₃ Sr ₆₇ (NO ₃) ₂	12m	46
Cadmium oxide, CdO (ref. standard)	8m	2	Calcium strontium nitrate, Ca ₆₇ Sr ₃₃ (NO ₃) ₂	12m	46
Cadmium selenide, (cadmoselite), CdSe (hexagonal)	7	12	Calcium sulfate (anhydrite), CaSO ₄ ..	4	65
Cadmium silicate, Cd ₂ SiO ₄	13m	19	Calcium sulfide (oldhamite), CaS ...	7	15
Cadmium silicate, Cd ₃ SiO ₅	13m	20	Calcium telluride, CaTe	4m	50
Cadmium sulfate, CdSO ₄	3m	20	Calcium titanium oxide (perovskite), CaTiO ₃	9m	17
Cadmium sulfate hydrate, 3CdSO ₄ ·8H ₂ O	6m	8	Calcium tungsten oxide, Ca ₃ WO ₆	9m	19
Cadmium sulfate hydrate, CdSO ₄ ·H ₂ O ..	6m	10	Calcium tungsten oxide, scheelite, CaWO ₄	6	23
Cadmium sulfide (greenockite), CdS	4	15	Carbon, diamond, C	2	5
Cadmium telluride, CdTe	3m	21	Cerium antimony, CeSb	4m	40
Cadmium tungsten oxide, CdWO ₄	2m	8	Cerium arsenate, CeAsO ₄	4m	8
Calcium, Ca	9m	68	Cerium arsenide, CeAs	4m	51
Calcium aluminum germanium oxide, Ca ₃ Al ₂ (GeO ₄) ₃	10	15	Cerium bismuth, CeBi	4m	46
Calcium aluminum hydroxide, Ca ₃ Al ₂ (OH) ₁₂	11m	16	Cerium cadmium, CeCd	5m	63
Calcium aluminum oxide, Ca ₃ Al ₂ O ₆ ...	5	10	Cerium(III) chloride, CeCl ₃	1m	8
Calcium aluminum oxide, (mayenite), Ca ₁₂ Al ₁₄ O ₃₃	9	20	Cerium cobalt, CeCo ₂	13m	50
Calcium aluminum sulfate hydrate (ettringite), Ca ₆ Al ₂ S ₃ O ₁₈ ·31H ₂ O ...	8	3	Cerium cobalt, Ce ₂₄ Co ₁₁	13m	51
Calcium bromide, CaBr ₂	11m	70	Cerium copper, CeCu ₆	7m	99
Calcium bromide hydrate, CaBr ₂ ·6H ₂ O ..	8	15	Cerium(III) fluoride, CeF ₃	8	17
Calcium carbonate (aragonite), CaCO ₃ (orthorhombic)	3	53	Cerium gallium, CeGa ₂	13m	54
Calcium carbonate (calcite), CaCO ₃ (hexagonal)	2	51	Cerium magnesium, CeMg ₃	13m	56
Calcium chloride (hydrophilite), CaCl ₂	11m	18	Cerium nickel, CeNi ₂	13m	58
Calcium chloride fluoride, CaClF ...	10m	17	Cerium niobium titanium oxide (aeschnite), CeNbTiO ₆	3m	24
Calcium chloride hydrate, CaCl ₂ ·4H ₂ O	11m	73	Cerium nitride, CeN	4m	51
Calcium chloride hydrate (antarcticite), CaCl ₂ ·6H ₂ O	12m	16	Cerium(IV) oxide (cerianite), CeO ₂ ..	1	56
Calcium chromium germanium oxide, Ca ₃ Cr ₂ (GeO ₄) ₃	10	16	Cerium phosphide, CeP	4m	52
Calcium chromium oxide, CaCrO ₄	7	13	Cerium thallium, CeTl	13m	59
Calcium chromium silicate (uvarovite), Ca ₃ Cr ₂ (SiO ₄) ₃	10	17	Cerium thallium, CeTl ₃	13m	60
Calcium fluoride (fluorite), CaF ₂ ..	1	69	Cerium thallium, Ce ₃ Tl	13m	61
			Cerium(III) vanadium oxide, CeVO ₄ ..	1m	9
			Cerium zinc, CeZn	5m	65

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Cesium aluminum sulfate hydrate, CsAl(SO ₄) ₂ ·12H ₂ O	6	25	Cesium selenium bromide, Cs ₂ SeBr ₆ ..	8	20
Cesium antimony fluoride, CsSbF ₆ ...	4m	9	Cesium silicon fluoride, Cs ₂ SiF ₆ ...	5	19
Cesium beryllium fluoride, CsBeF ₃ ..	9m	69	Cesium strontium chloride, CsSrCl ₃	6m	13
Cesium boron fluoride, CsBF ₄	8	22	Cesium sulfate, Cs ₂ SO ₄	7	17
Cesium bromate, CsBrO ₃	8	18	Cesium tellurium bromide, Cs ₂ TeBr ₆	9	24
Cesium bromide, CsBr	3	49	Cesium tin chloride, Cs ₂ SnCl ₆	5	16
Cesium cadmium bromide, CsCdBr ₃			Cesium vanadium sulfate hydrate, CsV(SO ₄) ₂ ·12H ₂ O	1m	11
(hexagonal)	10m	20	Cesium zinc sulfate hydrate, Cs ₂ Zn(SO ₄) ₂ ·6H ₂ O	7m	25
Cesium cadmium chloride, CsCdCl ₃			Chromium, Cr	5	20
(hexagonal)	5m	19	Chromium chloride, CrCl ₂	11m	77
Cesium calcium chloride, CsCaCl ₃ ...	5m	21	Chromium fluoride, Cr ₂ F ₅	7m	108
Cesium calcium fluoride, CsCaF ₃	8m	25	Chromium fluoride, CrF ₂	10m	81
Cesium calcium sulfate, Cs ₂ Ca ₂ (SO ₄) ₃	7m	12	Chromium(III) fluoride hydrate, CrF ₃ ·3H ₂ O	5m	25
Cesium cerium chloride, Cs ₂ CeCl ₆ ...	7m	101	Chromium iridium 3:1, Cr ₃ Ir	6m	14
Cesium chlorate, CsClO ₃	8	20	Chromium(III) oxide, Cr ₂ O ₃	5	22
Cesium chlorate, CsClO ₄ , (orthorhombic)	1m	10	Chromium phosphate, α-CrPO ₄	2m	12
Cesium chloride, CsCl	2	44	Chromium phosphate, β-CrPO ₄	9	26
Cesium chromium oxide, Cs ₂ CrO ₄	3m	25	Chromium rhodium 3:1, Cr ₃ Rh	6m	15
Cesium chromium sulfate hydrate, CsCr(SO ₄) ₂ ·12H ₂ O	8	21	Chromium silicide, Cr ₃ Si	6	29
Cesium cobalt(II) chloride, CsCoCl ₃	6m	11	Cobalt, Co (cubic)	4m	10
Cesium cobalt chloride, Cs ₂ CoCl ₄ ...	11m	19	Cobalt aluminum oxide, CoAl ₂ O ₄	9	27
Cesium copper(II) chloride, CsCuCl ₃	5m	22	Cobalt ammine iodide, Co(NH ₃) ₆ I ₃ ...	10m	83
Cesium copper chloride, Cs ₂ CuCl ₄ ...	11m	20	Cobalt antimony oxide, CoSb ₂ O ₆	5m	26
Cesium copper sulfate hydrate, Cs ₂ Cu(SO ₄) ₂ ·6H ₂ O	7m	14	Cobalt arsenide, CoAs ₂	4m	10
Cesium fluoride, CsF	3m	26	Cobalt arsenide (skutterudite), CoAs ₃	10	21
Cesium gallium sulfate hydrate, CsGa(SO ₄) ₂ ·12H ₂ O	8	23	Cobalt borate, Co ₃ (BO ₃) ₂	12m	20
Cesium germanium fluoride, Cs ₂ GeF ₆	5	17	Cobalt bromide hydrate, CoBr ₂ ·6H ₂ O	12m	21
Cesium iodide, CsI	4	47	Cobalt(II) carbonate (sphero- cobaltite), CoCO ₃	10	24
Cesium iodine bromide, CsI ₂ Br	7m	103	Cobalt chlorate hydrate, Co(ClO ₄) ₂ ·6H ₂ O	3m	28
Cesium iodine chloride, CsICl ₂	3	50	Cobalt chloride hydrate, CoCl ₂ ·2H ₂ O	11m	22
Cesium iron sulfate hydrate, Cs ₂ Fe(SO ₄) ₂ ·6H ₂ O	7m	16	Cobalt chloride hydrate, CoCl ₂ ·6H ₂ O	11m	23
Cesium iron sulfate hydrate, CsFe(SO ₄) ₂ ·12H ₂ O	6	28	Cobalt chromium oxide, CoCr ₂ O ₄	9m	21
Cesium lead(II) chloride, CsPbCl ₃			Cobalt dysprosium, Co ₂ Dy.....	13m	63
(tetragonal)	5m	24	Cobalt erbium, Co ₂ Er	13m	64
Cesium lead fluoride, CsPbF ₃	8m	26	Cobalt erbium, Co ₇ Er ₂	13m	65
Cesium lithium cobalt cyanide, CsLiCo(CN) ₆	10m	79	Cobalt fluoride, CoF ₂	10m	85
Cesium lithium fluoride, CsLiF ₂	7m	105	Cobalt fluoride hydrate, CoF ₂ ·4H ₂ O	11m	24
Cesium magnesium chromium oxide, Cs ₂ Mg ₂ (CrO ₄) ₃	8m	27	Cobalt gadolinium, CoGd ₃	13m	68
Cesium magnesium chromium oxide hydrate, Cs ₂ Mg(CrO ₄) ₂ ·6H ₂ O	8m	29	Cobalt gadolinium, Co ₂ Gd.....	13m	71
Cesium magnesium sulfate hydrate, Cs ₂ Mg(SO ₄) ₂ ·6H ₂ O	7m	18	Cobalt gadolinium, Co ₇ Gd ₂	13m	72
Cesium manganese fluoride, CsMnF ₃ ..	10m	21	Cobalt gallium manganese, Co ₂ GaMn...	13m	75
Cesium manganese sulfate hydrate, Cs ₂ Mn(SO ₄) ₂ ·6H ₂ O	7m	20	Cobalt gallium oxide, CoGa ₂ O ₄	10	27
Cesium mercury chloride, CsHgCl ₃ ...	7m	22	Cobalt gallium tantalum, Co ₂ GaTa....	13m	76
Cesium nickel(II) chloride, CsNiCl ₃	6m	12	Cobalt gallium titanium, Co ₂ GaTi....	13m	77
Cesium nickel sulfate hydrate, Cs ₂ Ni(SO ₄) ₂ ·6H ₂ O	7m	23	Cobalt gallium vanadium, Co ₂ GaV.....	13m	78
Cesium nitrate, CsNO ₃	9	25	Cobalt germanium manganese, Co ₂ GeMn.	13m	79
Cesium osmium(IV) bromide, Cs ₂ OsBr ₆	2m	10	Cobalt germanium oxide, Co ₂ GeO ₄	10	27
Cesium osmium chloride, Cs ₂ OsCl ₆ ...	2m	11	Cobalt germanium titanium, Co ₂ GeTi..	13m	80
Cesium platinum bromide, Cs ₂ PtBr ₆ ..	8	19	Cobalt indium, CoIn ₃	13m	81
Cesium platinum chloride, Cs ₂ PtCl ₆	5	14	Cobalt iodide, CoI ₂	4m	52
Cesium platinum fluoride, Cs ₂ PtF ₆ ..	6	27	Cobalt iron arsenide (safflorite), CoFeAs ₄	10	28
			Cobalt iron oxide, CoFe ₂ O ₄	9m	22
			Cobalt lanthanum, CoLa ₃	13m	83
			Cobalt lutetium, Co ₂ Lu.....	13m	86
			Cobalt mercury thiocyanate, Co[Hg(CNS) ₄]	2m	13
			Cobalt neodymium, Co ₂ Nd.....	13m	87
			Cobalt nickel tin, Co ₇₅ Ni ₇₅ Sn ₇₅ ..	13m	88

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Cobalt nitrate hydrate, α -Co(NO ₃) ₂ ·6H ₂ O	12m	22	Erbium vanadium oxide, ErVO ₄	5m	29
Cobalt(II) oxide, CoO	9	28	Europium arsenate, EuAsO ₄	3m	32
Cobalt(II,III) oxide, Co ₃ O ₄	9	29	Europium(III) chloride, EuCl ₃	1m	13
Cobalt phosphate, Co(PO ₃) ₂	13m	23	Europium gallium oxide, Eu ₃ Ga ₅ O ₁₂	2m	17
Cobalt samarium, Co ₅ Sm	13m	90	Europium nitride, EuN	4m	56
Cobalt silicate, Co ₂ SiO ₄ (orthorhombic)	4m	11	Europium oxide, EuO	4m	56
Cobalt silicon fluoride hydrate, CoSiF ₆ ·6H ₂ O	3m	27	Europium oxychloride, EuOCl	1m	13
Cobalt sulfate, β -CoSO ₄	2m	14	Europium phosphate, EuPO ₄	11m	26
Cobalt tin, Co ₃ Sn ₂	13m	92	Europium(III) vanadium oxide, EuVO ₄	4m	16
Cobalt titanium oxide, CoTiO ₃	4m	13	Gadolinium antimony, GdSb	4m	42
Cobalt tungsten oxide, CoWO ₄	4m	13	Gadolinium arsenate, GdAsO ₄	4m	17
Copper, Cu	1	15	Gadolinium arsenide, GdAs	4m	57
Copper aluminum, Cu ₉ Al ₄	11m	79	Gadolinium chloride hydrate, GdCl ₃ ·6H ₂ O	7m	118
Copper ammine selenate, Cu(NH ₃) ₄ SeO ₄	10m	87	Gadolinium fluoride, GdF ₃	1m	14
Copper ammine sulfate hydrate, Cu(NH ₃) ₄ SO ₄ ·H ₂ O	10m	90	Gadolinium gallium oxide, Gd ₃ Ga ₅ O ₁₂	2m	18
Copper antimony oxide, CuSb ₂ O ₆	5m	27	Gadolinium indium, GdIn	5m	67
Copper(I) bromide, CuBr	4	36	Gadolinium nitride, GdN	4m	57
Copper cadmium, Cu ₅ Cd ₈	11m	81	Gadolinium oxide, Gd ₂ O ₃	1m	16
Copper(I) chloride (nantokite), CuCl	4	35	Gadolinium oxychloride, GdOCl	1m	17
Copper fluoride hydrate, CuF ₂ ·2H ₂ O	11m	25	Gadolinium silver, GdAg	6m	87
Copper hydrogen phosphite hydrate, CuHPO ₃ ·2H ₂ O	11m	83	Gadolinium titanium oxide, Gd ₂ TiO ₅	8m	32
Copper hydroxide carbonate, azurite, Cu ₃ (OH) ₂ (CO ₃) ₂	10	30	Gadolinium vanadium oxide, GdVO ₄ ...	5m	30
Copper hydroxide carbonate (malachite), Cu ₂ (OH) ₂ CO ₃	10	31	Gallium, Ga	2	9
Copper imidazole nitrate, Cu(C ₃ H ₄ N ₂) ₄ (NO ₃) ₂	13m	24	Gallium antimony, GaSb	6	30
Copper(I) iodide (marchite), CuI ...	4	38	Gallium arsenide, GaAs	3m	33
Copper(I) oxide (cuprite), Cu ₂ O ...	2	23	Gallium magnesium, Ga ₂ Mg	12m	48
Copper(II) oxide (tenorite), CuO ...	1	49	Gallium magnesium, Ga ₅ Mg ₂	12m	51
Copper phosphate, α -Cu ₂ P ₂ O ₇	7m	113	Gallium oxide, α -Ga ₂ O ₃	4	25
Copper sulfate (chalcocyanite), CuSO ₄	3m	29	Gallium phosphate (α -quartz type), GaPO ₄	8	27
Copper(II) sulfide (covellite), CuS	4	13	Gallium phosphate hydrate, GaPO ₄ ·2H ₂ O	8m	34
Copper uranium oxide, CuUO ₄	10m	93	Germanium, Ge	1	18
Dysprosium antimony, DySb	4m	41	Germanium iodide, GeI ₂	4m	58
Dysprosium arsenate, DyAsO ₄	3m	30	Germanium(IV) iodide, GeI ₄	5	25
Dysprosium arsenide, DyAs	4m	53	Germanium oxide, GeO ₂ (hexagonal) (low form)	1	51
Dysprosium bismuth, DyBi	4m	47	Germanium oxide, GeO ₂ (tetragonal) (high form)	8	28
Dysprosium gallium oxide, Dy ₃ Ga ₅ O ₁₂	2m	15	Gold, Au	1	33
Dysprosium gold, DyAu	5m	66	Gold antimony 1:2 (aurostibite), AuSb ₂	7	18
Dysprosium nitride, DyN	4m	53	Gold(I) cyanide, AuCN	10	33
Dysprosium oxide, Dy ₂ O ₃	9	30	Gold potassium cyanide, AuK(CN) ₂ ...	8m	36
Dysprosium silver, DyAg	5m	66	Gold tin 1:1, AuSn	7	19
Dysprosium telluride, DyTe	4m	54	Gold titanium 1:3, AuTi ₃	6m	17
Dysprosium vanadium oxide, DyVO ₄ ...	4m	15	Hafnium, Hf	3	18
Erbium antimony, ErSb	4m	41	Holmium arsenate, HoAsO ₄	3m	34
Erbium arsenate, ErAsO ₄	3m	31	Holmium bismuth, HoBi	4m	48
Erbium arsenide, ErAs	4m	54	Holmium fluoride, HoF ₃	10m	23
Erbium bismuth, ErBi	4m	47	Holmium gold, HoAu	5m	68
Erbium gallium oxide, Er ₃ Ga ₅ O ₁₂ ...	1m	12	Holmium nitride, HoN	4m	58
Erbium manganese oxide, ErMnO ₃	2m	16	Holmium oxide, Ho ₂ O ₃	9	32
Erbium nitride, ErN	4m	55	Holmium selenide, HoSe	4m	59
Erbium oxide, Er ₂ O ₃	8	25	Holmium silver, HoAg	5m	68
Erbium phosphate, ErPO ₄	9	31	Holmium vanadium oxide, HoVO ₄	4m	18
Erbium silver, ErAg	5m	67	Hydrogen amidosulfate, H ₂ NSO ₃ H	7	54
Erbium telluride, ErTe	4m	55	Hydrogen arsenate, H ₅ As ₃ O ₁₀	7m	84
			Hydrogen borate, β -HBO ₂	9m	71
			Hydrogen borate (metaborite), HBO ₂ (cubic)	4m	27
			Hydrogen iodate, HIO ₃	5	28

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Hydrogen iodate, HI_3O_8	8m	104	Lead oxide (litharge), PbO (red, tetragonal).....	2	30
Hydrogen phosphate hydrate, $\text{H}_3\text{PO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$	12m	56	Lead oxide (massicot), PbO (yellow, orthorhombic).....	2	32
Hydrogen tellurate, H_6TeO_6	12m	34	Lead(II,III) oxide (minium), Pb_3O_4 ..	8	32
Indium, In	3	12	Lead oxide sulfate, $\text{Pb}_5\text{O}_4\text{SO}_4$	10m	27
Indium antimony, InSb	4	73	Lead oxybromide, $\text{Pb}_3\text{O}_2\text{Br}_2$	5m	32
Indium arsenide, InAs	3m	35	Lead selenide (clausthalite), PbSe ..	5	38
Indium oxide, In_2O_3	5	26	Lead strontium nitrate, $\text{Pb}_{.33}\text{Sr}_{.67}(\text{NO}_3)_2$	12m	53
Indium phosphate, InPO_4	8	29	Lead strontium nitrate, $\text{Pb}_{.67}\text{Sr}_{.33}(\text{NO}_3)_2$	12m	53
Indium sulfide, In_2S_3	11m	30	Lead sulfate (anglesite), PbSO_4	3	67
Iodine, I_2	3	16	Lead sulfide (galena), PbS	2	18
Iridium, Ir	4	9	Lead tin oxide, Pb_2SnO_4	10m	29
Iridium oxide, IrO_2	4m	19	Lead titanium oxide, PbTiO_3	5	39
Iridium titanium 1:3, IrTi_3	6m	20	Lead tungsten oxide (stolzite), PbWO_4	5m	34
Iron, $\alpha\text{-Fe}$	4	3	Lead uranium oxide, Pb_3UO_6	8m	109
Iron arsenide, FeAs	1m	19	Lithium aluminum, Li_3Al_4	10m	98
Iron arsenide (loellingite), FeAs_2	10	34	Lithium aluminum fluoride, $\alpha\text{-Li}_3\text{AlF}_6$	8m	111
Iron bromide, FeBr_2	4m	59	Lithium arsenate, Li_3AsO_4	2m	19
Iron chloride hydrate, $\text{FeCl}_2 \cdot 2\text{H}_2\text{O}$..	11m	32	Lithium azide, LiN_3	8m	113
Iron fluoride hydrate, $\text{FeF}_2 \cdot 4\text{H}_2\text{O}$...	11m	90	Lithium barium fluoride, LiBaF_3	5m	35
Iron hydroxide sulfate hydrate, butlerite, $\text{Fe}(\text{OH})\text{SO}_4 \cdot 2\text{H}_2\text{O}$	10m	95	Lithium beryllium fluoride, Li_2BeF_4	7m	126
Iron iodide, FeI_2	4m	60	Lithium borate, $\text{Li}_2\text{B}_4\text{O}_7$	8m	114
Iron(II,III) oxide (magnetite), Fe_3O_4	5m	31	Lithium bromide, LiBr	4	30
Iron sulfate hydrate (melanterite), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	8m	38	Lithium carbonate, Li_2CO_3	8m	42
Iron sulfide (pyrite), FeS_2	5	29	Lithium chlorate hydrate, $\text{LiClO}_4 \cdot 3\text{H}_2\text{O}$	8	34
Lanthanum antimony, LaSb	4m	42	Lithium chloride, LiCl	1	62
Lanthanum arsenate, LaAsO_4	3m	36	Lithium fluoride, LiF	1	61
Lanthanum arsenide, LaAs	4m	60	Lithium gallium oxide, LiGaO_2	10m	31
Lanthanum bismuth, LaBi	4m	48	Lithium hydroxide hydrate, $\text{LiOH} \cdot \text{H}_2\text{O}$	11m	92
Lanthanum borate, LaBO_3	1m	20	Lithium iodate, LiIO_3 (hexagonal) ..	7	26
Lanthanum cadmium, LaCd	5m	63	Lithium iodate, LiIO_3 (tetragonal)	10m	33
Lanthanum chloride, LaCl_3	1m	20	Lithium molybdenum oxide, Li_2MoO_4 (trigonal)	1m	23
Lanthanum fluoride, LaF_3	7	21	Lithium niobium oxide, LiNbO_3	6m	22
Lanthanum niobium titanium oxide, LaNbTiO_6	3m	37	Lithium nitrate, LiNO_3	7	27
Lanthanum nitrate hydrate, $\text{La}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$	8m	40	Lithium oxide, Li_2O	1m	25
Lanthanum nitride, LaN	4m	61	Lithium phosphate hydrate, $\text{Li}_3\text{P}_3\text{O}_9 \cdot 3\text{H}_2\text{O}$	2m	20
Lanthanum oxide, La_2O_3	3	33	Lithium phosphate, low form (lithio- phosphate), Li_3PO_4	4m	21
Lanthanum oxychloride, LaOCl	7	22	Lithium phosphate, high form, Li_3PO_4	3m	39
Lanthanum phosphide, LaP	5m	69	Lithium rubidium fluoride, LiRbF_2 ..	7m	128
Lanthanum selenide, LaSe	4m	61	Lithium selenide, Li_2Se	10m	100
Lanthanum zinc, LaZn	5m	70	Lithium silver bromide, $\text{Li}_{.2}\text{Ag}_{.8}\text{Br}$	12m	55
Lead, Pb	1	34	Lithium silver bromide, $\text{Li}_{.4}\text{Ag}_{.6}\text{Br}$	12m	55
Lead borate, PbB_4O_7	4m	19	Lithium silver bromide, $\text{Li}_{.6}\text{Ag}_{.4}\text{Br}$	12m	55
Lead bromide, PbBr_2	2	47	Lithium silver bromide, $\text{Li}_{.8}\text{Ag}_{.2}\text{Br}$	12m	55
Lead bromide chloride, PbBrCl	11m	33	Lithium sodium aluminum fluoride, cryolithionite, $\text{Li}_3\text{Na}_3\text{Al}_2\text{F}_{12}$	9m	23
Lead bromide fluoride, PbBrF	10m	25	Lithium sodium sulfate, LiNaSO_4	6m	24
Lead chloride (cotunnite), PbCl_2 ...	12m	23	Lithium sulfate, Li_2SO_4	6m	26
Lead carbonate (cerussite), PbCO_3	2	56	Lithium sulfate hydrate, $\text{Li}_2\text{SO}_4 \cdot \text{H}_2\text{O}$	4m	22
Lead chloride fluoride (matlockite), PbClF	13m	25	Lithium sulfide, Li_2S	10m	101
Lead fluoride, $\alpha\text{-PbF}_2$ (orthorhombic)	5	31			
Lead fluoride, $\beta\text{-PbF}_2$ (cubic)	5	33			
Lead fluoride iodide, PbFI	10m	26			
Lead hydroxide phosphate, $\text{Pb}_5(\text{PO}_4)_3\text{OH}$	8	33			
Lead(II) iodide, PbI_2	5	34			
Lead molybdenum oxide (wulfenite), PbMoO_4	7	23			
Lead nitrate, $\text{Pb}(\text{NO}_3)_2$	5	36			

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Lithium telluride, Li_2Te	10m	102	Magnesium lanthanum, MgLa	5m	69
Lithium tungsten oxide, Li_2WO_4 (trigonal)	1m	25	Magnesium lanthanum nitrate hydrate, $\text{Mg}_3\text{La}_2(\text{NO}_3)_{12} \cdot 24\text{H}_2\text{O}$	1m	22
Lithium tungsten oxide hydrate, $\text{Li}_2\text{WO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$	2m	20	Magnesium manganese oxide, MgMn_2O_4	10m	35
Lithium uranium fluoride, LiUF_5	7m	131	Magnesium mercury, MgHg	6m	84
Lutetium arsenate, LuAsO_4	5m	36	Magnesium molybdenum oxide, MgMoO_4	7m	28
Lutetium gallium oxide, $\text{Lu}_3\text{Ga}_5\text{O}_{12}$	2m	22	Magnesium nickel oxide, MgNiO_2	10m	36
Lutetium manganese oxide, LuMnO_3 ...	2m	23	Magnesium oxide (periclase), MgO ...	1	37
Lutetium nitride, LuN	4m	62	Magnesium phosphate, $\text{Mg}(\text{PO}_3)_2$	13m	26
Lutetium oxide, Lu_2O_3	1m	27	Magnesium phosphate, $\alpha\text{-Mg}_2\text{P}_2\text{O}_7$	9m	73
Lutetium vanadium oxide, LuVO_4	5m	37	Magnesium selenide, MgSe	5m	70
Magnesium, Mg	1	10	Magnesium selenite hydrate, $\text{MgSeO}_3 \cdot 6\text{H}_2\text{O}$	8m	116
Magnesium aluminum oxide (spinel), MgAl_2O_4	9m	25	Magnesium silicate, enstatite, MgSiO_3	6	32
Magnesium aluminum silicate (pyrope), $\text{Mg}_3\text{Al}_2(\text{SiO}_4)_3$	4m	24	Magnesium silicate (forsterite), Mg_2SiO_4	1	83
Magnesium aluminum silicate (low cordierite), $\text{Mg}_2\text{Al}_4\text{Si}_5\text{O}_{18}$ (orthorhombic)	1m	28	Magnesium sulfate hydrate (epsomite), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	7	30
Magnesium aluminum silicate (indialite) $\text{Mg}_2\text{Al}_4\text{Si}_5\text{O}_{18}$ (hexagonal)	1m	29	Magnesium sulfide, MgS	7	31
Magnesium ammonium phosphate hydrate, (struvite), $\text{MgNH}_4\text{PO}_4 \cdot 6\text{H}_2\text{O}$	3m	41	Magnesium sulfite hydrate, $\text{MgSO}_3 \cdot 6\text{H}_2\text{O}$	9m	26
Magnesium borate, $\text{Mg}_2\text{B}_2\text{O}_5$ (triclinic)	4m	25	Magnesium tin, Mg_2Sn	5	41
Magnesium bromide, MgBr_2	4m	62	Magnesium tin oxide, Mg_2SnO_4	10m	37
Magnesium bromide hydrate, $\text{MgBr}_2 \cdot 6\text{H}_2\text{O}$	11m	35	Magnesium titanium oxide (geikielite), MgTiO_3	5	43
Magnesium carbonate (magnesite), MgCO_3	7	28	Magnesium titanium oxide, Mg_2TiO_4 ..	12m	25
Magnesium cerium MgCe	5m	65	Magnesium tungsten oxide, MgWO_4	13m	27
Magnesium cerium nitrate hydrate, $\text{Mg}_3\text{Ce}_2(\text{NO}_3)_{12} \cdot 24\text{H}_2\text{O}$	10	20	Manganese, $\alpha\text{-Mn}$	7m	142
Magnesium chlorate hydrate, $\text{Mg}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$	7m	30	Manganese aluminum oxide (galaxite), MnAl_2O_4	9	35
Magnesium chloride (chloro- magnesite), MgCl_2	11m	94	Manganese bromide, MnBr_2	4m	63
Magnesium chloride hydrate, $\text{MgCl}_2 \cdot 12\text{H}_2\text{O}$	7m	135	Manganese(II) carbonate (rhodochrosite), MnCO_3	7	32
Magnesium chloride hydrate (bischofite), $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$	11m	37	Manganese chloride (scacchite), MnCl_2	8m	43
Magnesium chromium oxide (magnesiochromite), MgCr_2O_4	9	34	Manganese chloride hydrate, $\text{MnCl}_2 \cdot 2\text{H}_2\text{O}$	11m	38
Magnesium fluoride (sellaitite), MgF_2	4	33	Manganese chloride hydrate, $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	9m	28
Magnesium fluoride silicate (humite), $\text{Mg}_7\text{F}_2(\text{SiO}_4)_3$	1m	30	Manganese cobalt oxide, MnCo_2O_4	9m	30
Magnesium fluoride silicate (norbergite), $\text{Mg}_3\text{F}_2\text{SiO}_4$	10	39	Manganese fluoride, MnF_2	10m	105
Magnesium gallium oxide, MgGa_2O_4 ...	10	36	Manganese iodide, MnI_2	4m	63
Magnesium germanium oxide, Mg_2GeO_4 (cubic)	10	37	Manganese iron oxide (jacobsite), MnFe_2O_4	9	36
Magnesium germanium oxide, Mg_2GeO_4 (orthorhombic)	10	38	Manganese(II) oxide (manganosite), MnO	5	45
Magnesium gold, MgAu	6m	83	Manganese oxide (hausmannite), Mn_3O_4	10m	38
Magnesium hydrogen phosphate hydrate, newberyite, $\text{MgHPO}_4 \cdot 3\text{H}_2\text{O}$..	7m	139	Manganese oxide (bixbyite), $\alpha\text{-Mn}_2\text{O}_3$	11m	95
Magnesium hydroxide (brucite), $\text{Mg}(\text{OH})_2$	6	30	Manganese oxide (pyrolusite), $\beta\text{-MnO}_2$	10m	39
Magnesium iron hydroxide carbonate hydrate, pyroaurite, $\text{Mg}_6\text{Fe}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O}$, phase II	10m	104	Manganese oxide hydroxide, groutite, $\alpha\text{-MnOOH}$	11m	97
Magnesium iron hydroxide carbonate hydrate, sjögrenite, $\text{Mg}_6\text{Fe}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O}$, phase I	10m	103	Manganese selenide, MnSe	10	41
			Manganese sulfide (alabandite), $\alpha\text{-MnS}$	4	11
			Manganese(II) tungsten oxide (huebnerite), MnWO_4	2m	24
			Manganese vanadium oxide, $\text{Mn}_2\text{V}_2\text{O}_7$..	9m	75
			Mercury amide chloride, HgNH_2Cl	10m	40
			Mercury ammine chloride, $\text{Hg}(\text{NH}_3)_2\text{Cl}_2$	11m	39
			Mercury bromate, $\text{Hg}(\text{BrO}_3)_2$	10m	107
			Mercury(II) bromide, HgBr_2	10m	110
			Mercury(I) bromide, Hg_2Br_2	7	33

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Mercury(I) chloride (calomel)			Nickel sulfate hydrate (retgersite),		
Hg ₂ Cl ₂	13m	30	NiSO ₄ ·6H ₂ O	7	36
Mercury(II) chloride, HgCl ₂	13m	29	Nickel sulfide, millerite, NiS	1m	37
Mercury chloride sulfide,			Nickel tungsten oxide, NiWO ₄	2m	27
α-Hg ₃ Cl ₂ S ₂	8m	118	Niobium gold 3:1, Nb ₃ Au	6m	16
Mercury(II) cyanide, Hg(CN) ₂	6	35	Niobium iridium 3:1, Nb ₃ Ir	6m	19
Mercury(II) fluoride, HgF ₂	2m	25	Niobium osmium 3:1, Nb ₃ Os	6m	30
Mercury(I) iodide, HgI	4	49	Niobium oxychloride, NbOCl ₃	7m	148
Mercury(II) iodide, HgI ₂ (tetragonal)	7m	32	Niobium platinum 3:1, Nb ₃ Pt	6m	31
Mercury(II) oxide (montroydite), HgO	9	39	Niobium silicide, NbSi ₂	8	39
Mercury(II) selenide (tiemannite),			Osmium, Os	4	8
HgSe	7	35	Osmium titanium, OsTi	6m	85
Mercury(II) sulfide (cinnabar),			Palladium, Pd	1	21
HgS (hexagonal)	4	17	Palladium hydride, PdH _{0.706}	5m	72
Mercury(II) sulfide (metacinnabar),			Palladium oxide, PdO	4	27
HgS (cubic)	4	21	Phosphorus bromide, PBr ₇	7m	150
Molybdenum, Mo	1	20	Phosphorus oxide (stable form I),		
Molybdenum arsenide, Mo ₂ As ₃	10m	115	P ₂ O ₅ (orthorhombic)	9m	86
Molybdenum osmium 3:1, Mo ₃ Os	6m	28	Phosphorus oxide (stable form II),		
Molybdenum oxide (molybdate), MoO ₃	3	30	P ₂ O ₅ (orthorhombic)	9m	88
Molybdenum sulfide (molybdenite),			Phosphorus oxide (metastable form),		
MoS ₂	5	47	P ₄ O ₁₀ (rhombohedral)	9m	91
Neodymium antimony, NdSb	4m	43	Platinum, Pt	1	31
Neodymium arsenate, NdAsO ₄	4m	28	Platinum titanium 1:3, PtTi ₃	6m	33
Neodymium arsenide, NdAs	4m	64	Plutonium arsenide, PuAs	4m	65
Neodymium bismuth, NdBi	4m	49	Plutonium phosphide, PuP	4m	65
Neodymium borate, NdBO ₃	1m	32	Plutonium telluride, PuTe	4m	66
Neodymium chloride, NdCl ₃	1m	33	Potassium aluminum sulfate,		
Neodymium fluoride, NdF ₃	8	36	KAl(SO ₄) ₂	9m	31
Neodymium gallium oxide, Nd ₃ Ga ₅ O ₁₂	1m	34	Potassium aluminum sulfate hydrate,		
Neodymium oxide, Nd ₂ O ₃	4	26	(alum), KAl(SO ₄) ₂ ·12H ₂ O	6	36
Neodymium oxychloride, NdOCl	8	37	Potassium barium nickel nitrite,		
Neodymium phosphate, NdPO ₄	11m	40	K ₂ BaNi(NO ₂) ₆	9m	32
Neodymium selenide, NdSe	5m	71	Potassium borohydride, KBH ₄	9	44
Neodymium silver, NdAg	5m	71	Potassium bromate, KBrO ₃	7	38
Neodymium vanadium oxide, NdVO ₄	4m	30	Potassium bromide, KBr	1	66
Neptunium nitride, NpN	4m	64	Potassium bromide chloride,		
Nickel, Ni	1	13	KBr _{0.5} Cl _{0.5}	8m	46
Nickel acetate hydrate,			Potassium bromide iodide,		
Ni(C ₂ H ₃ O ₂) ₂ ·4H ₂ O	13m	31	KBr _{.33} I _{.67}	11m	44
Nickel aluminum, NiAl	6m	82	Potassium bromide iodide,		
Nickel aluminum oxide, NiAl ₂ O ₄	9	42	KBr _{.67} I _{.33}	11m	45
Nickel arsenide 1:2 (rammelsbergite),			Potassium cadmium fluoride, KCdF ₃ ..	8m	47
NiAs ₂	10	42	Potassium cadmium sulfate,		
Nickel arsenic sulfide			K ₂ Cd ₂ (SO ₄) ₃	7m	34
(gersdorffite), NiAsS	1m	35	Potassium calcium carbonate		
Nickel bromide, NiBr ₂	10m	119	(fairchildite), K ₂ Ca(CO ₃) ₂	8m	48
Nickel(II) carbonate, NiCO ₃			Potassium calcium chloride, KCaCl ₃ ..	7m	36
(trigonal)	1m	36	Potassium calcium fluoride, KCaF ₃ ..	8m	49
Nickel chloride, NiCl ₂	9m	81	Potassium calcium magnesium sulfate,		
Nickel chloride hydrate,			K ₂ CaMg(SO ₄) ₃	7m	37
NiCl ₂ ·6H ₂ O	11m	42	Potassium calcium nickel nitrite,		
Nickel fluoride, NiF ₂	10m	121	K ₂ CaNi(NO ₂) ₆	9m	33
Nickel fluoride hydrate, NiF ₂ ·4H ₂ O	11m	43	Potassium calcium sulfate,		
Nickel gallium oxide, NiGa ₂ O ₄	10	45	K ₂ Ca ₂ (SO ₄) ₃	7m	39
Nickel germanium oxide, Ni ₂ GeO ₄	9	43	Potassium cerium fluoride, β-KCeF ₄	12m	59
Nickel iron oxide (trevorite),			Potassium chlorate, KClO ₃	3m	42
NiFe ₂ O ₄	10	44	Potassium chlorate, KClO ₄	6	43
Nickel nitrate hydrate,			Potassium chloride (sylvite), KCl ..	1	65
Ni(NO ₃) ₂ ·6H ₂ O	12m	26	Potassium chromium oxide, K ₃ CrO ₈ ...	3m	44
Nickel(II) oxide (bunsenite), NiO ..	1	47	Potassium chromium oxide sulfate,		
Nickel phosphide, Ni ₁₂ P ₅	9m	83	K ₂ (CrO ₄) _{.33} (SO ₄) _{.67}	12m	28
Nickel silicon fluoride hydrate,			Potassium chromium oxide sulfate,		
NiSiF ₆ ·6H ₂ O	8	38	K ₂ (CrO ₄) _{.67} (SO ₄) _{.33}	12m	27
Nickel sulfate, NiSO ₄	2m	26			

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Potassium chromium sulfate hydrate, KCr(SO ₄) ₂ ·12H ₂ O	6	39	Potassium rhenium chloride, K ₂ ReCl ₆	2m	28
Potassium cobalt(II) fluoride, KCoF ₃	6m	37	Potassium rhenium oxide, KReO ₄	8	41
Potassium cobalt fluoride, K ₂ CoF ₄ ..	11m	46	Potassium rubidium chloride, K _{0.5} Rb _{0.5} Cl	8m	76
Potassium cobalt nitrite, K ₃ Co(NO ₂) ₆	9	45	Potassium rubidium chromium oxide, KRbCrO ₄	12m	29
Potassium cobalt(II) sulfate, K ₂ Co ₂ (SO ₄) ₃	6m	35	Potassium ruthenium chloride, K ₂ RuCl ₆	10	46
Potassium copper chloride, KCuCl ₃ ..	7m	41	Potassium ruthenium oxide chloride hydrate, K ₄ Ru ₂ OCl ₁₀ ·H ₂ O	10	47
Potassium copper chloride hydrate (mitscherlichite), K ₂ CuCl ₄ ·2H ₂ O ...	9m	34	Potassium selenate, K ₂ SeO ₄	9m	41
Potassium copper(II) fluoride, KCuF ₃	6m	38	Potassium selenide, K ₂ Se	10m	126
Potassium cyanate, KCNO	7	39	Potassium selenium bromide, K ₂ SeBr ₆	8	41
Potassium cyanide, KCN	1	77	Potassium silicon fluoride (hieratite), K ₂ SiF ₆	5	50
Potassium fluoride, KF	1	64	Potassium silver cyanide, KAg(CN) ₂	8m	78
Potassium germanium fluoride, K ₂ GeF ₆	6	41	Potassium sodium aluminum fluoride (elpasolite), K ₂ NaAlF ₆	9m	43
Potassium hydrogen arsenate, KH ₂ AsO ₄	1m	38	Potassium sodium bromide, K ₂ Na ₈ Br	12m	62
Potassium hydrogen phosphate, KH ₂ PO ₄	3	69	Potassium sodium bromide, K ₄ Na ₆ Br	12m	62
Potassium hydroxide, KOH at 300 °C	4m	66	Potassium sodium bromide, K ₆ Na ₄ Br	12m	62
Potassium iodate, KIO ₄	7	41	Potassium sodium bromide, K ₈ Na ₂ Br	12m	62
Potassium iodide, KI	1	68	Potassium sodium chloride, K ₂ Na ₈ Cl	12m	63
Potassium iron cyanide, K ₃ Fe(CN) ₆ ..	9m	35	Potassium sodium chloride, K ₄ Na ₆ Cl	12m	63
Potassium iron(II) fluoride, KFeF ₃	6m	39	Potassium sodium chloride, K ₆ Na ₄ Cl	12m	63
Potassium iron fluoride, K ₃ FeF ₆	9m	37	Potassium sodium chloride, K ₈ Na ₂ Cl	12m	63
Potassium lead chloride, KPb ₂ Cl ₅	13m	33	Potassium sodium sulfate, K _{6.7} Na _{1.33} SO ₄	6m	48
Potassium lithium sulfate, KLiSO ₄	3m	43	Potassium sodium sulfate, KNaSO ₄ ...	6m	50
Potassium magnesium chloride hydrate (carnallite), KMgCl ₃ ·6H ₂ O.....	8m	50	Potassium sodium sulfate (aphthitalite), K ₃ Na(SO ₄) ₂	6m	52
Potassium magnesium chromium oxide, K ₂ Mg ₂ (CrO ₄) ₃	8m	52	Potassium sulfate, K ₂ S ₂ O ₇	9m	99
Potassium magnesium fluoride, KMgF ₃	6m	42	Potassium sulfate (arcanite), K ₂ SO ₄	3	62
Potassium magnesium fluoride, K ₂ MgF ₄	10m	42	Potassium sulfide, K ₂ S	10m	127
Potassium magnesium selenate hydrate, K ₂ Mg(SeO ₄) ₂ ·6H ₂ O	10m	43	Potassium telluride, K ₂ Te	10m	128
Potassium magnesium sulfate (langbeinite), K ₂ Mg ₂ (SO ₄) ₃	6m	40	Potassium thiocyanate, KCNS	8	44
Potassium magnesium sulfate hydrate (picromerite), K ₂ Mg(SO ₄) ₂ ·6H ₂ O	8m	54	Potassium tin chloride, K ₂ SnCl ₆	6	38
Potassium manganese(II) fluoride, KMnF ₃	6m	45	Potassium titanium fluoride, K ₂ TiF ₆	7	40
Potassium manganese oxide, KMnO ₄ ...	7	42	Potassium tungsten oxide, K ₂ WO ₄	11m	47
Potassium manganese(II) sulfate (manganolangbeinite), K ₂ Mn ₂ (SO ₄) ₃	6m	43	Potassium vanadium oxide, KV ₃ O ₈	8m	56
Potassium molybdenum oxide phosphate hydrate, K ₂ (MoO ₃) ₁₂ PO ₄ ·4H ₂ O	8	43	Potassium zinc bromide hydrate, KZnBr ₃ ·2H ₂ O	11m	104
Potassium nickel fluoride, KNiF ₃ ...	7m	42	Potassium zinc fluoride, KZnF ₃	5	51
Potassium nickel fluoride, K ₂ NiF ₄ ..	10m	45	Potassium zinc fluoride, K ₂ ZnF ₄	10m	46
Potassium nickel(II) sulfate, K ₂ Ni ₂ (SO ₄) ₃	6m	46	Potassium zinc iodide hydrate, KZnI ₃ ·2H ₂ O	11m	107
Potassium niobium fluoride, K ₂ NbF ₇	8m	120	Potassium zinc sulfate, K ₂ Zn ₂ (SO ₄) ₃	6m	54
Potassium nitrate (niter), KNO ₃	3	58	Potassium zinc sulfate hydrate, K ₂ Zn(SO ₄) ₂ ·6H ₂ O	7m	43
Potassium nitrite, KNO ₂	9m	38	Potassium zinc vanadium oxide hydrate, K ₂ Zn ₂ V ₁₀ O ₂₈ ·16H ₂ O	3m	45
Potassium nitroso ruthenium chloride, K ₂ (NO)RuCl ₅	2m	29	Potassium zirconium fluoride, K ₃ ZrF ₇	9	46
Potassium oxide, K ₂ O	10m	125	Praseodymium antimony, PrSb	4m	43
Potassium platinum bromide, K ₂ PtBr ₆	8	40	Praseodymium arsenate, PrAsO ₄	4m	32
Potassium platinum chloride, K ₂ PtCl ₆	13m	34	Praseodymium arsenide, PrAs	4m	67
Potassium platinum fluoride, K ₂ PtF ₆	6	42			

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Praseodymium bismuth, PrBi	4m	49	Rubidium nitrate, RbNO ₃ (trigonal)	5m	45
Praseodymium cadmium, PrCd	5m	64	Rubidium platinum chloride, Rb ₂ PtCl ₆	5	53
Praseodymium chloride, PrCl ₃	1m	39	Rubidium platinum fluoride, Rb ₂ PtF ₆	6	48
Praseodymium fluoride, PrF ₃	5	52	Rubidium selenate, Rb ₂ SeO ₄	9m	44
Praseodymium oxychloride, PrOCl	9	47	Rubidium silicon fluoride, Rb ₂ SiF ₆	6	49
Praseodymium sulfide, PrS	4m	67	Rubidium strontium chloride, RbSrCl ₃	7m	54
Praseodymium vanadium oxide, PrVO ₄	5m	40	Rubidium sulfate, Rb ₂ SO ₄	8	48
Praseodymium zinc, PrZn	5m	72	Rubidium tellurium bromide, Rb ₂ TeBr ₆	8	46
Rhenium, Re	2	13	Rubidium tellurium chloride, Rb ₂ TeCl ₆	8	48
Rhodium, Rh	3	9	Rubidium tin chloride, Rb ₂ SnCl ₆	6	46
Rubidium aluminum sulfate hydrate, RbAl(SO ₄) ₂ ·12H ₂ O	6	44	Rubidium zinc fluoride, RbZnF ₃	7m	57
Rubidium amide, RbNH ₂	5m	73	Rubidium zinc sulfate hydrate, Rb ₂ Zn(SO ₄) ₂ ·6H ₂ O	7m	55
Rubidium bromate, RbBrO ₃	8	45	Ruthenium, Ru	4	5
Rubidium bromide, RbBr	7	43	Ruthenium titanium, RuTi	6m	86
Rubidium cadmium chloride, high form, RbCdCl ₃ (tetragonal)	5m	43	Samarium arsenate, SmAsO ₄	4m	33
Rubidium cadmium chloride, low form, RbCdCl ₃ (orthorhombic)	5m	41	Samarium arsenide, SmAs	4m	68
Rubidium cadmium sulfate, Rb ₂ Cd ₂ (SO ₄) ₃	7m	45	Samarium chloride, SmCl ₃	1m	40
Rubidium calcium chloride, RbCaCl ₃	7m	47	Samarium fluoride, SmF ₃	1m	41
Rubidium calcium fluoride, RbCaF ₃	8m	57	Samarium gallium oxide, Sm ₃ Ga ₅ O ₁₂ ..	1m	42
Rubidium calcium sulfate, Rb ₂ Ca ₂ (SO ₄) ₃	7m	48	Samarium oxide, Sm ₂ O ₃ (cubic)	4m	34
Rubidium chlorate, RbClO ₃	8	47	Samarium oxychloride, SmOCl	1m	43
Rubidium chlorate, RbClO ₄	2m	30	Samarium silver, SmAg	5m	73
Rubidium chloride, RbCl	4	41	Samarium tin oxide, Sm ₂ Sn ₂ O ₇	8m	77
Rubidium chromium oxide, Rb ₂ CrO ₄ ...	3m	46	Samarium vanadium oxide, SmVO ₄	5m	47
Rubidium chromium sulfate hydrate, RbCr(SO ₄) ₂ ·12H ₂ O	6	47	Scandium antimony, ScSb	4m	44
Rubidium cobalt(II) chloride, RbCoCl ₃	6m	57	Scandium arsenate, ScAsO ₄	4m	35
Rubidium cobalt fluoride, RbCoF ₃ ...	8m	58	Scandium arsenide, ScAs	4m	68
Rubidium cobalt sulfate, Rb ₂ Co ₂ (SO ₄) ₃	8m	59	Scandium oxide, Sc ₂ O ₃	3	27
Rubidium copper chloride hydrate, Rb ₂ CuCl ₄ ·2H ₂ O	10m	47	Scandium phosphate, ScPO ₄	8	50
Rubidium copper sulfate hydrate, Rb ₂ Cu(SO ₄) ₂ ·6H ₂ O	8m	61	Scandium silicate (thortveitite), Sc ₂ Si ₂ O ₇	7m	58
Rubidium fluoride, RbF	8m	63	Selenium, Se	5	54
Rubidium iodate, RbIO ₄	2m	31	Selenium oxide (selenolite), SeO ₂ ..	7m	60
Rubidium iodide, RbI	4	43	Silicon, Si	13m	35
Rubidium iron sulfate hydrate, Rb ₂ Fe(SO ₄) ₂ ·6H ₂ O	8m	64	Silicon, Si (reference standard) ...	12m	2
Rubidium magnesium chromium oxide, Rb ₂ Mg ₂ (CrO ₄) ₃	8m	66	Silicon oxide (α or low cristobalite), SiO ₂ (tetragonal)	10	48
Rubidium magnesium chromium oxide hydrate, Rb ₂ Mg(CrO ₄) ₂ ·6H ₂ O	8m	68	Silicon oxide (α or low quartz), SiO ₂ (hexagonal)	3	24
Rubidium magnesium sulfate, Rb ₂ Mg ₂ (SO ₄) ₃	7m	50	Silicon oxide (β or high cristobalite), SiO ₂ (cubic)	1	42
Rubidium magnesium sulfate hydrate, Rb ₂ Mg(SO ₄) ₂ ·6H ₂ O	8m	70	Silver, Ag	1	23
Rubidium manganese(II) fluoride, RbMnF ₃	5m	44	Silver, Ag (reference standard)	8m	2
Rubidium manganese sulfate, Rb ₂ Mn ₂ (SO ₄) ₃	7m	52	Silver antimony sulfide, AgSbS ₂ (cubic)	5m	48
Rubidium nickel(II) chloride, RbNiCl ₃	6m	58	Silver antimony sulfide (miargyrite), AgSbS ₂ (monoclinic)	5m	49
Rubidium nickel sulfate, Rb ₂ Ni ₂ (SO ₄) ₃	8m	72	Silver antimony sulfide (pyrargyrite), Ag ₃ SbS ₃ (trigonal)	5m	51
Rubidium nickel sulfate hydrate, Rb ₂ Ni(SO ₄) ₂ ·6H ₂ O	8m	74	Silver antimony telluride, AgSbTe ₂	3m	47
			Silver arsenate, Ag ₃ AsO ₄	5	56
			Silver arsenic sulfide, xanthoconite, Ag ₃ AsS ₃	8m	126
			Silver bromate, AgBrO ₃	5	57
			Silver bromide (bromargyrite), AgBr	4	46
			Silver carbonate, Ag ₂ CO ₃	13m	36
			Silver chlorate, AgClO ₃	7	44
			Silver chloride (chlorargyrite), AgCl	4	44

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Silver chromium oxide, Ag_2CrO_4	12m	30	Sodium chromium oxide sulfate, $\text{Na}_4(\text{CrO}_4)(\text{SO}_4)$	11m	55
Silver cyanide, AgCN	9m	48	Sodium cobalt(II) sulfate hydrate, $\text{Na}_2\text{Co}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O}$	6m	61
Silver fluoride, Ag_2F	5m	53	Sodium cyanate, NaCNO	2m	33
Silver iodate, AgIO_4	9	49	Sodium cyanide, NaCN (cubic)	1	78
Silver iodide (iodargyrite), AgI (hexagonal)	8	51	Sodium cyanide, NaCN (orthorhombic) at 6 °C	1	79
Silver iodide, $\gamma\text{-AgI}$ (cubic)	9	48	Sodium fluoride (villiaumite), NaF	1	63
Silver manganese oxide, AgMnO_4	7m	155	Sodium hydrogen fluoride, NaHF_2	5	63
Silver molybdenum oxide, Ag_2MoO_4 ...	7	45	Sodium hydrogen phosphate, $\text{Na}_3\text{H}(\text{PO}_3)_4$	10m	130
Silver nitrate, AgNO_3	5	59	Sodium hydrogen silicate hydrate, $\text{Na}_2\text{H}_2\text{SiO}_4 \cdot 4\text{H}_2\text{O}$	7m	163
Silver nitrite, AgNO_2	5	60	Sodium hydrogen sulfate hydrate, $\text{NaHSO}_4 \cdot \text{H}_2\text{O}$	9m	52
Silver oxide, Ag_2O	1m	45	Sodium hydroxide, NaOH at 300 °C ...	4m	69
Silver(II) oxide nitrate, $\text{Ag}_7\text{O}_8\text{NO}_3$	4	61	Sodium iodate, NaIO_3	7	47
Silver phosphate, Ag_3PO_4	5	62	Sodium iodate, NaIO_4	7	48
Silver rhenium oxide, AgReO_4	8	53	Sodium iodide, NaI	4	31
Silver selenate, Ag_2SeO_4	2m	32	Sodium iron fluoride, Na_3FeF_6	9m	54
Silver sodium chloride, $\text{Ag}_{0.5}\text{Na}_{0.5}\text{Cl}$	8m	79	Sodium lanthanum fluoride silicate, $(\text{Na}_2\text{La}_8)\text{F}_2(\text{SiO}_4)_6$	7m	64
Silver sulfate, Ag_2SO_4	13m	37	Sodium lanthanum molybdenum oxide, $\text{NaLa}(\text{MoO}_4)_2$	10m	49
Silver sulfide (acanthite), Ag_2S	10	51	Sodium magnesium aluminum boron hydroxide silicate, dravite, $\text{NaMg}_3\text{Al}_6\text{B}_3(\text{OH})_4\text{Si}_6\text{O}_{27}$	3m	47
Sodium, Na	9m	105	Sodium magnesium carbonate (eitelite), $\text{Na}_2\text{Mg}(\text{CO}_3)_2$	11m	56
Sodium aluminum chloride silicate, sodalite, $\text{Na}_8\text{Al}_6\text{Cl}_{12}(\text{SiO}_4)_6$	7m	158	Sodium magnesium sulfate hydrate, bloedite, $\text{Na}_2\text{Mg}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O}$	6m	63
Sodium azide, $\alpha\text{-NaN}_3$, at -90 to -100 °C	8m	129	Sodium manganese(II) fluoride, NaMnF_3	6m	65
Sodium azide, $\beta\text{-NaN}_3$	8m	130	Sodium mercury(II) chloride hydrate, $\text{NaHgCl}_3 \cdot 2\text{H}_2\text{O}$	6m	66
Sodium beryllium calcium fluoride silicate, leucophanite, $\text{NaBeCaFSi}_2\text{O}_6$	8m	138	Sodium molybdenum oxide, Na_2MoO_4 ...	1m	46
Sodium borate, $\text{Na}_2\text{B}_8\text{O}_{13}$	7m	160	Sodium molybdenum oxide, $\text{Na}_2\text{Mo}_2\text{O}_7$..	9m	110
Sodium boron hydride, NaBH_4	9	51	Sodium neodymium fluoride silicate, $(\text{Na}_2\text{Nd}_8)\text{F}_2(\text{SiO}_4)_6$	7m	66
Sodium bromate, NaBrO_3	5	65	Sodium nickel(II) sulfate hydrate, $\text{Na}_2\text{Ni}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O}$	6m	68
Sodium bromide, NaBr	3	47	Sodium nitrate (soda-niter), NaNO_3	6	50
Sodium bromide chloride, $\text{NaBr}_{.33}\text{Cl}_{.67}$	11m	49	Sodium nitrite, NaNO_2	4	62
Sodium bromide chloride, $\text{NaBr}_{.67}\text{Cl}_{.33}$	11m	50	Sodium oxide, Na_2O	10m	134
Sodium calcium aluminum fluoride hydrate, thomsenolite, $\text{NaCaAlF}_6 \cdot \text{H}_2\text{O}$	8m	132	Sodium phosphate, $\text{Na}_3\text{P}_3\text{O}_9$	3m	49
Sodium calcium beryllium aluminum fluorosilicate, meliphanite, $(\text{Na}_{0.63}\text{Ca}_{1.37})\text{Be}(\text{Al}_{0.13}\text{Si}_{1.87})$ $(\text{O}_{6.25}\text{F}_{0.75})$	8m	135	Sodium phosphate hydrate, $\text{Na}_3\text{P}_3\text{O}_9 \cdot \text{H}_2\text{O}$	3m	50
Sodium calcium carbonate hydrate, pirssonite, $\text{Na}_2\text{Ca}(\text{CO}_3)_2 \cdot 2\text{H}_2\text{O}$	9m	106	Sodium phosphate hydrate, $\alpha\text{-Na}_4\text{P}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (monoclinic)	13m	39
Sodium calcium silicate, $\text{Na}_2\text{CaSiO}_4$	10m	48	Sodium phosphate hydrate, $\beta\text{-Na}_4\text{P}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$ (triclinic)	2m	35
Sodium calcium sulfate (glauberite), $\text{Na}_2\text{Ca}(\text{SO}_4)_2$	6m	59	Sodium phosphate hydrate, $\text{Na}_6\text{P}_6\text{O}_{18} \cdot 6\text{H}_2\text{O}$	5m	54
Sodium carbonate hydrate (thermo- natrite), $\text{Na}_2\text{CO}_3 \cdot \text{H}_2\text{O}$	8	54	Sodium praseodymium fluoride silicate, $(\text{Na}_2\text{Pr}_8)\text{F}_2(\text{SiO}_4)_6$	7m	68
Sodium carbonate sulfate, $\text{Na}_4\text{CO}_3\text{SO}_4$	11m	51	Sodium selenate, Na_2SeO_4	9m	55
Sodium carbonate sulfate (burkeite), $\text{Na}_6\text{CO}_3(\text{SO}_4)_2$	11m	52	Sodium selenide, Na_2Se	10m	135
Sodium carbonate sulfate, $\text{Na}_6\text{CO}_3(\text{SO}_4)_2$	11m	53	Sodium silicate, α (III), $\text{Na}_2\text{Si}_2\text{O}_5$..	8m	141
Sodium carbonate sulfate, $\text{Na}_6(\text{CO}_3)_2\text{SO}_4$	11m	54	Sodium silicate, $\beta\text{-Na}_2\text{Si}_2\text{O}_5$	10m	136
Sodium chlorate, NaClO_3	3	51	Sodium sulfate, Na_2SO_4	11m	57
Sodium chlorate, NaClO_4 (orthorhombic)	7	49	Sodium sulfate (thenardite), Na_2SO_4	2	59
Sodium chloride (halite), NaCl	2	41	Sodium sulfide, Na_2S	10m	140
Sodium chromium oxide, Na_2CrO_4	9m	48	Sodium sulfite, Na_2SO_3	3	60
Sodium chromium oxide hydrate, $\text{Na}_2\text{CrO}_4 \cdot 4\text{H}_2\text{O}$	9m	50	Sodium telluride, Na_2Te	10m	141
Sodium chromium oxide hydrate, $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot 2\text{H}_2\text{O}$	7m	62	Sodium tin fluoride, NaSn_2F_5	7m	166

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Sodium tungsten oxide, Na_2WO_4	1m	47	Tellurium, Te	1	26
Sodium tungsten(VI) oxide hydrate, $\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$	2m	33	Tellurium(IV) oxide (paratellurite), TeO_2 (tetragonal)	7	56
Sodium zinc fluoride, NaZnF_3	6m	74	Tellurium(IV) oxide, paratellurite, TeO_2 (tetragonal)	10	55
Sodium zinc sulfate hydrate, $\text{Na}_2\text{Zn}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O}$	6m	72	Tellurium(IV) oxide, tellurite, TeO_2 (orthorhombic)	9	57
Sodium zirconium fluoride, $\text{Na}_7\text{Zr}_6\text{F}_{31}$	8m	144	Terbium antimony, TbSb	5m	61
Strontium aluminum hydroxide, $\text{Sr}_3\text{Al}_2(\text{OH})_{12}$	10m	50	Terbium arsenate, TbAsO_4	3m	54
Strontium aluminum oxide, $\text{Sr}_3\text{Al}_2\text{O}_6$	10m	52	Terbium arsenide, TbAs	5m	75
Strontium arsenate, $\text{Sr}_3(\text{AsO}_4)_2$	2m	36	Terbium nitride, TbN	4m	70
Strontium azide, $\text{Sr}(\text{N}_3)_2$	8m	146	Terbium phosphide, TbP	5m	76
Strontium borate, SrB_2O_4	3m	53	Terbium selenide, TbSe	5m	76
Strontium borate, SrB_4O_7	4m	36	Terbium silver, TbAg	5m	74
Strontium bromide fluoride, SrBrF ..	10m	54	Terbium sulfide, TbS	5m	77
Strontium bromide hydrate, $\text{SrBr}_2 \cdot 6\text{H}_2\text{O}$	4	60	Terbium telluride, TbTe	5m	77
Strontium carbonate (strontianite), SrCO_3	3	56	Terbium vanadium oxide, TbVO_4	5m	56
Strontium chloride, SrCl_2	4	40	Thallium aluminum sulfate hydrate, $\text{TlAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$	6	53
Strontium chloride fluoride, SrClF ..	10m	55	Thallium(I) arsenate, Tl_3AsO_4	2m	37
Strontium chloride hydrate, $\text{SrCl}_2 \cdot 2\text{H}_2\text{O}$	11m	58	Thallium azide, TlN_3	8m	82
Strontium chloride hydrate, $\text{SrCl}_2 \cdot 6\text{H}_2\text{O}$	4	58	Thallium(I) bromate, TlBrO_3	8	60
Strontium chlorite hydroxide phosphate, $\text{Sr}_5\text{Cl}_{.65}(\text{OH})_{.35}(\text{PO}_4)_3$..	11m	60	Thallium bromide, TlBr	7	57
Strontium fluoride, SrF_2	5	67	Thallium cadmium sulfate, $\text{Tl}_2\text{Cd}_2(\text{SO}_4)_3$	8m	83
Strontium hydroxide, $\text{Sr}(\text{OH})_2$	13m	41	Thallium(I) chlorate, TlClO_4	2m	38
Strontium hydroxide hydrate, $\text{Sr}(\text{OH})_2 \cdot \text{H}_2\text{O}$	13m	42	Thallium(I) chlorate, TlClO_3	8	61
Strontium hydroxide hydrate, $\text{Sr}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$	13m	43	Thallium(I) chloride, TlCl	4	51
Strontium indium hydroxide, $\text{Sr}_3\text{In}_2(\text{OH})_{12}$	6m	76	Thallium chromium oxide, Tl_2CrO_4 ...	3m	54
Strontium iodide hydrate, $\text{SrI}_2 \cdot 6\text{H}_2\text{O}$	8	58	Thallium chromium sulfate hydrate, $\text{TlCr}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$	6	55
Strontium manganese oxide, SrMnO_3 (cubic)	10m	56	Thallium cobalt sulfate, $\text{Tl}_2\text{Co}_2(\text{SO}_4)_3$	8m	85
Strontium manganese oxide, SrMnO_3 (hexagonal)	10m	58	Thallium cobalt sulfate hydrate, $\text{Tl}_2\text{Co}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$	7m	70
Strontium molybdenum oxide, SrMoO_4 ..	7	50	Thallium copper sulfate hydrate, $\text{Tl}_2\text{Cu}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$	7m	72
Strontium nitrate, $\text{Sr}(\text{NO}_3)_2$	12m	31	Thallium gallium sulfate hydrate, $\text{TlGa}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$	6	57
Strontium oxide, SrO	5	68	Thallium(I) iodate, TlIO_3	8	62
Strontium oxide, SrO_2	6	52	Thallium(I) iodide, TlI (orthorhombic)	4	53
Strontium oxide hydrate, $\text{SrO}_2 \cdot 8\text{H}_2\text{O}$..	11m	61	Thallium iron sulfate hydrate, $\text{Tl}_2\text{Fe}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$	8m	87
Strontium phosphate, $\alpha\text{-Sr}_2\text{P}_2\text{O}_7$	11m	62	Thallium magnesium chromium oxide, $\text{Tl}_2\text{Mg}_2(\text{CrO}_4)_3$	8m	89
Strontium phosphate, $\alpha\text{-Sr}_3(\text{PO}_4)_2$...	11m	64	Thallium magnesium sulfate hydrate, $\text{Tl}_2\text{Mg}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$	7m	74
Strontium scandium oxide hydrate, $\text{Sr}_3\text{Sc}_2\text{O}_6 \cdot 6\text{H}_2\text{O}$	6m	78	Thallium manganese sulfate, $\text{Tl}_2\text{Mn}_2(\text{SO}_4)_3$	7m	76
Strontium silicate, Sr_3SiO_5	13m	44	Thallium nickel sulfate hydrate, $\text{Tl}_2\text{Ni}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$	7m	78
Strontium sulfate (celestite), SrSO_4	2	61	Thallium(I) nitrate, TlNO_3	6	58
Strontium sulfide, SrS	7	52	Thallium(III) oxide, Tl_2O_3	2	28
Strontium telluride, SrTe	4m	69	Thallium(I) phosphate, Tl_3PO_4	7	58
Strontium tin oxide, SrSnO_3	8m	80	Thallium(III) phosphate, TlPO_4	7	59
Strontium titanium oxide, SrTiO_3 ...	3	44	Thallium platinum chloride, Tl_2PtCl_6 ..	5	70
Strontium tungsten oxide, SrWO_4	7	53	Thallium silicon fluoride, Tl_2SiF_6	6	56
Strontium tungsten oxide, Sr_2WO_5 ...	12m	32	Thallium(I) sulfate, Tl_2SO_4	6	59
Strontium zirconium oxide, SrZrO_3 ..	9	51	Thallium(I) thiocyanate, TlCNS	8	63
Sulfamic acid, $\text{H}_2\text{NSO}_3\text{H}$	7	54	Thallium tin chloride, Tl_2SnCl_6	6	54
Sulfur, S (orthorhombic)	9	54	Thallium(I) tungsten oxide, Tl_2WO_4 ..	1m	48
Tantalum, Ta	1	29	Thallium zinc sulfate hydrate, $\text{Tl}_2\text{Zn}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$	7m	80
Tantalum silicide, TaSi_2	8	59	Thorium antimony, ThSb	4m	44

CUMULATIVE INORGANIC INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Thorium arsenide, ThAs	4m	70	Yttrium titanium oxide, Y ₂ TiO ₅	11m	113
Thorium cobalt, Th ₂ Co ₁₇	12m	64	Yttrium vanadium oxide, YVO ₄	5m	59
Thorium iron, Th ₂ Fe ₁₇	12m	67	Zinc, Zn	1	16
Thorium oxide (thorianite), ThO ₂ ...	1	57	Zinc aluminum oxide (gahnite),		
Thulium antimony, TmSb	4m	45	ZnAl ₂ O ₄	2	38
Thulium arsenate, TmAsO ₄	3m	56	Zinc ammine bromide, Zn(NH ₃) ₂ Br ₂ ...	11m	68
Thulium arsenide, TmAs	4m	71	Zinc ammine chloride, Zn(NH ₃) ₂ Cl ₂ ..	10m	59
Thulium nitride, TmN	4m	71	Zinc antimony oxide, ZnSb ₂ O ₄	4m	39
Thulium oxide, Tm ₂ O ₃	9	58	Zinc borate, ZnB ₂ O ₄	1	83
Thulium silver, TmAg	5m	74	Zinc borate, Zn ₄ B ₆ O ₁₃	13m	48
Thulium telluride, TmTe	4m	72	Zinc carbonate, smithsonite, ZnCO ₃	8	69
Thulium vanadium oxide, TmVO ₄	5m	57	Zinc chromium oxide, ZnCr ₂ O ₄	9m	59
Tin, α-Sn (cubic)	2	12	Zinc cobalt oxide, ZnCo ₂ O ₄	10m	60
Tin, β-Sn (tetragonal)	1	24	Zinc cyanide, Zn(CN) ₂	5	73
Tin arsenide, SnAs	4m	37	Zinc fluoride, ZnF ₂	6	60
Tin(II) fluoride, SnF ₂	3m	51	Zinc fluoride hydrate,		
Tin hydrogen phosphate, SnHPO ₄	13m	46	ZnF ₂ ·4H ₂ O	11m	69
Tin(IV) iodide, SnI ₄	5	71	Zinc germanium oxide, Zn ₂ GeO ₄	10	56
Tin(II) oxide, (romarchite), SnO....	4	28	Zinc hydroxide silicate hydrate,		
Tin(IV) oxide (cassiterite), SnO ₂ ..	1	54	hemimorphite, Zn ₄ (OH) ₂ Si ₂ O ₇ ·H ₂ O ...	2	62
Tin sulfide (berndtite), β-SnS ₂	9m	57	Zinc iodide, ZnI ₂	9	60
Tin(II) telluride, SnTe	7	61	Zinc iron oxide (franklinite),		
Titanium, Ti	3	1	ZnFe ₂ O ₄	9m	60
Titanium oxide (anatase), TiO ₂	7m	82	Zinc manganese oxide (hetaerolite),		
Titanium oxide, brookite, TiO ₂			ZnMn ₂ O ₄	10m	61
(orthorhombic)	3m	57	Zinc molybdenum oxide, Zn ₂ Mo ₃ O ₈	7m	173
Titanium oxide (rutile), TiO ₂	7m	83	Zinc nitrate hydrate,		
Titanium(III) oxide, TiO _{1.515}	9	59	α-Zn(NO ₃) ₂ ·6H ₂ O	12m	36
Titanium silicide, Ti ₅ Si ₃	8	64	Zinc oxide (zincite), ZnO	2	25
Titanium sulfide, TiS ₂	4m	72	Zinc selenide, ZnSe	3	23
Titanium sulfide, Ti ₂ S	8m	149	Zinc silicate (willemite), Zn ₂ SiO ₄	7	62
Tungsten, W	1	28	Zinc silicon fluoride hydrate,		
Tungsten, W (reference standard) ...	8m	2	ZnSiF ₆ ·6H ₂ O	8	70
Tungsten sulfide (tungstenite), WS ₂	8	65	Zinc sulfate (zinkosite), ZnSO ₄	7	64
Uranium oxide, UO	5m	78	Zinc sulfate hydrate (goslarite),		
Uranium oxide (uraninite), UO ₂	2	33	ZnSO ₄ ·7H ₂ O	8	71
Uranium selenide, USe	5m	78	Zinc sulfide (wurtzite), α-ZnS		
Uranium telluride, UTe	4m	73	(hexagonal)	2	14
Vanadium, V	9m	58	Zinc sulfide (sphaelerite), β-ZnS		
Vanadium gold 3:1, V ₃ Au	6m	18	(cubic)	2	16
Vanadium iridium 3:1, V ₃ Ir	6m	21	Zinc telluride, ZnTe	3m	58
Vanadium(V) oxide, V ₂ O ₅	8	66	Zinc tin oxide, Zn ₂ SnO ₄	10m	62
Vanadium palladium 3:1, V ₃ Pd	6m	32	Zinc titanium oxide, ZnTiO ₃	13m	49
Vanadium platinum 3:1, V ₃ Pt	6m	34	Zinc titanium oxide, Zn ₂ TiO ₄	12m	37
Vanadium rhodium 3:1, V ₃ Rh	6m	56	Zinc tungsten oxide (sanmartinite),		
Ytterbium antimony, YbSb	4m	45	ZnWO ₄	2m	40
Ytterbium arsenate, YbAsO ₄	4m	38	Zirconium, α-Zr	2	11
Ytterbium arsenide, YbAs	4m	73	Zirconium hydride, ZrH ₂	5m	60
Ytterbium gallium oxide, Yb ₃ Ga ₅ O ₁₂	1m	49	Zirconium iodate, Zr(IO ₃) ₄	1m	51
Ytterbium nitride, YbN	4m	74	Zirconium nitride, ZrN	5m	80
Ytterbium oxide, Yb ₂ O ₃	6m	80	Zirconium oxide, ZrO	5m	81
Ytterbium selenide, YbSe	5m	79	Zirconium phosphide, ZrP	4m	75
Ytterbium telluride, YbTe	5m	79	Zirconium silicate, zircon, ZrSiO ₄	4	68
Ytterbium(III) vanadium oxide, YbVO ₄	5m	58	Zirconium sulfate hydrate		
Yttrium antimony, YSb	4m	46	(zircosulfate), Zr(SO ₄) ₂ ·4H ₂ O	7	66
Yttrium arsenate, YAsO ₄	2m	39			
Yttrium arsenide, YAs	4m	74			
Yttrium gallium oxide, Y ₃ Ga ₅ O ₁₂	1m	50			
Yttrium nickel, YNi ₃	10m	123			
Yttrium oxide, Y ₂ O ₃	3	28			
Yttrium oxychloride, YOCl	1m	51			
Yttrium phosphate (xenotime), YPO ₄	8	67			
Yttrium silver, YAg	5m	75			
Yttrium sulfide, YS	5m	80			
Yttrium telluride, YTe	4m	75			

CUMULATIVE ORGANIC INDEX

	Vol. or Sec.	Page		Vol. or Sec.	Page
4-Acetyl-2'-fluorodiphenyl, $C_{14}H_{11}FO$	8m	91	Nickel acetate hydrate, $Ni(C_2H_3O_2)_2 \cdot 4H_2O$	13m	31
Alanine, L, $CH_3CHNH_2CO_2H$	8m	93	Nickel hexaimidazole nitrate, $Ni(C_3H_4N_2)_6(NO_3)_2$	7m	27
Ammonium acetate, $NH_4 \cdot CH_3CO_2$	8m	95	Nickel tetrapyrazole chloride, $Ni(C_3H_4N_2)_4Cl_2$	8m	44
Ammonium formate, NH_4HCO_2	11m	9	Octahydro-1,3,5,7-tetranitro- 1,3,5,7-tetrazocine (α -HMX), $C_4H_8N_8O_8$	11m	100
Ammonium oxalate hydrate (oxammite), $(NH_4)_2C_2O_4 \cdot H_2O$	7	5	Octahydro-1,3,5,7-tetranitro- 1,3,5,7-tetrazocine (β -HMX), $C_4H_8N_8O_8$	11m	102
Ammonium yttrium oxalate hydrate, $NH_4Y(C_2O_4)_2 \cdot H_2O$	8m	97	Palladium bis-(N-isopropyl-3-ethyl- salicylaldiminate), $Pd(C_{12}H_{16}NO)_2$	7m	144
Ascorbic acid, L- $C_6H_8O_6$	8m	99	Pimelic acid, $(CH_2)_5(COOH)_2$	7m	153
Azobenzene, $C_6H_5NNC_6H_5$	7m	86	Potassium formate-formic acid complex, $KO_2CH \cdot HO_2CH$	9m	93
Cadmium hexaimidazole nitrate, $Cd(C_3H_4N_2)_6(NO_3)_2$	8m	23	Potassium hydrogen o-phthalate, $C_6H_4(COOH)(COOK)$	4m	30
Calcium formate, $Ca(HCO_2)_2$	8	16	Potassium oxalate hydrate, $K_2C_2O_4 \cdot H_2O$	9m	39
Calcium malate hydrate, $Ca(O_2C)_2(CH_2CHOH) \cdot 2H_2O$	10m	76	Potassium oxalate perhydrate, $K_2C_2O_4 \cdot H_2O_2$	9m	96
Cobalt acetate hydrate, $Co(C_2H_3O_2)_2 \cdot 4H_2O$	12m	19	Reserpine, $C_{33}H_{40}N_2O_9$	8m	123
Copper glutamate hydrate, $Cu(O_2C)_2(H_2NCHCH_2CH_2) \cdot 2H_2O$	7m	110	Rubidium oxalate perhydrate, $Rb_2C_2O_4 \cdot H_2O_2$	9m	102
Copper imidazole nitrate, $Cu(C_3H_4N_2)_4(NO_3)_2$	13m	24	Silver oxalate, $Ag_2C_2O_4$	9m	47
Copper tetrapyrazole chloride, $Cu(C_3H_4N_2)_4Cl_2$	8m	31	Sodium D-tartrate hydrate, $(CHOH-CO_2Na)_2 \cdot 2H_2O$	11m	110
Cysteine, L, $HSCH_2 \cdot CH(NH_2) \cdot COOH$	11m	86	Sodium oxalate, $Na_2C_2O_4$	6m	70
Dibenzoylmethane, $(C_6H_5CO)_2CH_2$	7m	115	Strontium formate, $Sr(CHO_2)_2$	8	55
bis-(o-Dodecacarborane), $C_{40}B_{20}H_{22}$..	6m	7	Strontium formate hydrate, $Sr(CHO_2)_2 \cdot 2H_2O$ (orthorhombic)	8	56
Glucose, D, α , (dextrose), $C_6H_{12}O_6$..	11m	28	Sucrose, $C_{12}H_{22}O_{11}$	11m	66
Glyoxime, $H_2C_2(NO_2)_2$	8m	102	Tartaric acid, D, $(CHOHCO_2H)_2$	7m	168
Hexamethylenediammonium adipate, $(CH_2)_4(CO_2H_3N)_2(CH_2)_6$	7m	121	Trimethylammonium chloride, $(CH_3)_3NHCl$	9m	113
Holmium ethylsulfate hydrate, $Ho[(C_2H_5)SO_4]_3 \cdot 9H_2O$	1m	18	2,4,6-Trinitrophenetole, $C_2H_5OC_6H_2(NO_2)_3$	8m	152
Hydroquinone, γ - HOC_6H_4OH	8m	107	Urea, $CO(NH_2)_2$	7	61
Iron oxalate hydrate (humboldtine), $FeC_2O_4 \cdot 2H_2O$	10m	24	Uric acid, $C_5H_4N_4O_3$	8m	154
Lead formate, $Pb(HCO_2)_2$	8	30	Zinc diimidazole chloride, $Zn(C_3H_4N_2)_2Cl_2$	7m	123
Lithium oxalate, $Li_2C_2O_4$	10m	34	Zinc glutamate hydrate, $Zn(O_2CCHNH_2CH_2CH_2CO_2) \cdot 2H_2O$	7m	170
Mercury o-phthalate, $C_6H_4(CO_2Hg)_2$..	10m	113			
Methyl sulfonanilide, $C_6H_5NHSO_2CH_3$	9m	78			
N-Methylphenazinium-7,7,8,8-tetra- cyanoquinodimethanide, $C_{25}H_{15}N_6$...	7m	146			
2-Naphthylamine, N-phenyl-, $C_{10}H_7NHC_6H_5$	6m	29			
Neodymium ethylsulfate hydrate, $Nd[(C_2H_5)SO_4]_3 \cdot 9H_2O$	9	41			

CUMULATIVE MINERAL INDEX

	Vol. or Sec.	Page		Vol. or Sec.	Page
Acanthite, Ag_2S (monoclinic).....	10	51	Cryptohalite, $(\text{NH}_4)_2\text{SiF}_6$	5	5
Aeschnite CeNbTiO_6	3m	24	Cuprite, Cu_2O	2	23
Alabandite, MnS	4	11	*Diamond, C	2	5
Alum, $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$	6	36	*Diaspore, $\text{Al}_2\text{O}_3 \cdot \text{H}_2\text{O}$	3	41
Anatase, TiO_2	7m	82	Diopside, $\text{CaMg}(\text{SiO}_3)_2$	5m	17
Andradite, $\text{Ca}_3\text{Fe}_2\text{Si}_3\text{O}_{12}$	9	22	*Dravite, $\text{NaMg}_3\text{Al}_6\text{B}_3\text{Si}_6\text{O}_{27}(\text{OH})_4$	3m	47
Anglesite, PbSO_4	3	67	Eitelite, $\text{Na}_2\text{Mg}(\text{CO}_3)_2$	11m	56
Anhydrite, CaSO_4	4	65	Elpasolite, K_2NaAlF_6	9m	43
Antarcticite, $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$	12m	16	*Enstatite, MgSiO_3	6	32
Antimony, Sb.....	3	14	Epsomite, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	7	30
Aphthitalite, $\text{K}_3\text{Na}(\text{SO}_4)_2$	6m	52	Eskolaite, Cr_2O_3	5	22
Aragonite, CaCO_3	3	53	Ettringite, $\text{Ca}_6\text{Al}_2\text{S}_3\text{O}_{18} \cdot 31\text{H}_2\text{O}$	8	3
Arcanite, K_2SO_4	3	62	Fairchildite, $\text{K}_2\text{Ca}(\text{CO}_3)_2$	8m	48
Arsenic, As	3	6	Fluorapatite, $\text{Ca}_5\text{F}(\text{PO}_4)_3$	3m	22
Arsenolite, As_2O_3	1	51	Fluorite, CaF_2	1	69
Aurostibite, AuSb_2	7	18	Forsterite, Mg_2SiO_4	1	83
*Azurite, $\text{Cu}_3(\text{OH})_2(\text{CO}_3)_2$	10	30	Franklinite, ZnFe_2O_4	9m	60
Baryte, BaSO_4	10m	12	Fresnoite, $\text{Ba}_2\text{TiSi}_2\text{O}_8$	9m	14
Berlinite, AlPO_4	10	3	Gahnite, ZnAl_2O_4	2	38
Berndtite, SnS_2	9m	57	Galaxite, MnAl_2O_4	9	35
*Beryl, $\text{Be}_3\text{Al}_2\text{Si}_6\text{O}_{18}$	9	13	Galena, PbS	2	18
Bischofite, $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$	11m	37	Geikielite, MgTiO_3	5	43
Bismite, $\alpha\text{-Bi}_2\text{O}_3$	3m	17	Gersdorffite, NiAsS	1m	35
Bismoclite, BiOCl	4	54	Glauberite, $\text{Na}_2\text{Ca}(\text{SO}_4)_2$	6m	59
Bismuth, Bi	3	20	Gold, Au	1	33
Bismuthinite, Bi_2S_3	5m	13	Goslarite, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	8	71
Bixbyite, $\alpha\text{-Mn}_2\text{O}_3$	11m	95	Greenockite, CdS	4	15
*Bloedite, $\text{Na}_2\text{Mg}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O}$	6m	63	*Groutite, $\text{MnO}(\text{OH})$	11m	97
Boehmite, $\text{Al}_2\text{O}_3 \cdot \text{H}_2\text{O}$	3	38	Halite, NaCl	2	41
Bromargyrite, AgBr	4	46	Hausmannite, Mn_3O_4	10m	38
Bromellite, BeO	1	36	*Hemimorphite, $\text{Zn}_4(\text{OH})_2\text{Si}_2\text{O}_7 \cdot \text{H}_2\text{O}$...	2	62
*Brookite, TiO_2	3m	57	Hetaerolite, ZnMn_2O_4	10m	61
Brucite, $\text{Mg}(\text{OH})_2$	6	30	Hieratite, K_2SiF_6	5	50
Bunsenite, NiO	1	47	Huebnerite, MnWO_4	2m	24
Burkeite, $\text{Na}_6\text{CO}_3(\text{SO}_4)_2$	11m	52	Humboldtine, $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$	10m	24
*Butlerite, $\text{Fe}(\text{OH})\text{SO}_4 \cdot 2\text{H}_2\text{O}$	10m	95	Humite, $3\text{Mg}_2\text{SiO}_4 \cdot \text{MgF}_2$	1m	30
Cadmoselite, CdSe	7	12	Hydrophilite, CaCl_2	11m	18
Calcite, CaCO_3	2	51	Indialite, $\text{Mg}_2\text{Al}_4\text{Si}_5\text{O}_{18}$ (hexagonal)	1m	29
Calomel, Hg_2Cl_2	13m	30	Iodargyrite, AgI	8	51
Carnallite, $\text{KMgCl}_3 \cdot 6\text{H}_2\text{O}$	8m	50	Iron, $\alpha\text{-Fe}$	4	3
Cassiterite, SnO_2	1	54	Jacobsite, MnFe_2O_4	9	36
Celestite, SrSO_4	2	61	*Julgoidite, $\text{Ca}_2\text{Fe}_3\text{Si}_3\text{O}_{10}(\text{OH},\text{O})_2(\text{OH})_2$	10m	72
Cerianite, CeO_2	1	56	Langbeinite, $\text{K}_2\text{Mg}_2(\text{SO}_4)_3$	6m	40
Cerussite, PbCO_3	2	56	Lead, Pb	1	34
Cervantite, Sb_2O_4	10	8	*Leucophanite, $\text{NaCaBeFSi}_2\text{O}_6$	8m	138
Chalcocyanite, CuSO_4	3m	29	Litharge, PbO (red)	2	30
Chloraluminite, $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$	7	3	Lithiophosphate, Li_3PO_4	4m	21
Chlorargyrite, AgCl	4	44	Loellingite, FeAs_2	10	34
Chloromagnesite, MgCl_2	11m	94	Macedonite, PbTiO_3	5	39
Chromatite, CaCrO_4	7	13	Magnesiochromite, MgCr_2O_4	9	34
Chrysoberyl, BeAl_2O_4	9	10	Magnesite, MgCO_3	7	28
Cinnabar, HgS	4	17	Magnetite, Fe_3O_4	5m	31
*Claudetite, As_2O_3	3m	9	Malachite, $\text{Cu}_2(\text{OH})_2\text{CO}_3$	10	31
Clausthalite, PbSe	5	38	Manganolangbeinite, $\text{K}_2\text{Mn}_2(\text{SO}_4)_3$	6m	43
Copper, Cu	1	15	Manganosite, MnO	5	45
Cordierite, $\text{Mg}_2\text{Al}_4\text{Si}_5\text{O}_{18}$ (orthorhombic)	1m	28	Marshite, CuI	4	38
Corundum, Al_2O_3	9	3	Mascagnite, $(\text{NH}_4)_2\text{SO}_4$	9	8
Cotunnite, PbCl_2	12m	23	Massicot, PbO (yellow)	2	32
Covellite, CuS	4	13	Matlockite, PbFCl	13m	25
Cristobalite (α or low) SiO_2	10	48	Mayenite, $\text{Ca}_{12}\text{Al}_{14}\text{O}_{33}$	9	20
Cristobalite (β or high) SiO_2	1	42	Melanterite, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	8m	38
*Cryolithionite, $\text{Li}_3\text{Na}_3\text{Al}_2\text{F}_{12}$	9m	23	*Meliphanite, $\text{Na}_{.63}\text{Ca}_{1.37}\text{BeAl}_{.13}\text{Si}_{1.87}\text{O}_{6.25}\text{F}_{.75}$..	8m	135
			Metaborite, HBO_2 (cubic)	4m	27
			Metacinnabar, HgS	4	21
			Miargyrite, AgSbS_2	5m	49
			*Millerite, NiS	1m	37

*Natural mineral.

CUMULATIVE MINERAL INDEX - Continued

	Vol. or Sec.	Page		Vol. or Sec.	Page
Minium, Pb_3O_4	8	32	Silver, Ag (reference standard)	8m	2
Mitscherlichite, $K_2CuCl_4 \cdot 2H_2O$	9m	34	*Sjögrenite, $Mg_6Fe_2CO_3(OH)_{16} \cdot 4H_2O$, phase I	10m	103
Molybdenite, MoS_2	5	47	Skutterudite, $CoAs_3$	10	21
Molybdate, MoO_3	3	30	*Smithsonite, $ZnCO_3$	8	69
Montroydite, HgO	9	39	*Sodalite, $Na_8Si_6Al_6O_{24}Cl_2$	7m	158
Mullite, $Al_6Si_2O_{13}$	3m	3	Soda-niter, $NaNO_3$	6	50
Nantokite, $CuCl$	4	35	Sphaerocobaltite, $CoCO_3$	10	24
*Newberyite, $MgHPO_4 \cdot 3H_2O$	7m	139	Sphalerite, ZnS	2	16
Niter, KNO_3	3	58	Spinel, $MgAl_2O_4$	9m	25
Nitrammite, NH_4NO_3	7	4	Stibnite, Sb_2S_3	5	6
Nitrobarite, $Ba(NO_3)_2$	11m	14	Stolzite, $PbWO_4$	5m	34
Norbergite, $Mg_2SiO_4 \cdot MgF_2$	10	39	Strontianite, $SrCO_3$	3	56
Oldhamite, CaS	7	15	Struvite, $MgNH_4PO_4 \cdot 6H_2O$	3m	41
Otavite, $CdCO_3$	7	11	Sulfur, S (orthorhombic)	9	54
Oxammite, $(NH_4)_2C_2O_4 \cdot H_2O$	7	5	Sylvite, KCl	1	65
Palladium, Pd	1	21	*Tellurite, TeO_2	9	57
*Paratellurite, TeO_2	10	55	Tellurium, Te	1	26
Paratellurite, TeO_2	7	56	Tellurobismuthite, Bi_2Te_3	3m	16
Periclase, MgO	1	37	Tenorite, CuO	1	49
Perovskite, $CaTiO_3$	9m	17	Teschemacherite, NH_4HCO_3	9	5
*Phenakite, Be_2SiO_4	8	11	Thenardite, Na_2SO_4	2	59
Picromerite, $K_2Mg(SO_4)_2 \cdot 6H_2O$	8m	54	Thermonatrite, $Na_2CO_3 \cdot H_2O$	8	54
*Pirssonite, $Na_2Ca(CO_3)_2 \cdot 2H_2O$	9m	106	*Thomsenolite, $NaCaAlF_6 \cdot H_2O$	8m	132
Platinum, Pt	1	31	Thorianite, ThO_2	1	57
Portlandite, $Ca(OH)_2$	1	58	Thortveitite, $Sc_2Si_2O_7$	7m	58
Powellite, $CaMoO_4$	6	22	Tiemannite, $HgSe$	7	35
Pyrargyrite, Ag_3SbS_3	5m	51	Tin, α -Sn (cubic)	2	12
Pyrite, FeS_2	5	29	Tin, β -Sn (tetragonal)	1	24
*Pyroaurite, $Mg_6Fe_2CO_3(OH)_{16} \cdot 4H_2O$, phase II	10m	104	*Topaz, $Al_2SiO_4(F,OH)_2$	1m	4
Pyrolusite, β - MnO_2	10m	39	Trevorite, $NiFe_2O_4$	10	44
Pyrope, $Mg_3Al_2(SiO_4)_3$	4m	24	Tschermigite, $NH_4Al(SO_4)_2 \cdot 12H_2O$	6	3
*Quartz, SiO_2 (α or low)	3	24	Tungstenite, WS_2	8	65
Rammelsbergite, $NiAs_2$	10	42	Uraninite, UO_2	2	33
Retgersite, $NiSO_4 \cdot 6H_2O$	7	36	Uvarovite, $Ca_3Cr_2(SiO_4)_3$	10	17
Rhodochrosite, $MnCO_3$	7	32	*Valentinite, Sb_2O_3	10	6
Romarchite, SnO	4	28	Valentinite, Sb_2O_3	10	6
Rutile, TiO_2	7m	83	Villiaumite, NaF	1	63
Safflorite, $CoFeAs_4$	10	28	Willemite, Zn_2SiO_4	7	62
Sal-ammoniac, NH_4Cl	1	59	Witherite, $BaCO_3$	2	54
Sanbornite, β - $BaSi_2O_5$	13m	10	Wulfenite, $PbMoO_4$	7	23
Sanmartinite, $ZnWO_4$	2m	40	Wurtzite, ZnS	2	14
Scacchite, $MnCl_2$	8m	43	*Xanthoconite, Ag_3AsS_3	8m	126
*Scheelite, $CaWO_4$	6	23	Xenotime, YPO_4	8	67
Selenium, Se	5	54	Zinc, Zn	1	16
Selenolite, SeO_2	7m	60	Zincite, ZnO	2	25
Sellaite, MgF_2	4	33	Zinkosite, $ZnSO_4$	7	64
Senarmontite, Sb_2O_3	3	31	*Zircon, $ZrSiO_4$	4	68
Shcherbinaite, V_2O_5	8	66	Zircosulfate, $Zr(SO_4)_2 \cdot 4H_2O$	7	66
Silver, Ag	1	23			

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET		1. PUBLICATION OR REPORT NO. NBS-MN25, Section 13	2. Gov't Accession No.	3. Recipient's Accession No.
4. TITLE AND SUBTITLE Standard X-ray Diffraction Powder Patterns Section 13 - Data for 58 Substances			5. Publication Date June 1976	
			6. Performing Organization Code	
7. AUTHOR(S) Marlene C. Morris, Howard F. McMurdie, Eloise H. Evans Boris Paretzkin, Johan H. de Groot, Camden R. Hubbard and Simon J. Carmel			8. Performing Organ. Report No. J. Carmel	
9. PERFORMING ORGANIZATION NAME AND ADDRESS NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, D.C. 20234			10. Project/Task/Work Unit No. 3130163	
			11. Contract/Grant No.	
12. Sponsoring Organization Name and Complete Address (Street, City, State, ZIP) Same as Item 9.			13. Type of Report & Period Covered Interim	
			14. Sponsoring Agency Code	
15. SUPPLEMENTARY NOTES Library of Congress Catalog Card Number: 53-61386				
16. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here.) Standard x-ray diffraction patterns are presented for 58 substances. Thirty-one of these patterns represent experimental data and 27 are calculated. The experimental x-ray powder diffraction patterns were obtained with an x-ray diffractometer. All d-values were assigned Miller indices determined by comparison with computed inter-planar spacings consistent with space group extinctions. The densities and lattice constants were calculated, and the refractive indices were measured whenever possible. The calculated x-ray powder diffraction patterns were computed from published crystal structure data. Both peak height and integrated intensities are reported for the calculated patterns.				
17. KEY WORDS (six to twelve entries; alphabetical order; capitalize only the first letter of the first key word unless a proper name; separated by semicolons) Crystal structure; integrated intensities; lattice constants; peak intensities; powder patterns; reference intensities; standard; x-ray diffraction.				
18. AVAILABILITY <input checked="" type="checkbox"/> Unlimited <input type="checkbox"/> For Official Distribution. Do Not Release to NTIS <input checked="" type="checkbox"/> Order From Sup. of Doc., U.S. Government Printing Office Washington, D.C. 20402, SD Cat. No. C13, 44:25/Sec. 13 <input type="checkbox"/> Order From National Technical Information Service (NTIS) Springfield, Virginia 22151		19. SECURITY CLASS (THIS REPORT) UNCLASSIFIED		21. NO. OF PAGES 114
		20. SECURITY CLASS (THIS PAGE) UNCLASSIFIED		22. Price \$1.80

SINGLE CRYSTAL DATA

REVISED! UPDATED!


In 1954, the first edition of CRYSTAL DATA (Determinative Tables and Systematic Tables) was published as Memoir 60 of the Geological Society of America. In 1960, the second edition of the Determinative Tables was issued as Monograph 5 of the American Crystallographic Association, and in 1967, the Systematic Tables were issued as Monograph 6. These editions proved extremely valuable to crystallographers throughout the world. Recognizing the need for updated crystallographic information, the National Bureau of Standards Office of Standard Reference Data has sponsored the issuance of a new edition.

This, the THIRD EDITION, should be of particular interest not only to crystallographers but also to chemists, mineralogists, physicists and individuals in related fields of study. The current edition, which comprises two volumes, Organic and Inorganic, is a thoroughly revised and updated work, containing over 25,000 entries.

The entries are listed, *within each crystal system*, according to increasing values of a determinative number: a/b ratio in trimetric systems, c/a ratio in dimetric systems, and cubic cell edge a , in the isometric system. In addition, the following information is given:

axial ratio(s) and interaxial angles not fixed by symmetry, cell dimensions, space group or diffraction aspect, number of formula units per unit cell, crystal structure, (whether determined), measured density and x-ray calculated density. Also listed is the name of the compound and synonym(s), chemical formula, literature reference and transformation matrix. When available, the crystal structure type, crystal habit, cleavages, twinning, color, optical properties, indices of refraction, optical orientation, melting point and transition point are also listed.

THIS EDITION culminates years of effort by J. D. H. Donnay, Johns Hopkins University, Helen M. Ondik, National Bureau of Standards, Sten Samson, California Institute of Technology, Quintin Johnson, Lawrence Radiation Laboratory, Melvin H. Mueller, Argonne National Laboratory, Gerard M. Wolten, Aerospace Corporation, Mary E. Mrose, U.S. Geological Survey, Olga Kennard and David G. Watson, Cambridge University, England and Murray Vernon King, Massachusetts General Hospital.



INORGANIC VOLUME \$50.00

ORGANIC VOLUME \$30.00

Plus shipping and handling

Shipments are made via insured parcel post. Additional charges for shipments by air or commercial carrier.

TERMS: Domestic—30 days Foreign—prepayment required. Address all orders to:

JOINT COMMITTEE ON POWDER DIFFRACTION STANDARDS 1601 Park Lane, Swarthmore, Pennsylvania 19081

Please accept my order for CRYSTAL DATA, DETERMINATIVE TABLES, Third Edition, Donnay/Ondik.

☐ Organic Volume

☐ Inorganic Volume

Ship to: _____

Signature _____



JCPDS

PERIODICALS

JOURNAL OF RESEARCH reports National Bureau of Standards research and development in physics, mathematics, and chemistry. It is published in two sections, available separately:

• **Physics and Chemistry (Section A)**

Papers of interest primarily to scientists working in these fields. This section covers a broad range of physical and chemical research, with major emphasis on standards of physical measurement, fundamental constants, and properties of matter. Issued six times a year. Annual subscription: Domestic, \$17.00; Foreign, \$21.25.

• **Mathematical Sciences (Section B)**

Studies and compilations designed mainly for the mathematician and theoretical physicist. Topics in mathematical statistics, theory of experiment design, numerical analysis, theoretical physics and chemistry, logical design and programming of computers and computer systems. Short numerical tables. Issued quarterly. Annual subscription: Domestic, \$9.00; Foreign, \$11.25.

DIMENSIONS/NBS (formerly *Technical News Bulletin*)—This monthly magazine is published to inform scientists, engineers, businessmen, industry, teachers, students, and consumers of the latest advances in science and technology, with primary emphasis on the work at NBS. The magazine highlights and reviews such issues as energy research, fire protection, building technology, metric conversion, pollution abatement, health and safety, and consumer product performance. In addition, it reports the results of Bureau programs in measurement standards and techniques, properties of matter and materials, engineering standards and services, instrumentation, and automatic data processing.

Annual subscription: Domestic, \$9.45; Foreign, \$11.85.

NONPERIODICALS

Monographs—Major contributions to the technical literature on various subjects related to the Bureau's scientific and technical activities.

Handbooks—Recommended codes of engineering and industrial practice (including safety codes) developed in cooperation with interested industries, professional organizations, and regulatory bodies.

Special Publications—Include proceedings of conferences sponsored by NBS, NBS annual reports, and other special publications appropriate to this grouping such as wall charts, pocket cards, and bibliographies.

Applied Mathematics Series—Mathematical tables, manuals, and studies of special interest to physicists, engineers, chemists, biologists, mathematicians, computer programmers, and others engaged in scientific and technical work.

National Standard Reference Data Series—Provides quantitative data on the physical and chemical properties of materials, compiled from the world's literature and critically evaluated. Developed under a world-wide

program coordinated by NBS. Program under authority of National Standard Data Act (Public Law 90-396).

NOTE: At present the principal publication outlet for these data is the *Journal of Physical and Chemical Reference Data* (JPCRD) published quarterly for NBS by the American Chemical Society (ACS) and the American Institute of Physics (AIP). Subscriptions, reprints, and supplements available from ACS, 1155 Sixteenth St. N. W., Wash. D. C. 20056.

Building Science Series—Disseminates technical information developed at the Bureau on building materials, components, systems, and whole structures. The series presents research results, test methods, and performance criteria related to the structural and environmental functions and the durability and safety characteristics of building elements and systems.

Technical Notes—Studies or reports which are complete in themselves but restrictive in their treatment of a subject. Analogous to monographs but not so comprehensive in scope or definitive in treatment of the subject area. Often serve as a vehicle for final reports of work performed at NBS under the sponsorship of other government agencies.

Voluntary Product Standards—Developed under procedures published by the Department of Commerce in Part 10, Title 15, of the Code of Federal Regulations. The purpose of the standards is to establish nationally recognized requirements for products, and to provide all concerned interests with a basis for common understanding of the characteristics of the products. NBS administers this program as a supplement to the activities of the private sector standardizing organizations.

Federal Information Processing Standards Publications (FIPS PUBS)—Publications in this series collectively constitute the Federal Information Processing Standards Register. Register serves as the official source of information in the Federal Government regarding standards issued by NBS pursuant to the Federal Property and Administrative Services Act of 1949 as amended, Public Law 89-306 (79 Stat. 1127), and as implemented by Executive Order 11717 (38 FR 12315, dated May 11, 1973) and Part 6 of Title 15 CFR (Code of Federal Regulations).

Consumer Information Series—Practical information, based on NBS research and experience, covering areas of interest to the consumer. Easily understandable language and illustrations provide useful background knowledge for shopping in today's technological marketplace.

NBS Interagency Reports (NBSIR)—A special series of interim or final reports on work performed by NBS for outside sponsors (both government and non-government). In general, initial distribution is handled by the sponsor; public distribution is by the National Technical Information Service (Springfield, Va. 22161) in paper copy or microfiche form.

Order NBS publications (except NBSIR's and Bibliographic Subscription Services) from: Superintendent of Documents, Government Printing Office, Washington, D.C. 20402.

BIBLIOGRAPHIC SUBSCRIPTION SERVICES

The following current-awareness and literature-survey bibliographies are issued periodically by the Bureau:

Cryogenic Data Center Current Awareness Service

A literature survey issued biweekly. Annual subscription: Domestic, \$20.00; foreign, \$25.00.

Liquefied Natural Gas. A literature survey issued quarterly. Annual subscription: \$20.00.

Superconducting Devices and Materials. A literature

survey issued quarterly. Annual subscription: \$20.00. Send subscription orders and remittances for the preceding bibliographic services to National Bureau of Standards, Cryogenic Data Center (275.02) Boulder, Colorado 80302.

Electromagnetic Metrology Current Awareness Service Issued monthly. Annual subscription: \$24.00. Send subscription order and remittance to Electromagnetics Division, National Bureau of Standards, Boulder, Colo. 80302.

U.S. DEPARTMENT OF COMMERCE
National Bureau of Standards
Washington, D.C. 20234

OFFICIAL BUSINESS

Penalty for Private Use, \$300

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-215



SPECIAL FOURTH-CLASS RATE
BOOK



75 YEARS
NBS
1901-1976